This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.

+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.

+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.

+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`

NEDL TRANSFER
HN 3FVR U

KF3905



A

TREATISE

ON THE

DIFFERENTIAL AND INTEGRAL CALCULUS,

AND THE

CALCULUS OF VARIATIONS.

BY THE

REV. T. G. HALL, M.A.

PROFESSOR OF MATHEMATICS IN KING'S COLLEGE, LONDON;
PREBENDARY OF ST. PAUL'S;
AND LATE
FELLOW AND TUTOR OF MAGDALENE COLLEGE, CAMBRIDGE.

THE FOURTH EDITION.

LONDON:
JOHN W. PARKER, WEST STRAND.

M.DCCC.XLVI.

KF 3905

HARVARD
UNIVERSITY
LIBRARY

CAMBRIDGE:

PRINTED AT THE UNIVERSITY PRESS.

PREFACE.

A KNOWLEDGE of the principles of this branch of the Pure Mathematics is absolutely necessary, before any one can successfully undertake the perusal of works on Natural Philosophy, in which the effects of the observed laws that govern the material world are reduced to calculation.

For Students deficient in this knowledge, yet anxious to obtain as much as may enable them to master the chief analytical difficulties incident to the study of Elementary Treatises on the Mixed Mathematics, this book has been written: and with the hope that by its means, a subject of high interest may be rendered accessible to an increased number of readers.

The ample Table of Contents which accompanies this work will sufficiently exhibit its plan, which in its arrangement differs in part from that of former editions. The Theory of the Functions of two Variables and its application to Maxima and Minima are earlier introduced—and are placed before those chapters which treat of Curves and their properties. To this, the Fourth Edition, many additional Problems have been given; every chapter prior to its going to the press has been attentively examined, and in almost all some improvement has been made, and to most some new article added, illustrative of the subject and the result of the long experience of the author.

A few words may be here added, in explanation of the principles adopted in laying down the definitions.

By a method, similar to that of M. Poisson, it is shewn that $u_1 = f(x+h)$ can always be expanded in a series of ascending integral powers of h, which may be written under the convenient form of the equation

$$u_1 = u + Ah + Uh^2.$$

The term Ah, the first term of the difference between $u_1 - u$, is defined to be the differential of u: and A the coefficient of h is called the differential coefficient: and from these definitions, the rules for Differentiation are in general derived.

But since when the general form of $f(x+h)$ has been demonstrated, we see that $\frac{u_1 - u}{h}$ is equal to A, when $h = 0$: we may therefore find the differential coefficient, by dividing both sides of the equation $u_1 - u = \phi(x, h)$ (not expanded), by h, and then make $h = 0$. This method which sometimes diminishes the algebraical labour of finding A, is in few instances made use of. It is in fact the method of Limits, often useful in the application of the Calculus, and to which the idea of series is so necessary an auxiliary.

In truth the notion of a series seems inseparably connected with the method of Limits—to which in the Differential Calculus it gives a clearness and precision, of which it stands much in need. For to say that when $u = f(x)$; and u_1 is its value when x becomes x_1 the differential Coefficient is the value to which $\frac{u_1 - u}{x_1 - x}$ tends while x_1 continually approaches x, without first exhibiting the relation which exists between u_1 and u, is to use a mysterious obscurity which must heavily tax the faith or the credulousness of the reader. But put for u_1 its expansion and the difficulty vanishes.

It has been said that $f(x+h)$ produces diverging series—if by this it is meant, that Ah is improperly determined, the objection lies equally against the method of Limits; and although a preface is not the place to enter into such discussions, it may be remarked that

it is one thing to have a diverging series, arising from a known function, and another to determine by a limited number of terms the value of such a series.

If however it be granted that the method of Limits, and that used in our definitions, will when u is a function of one variable only, lead with equal facility to the same results; this is not the case, with the former method when u is a function of two or a greater number of variables—either some new definition, or some new hypothesis is necessary, in order to arrive at the equation $du = \left(\frac{du}{dx}\right) dx + \left(\frac{du}{dy}\right) dy$, u being a function of x and y, yet how readily is the same result obtained from the expansion of $f(y+k,\ x+h)$. To these remarks I may add, that the experience of twenty years, has convinced me that the method of series is that which is best suited to the instruction of a class, and the sale of three large editions, has also proved that my opinion is numerically at least well supported.

The symbol $\frac{du}{dx}$ for the differential coefficient of $u = f(x)$, invented by Leibnitz, and used almost without exception by the continental writers, is here retained—I mention the fact, since the notation $d_x u$ for the same term has been revived by some Cambridge Mathematicians.—I do not pretend to decide the question which of the two, $\frac{du}{dx}$ or $d_x u$, estimated by its power of best representing the differential coefficient ought to be preferred, but I see that the latter is, to say the least, an imperfect notation; and is liable to the objection that the suffix x, in the calculus of finite differences has a meaning entirely different from that indicated by the x in d_x. But the most important objection is that already alluded to, namely, that when a student has become acquainted with the proposed notation from his elementary books, his eye must be familiarized with that of Leibnitz, before the works of Lacroix and Laplace can be read with advantage.

Lastly, if it be considered necessary to offer an inducement to any one to enter upon the study of a science—which is the result of one of Newton's most brilliant discoveries, let him know "that it is a high privilege, not a duty, to study this language of pure unmixed truth. The laws by which God has thought good to govern the universe are surely subjects of lofty contemplation, and the study of that symbolical language by which alone these laws can be fully decyphered, is well deserving of his noblest efforts*."

* Professor Sedgwick on the Studies of the University.

KING'S COLLEGE, LONDON,
June 1, 1846.

TABLE OF CONTENTS.

DIFFERENTIAL CALCULUS.

CHAPTER I.

	PAGE
DEFINITIONS and Principles	1
To prove that $u_1 = f(x+h) = u + Ah + Uh^2$	4
Definitions of Differential and Differential Coefficient	6
Rules for finding the Differential Coefficient	8
Examples	12

CHAPTER II.

Differentiation of Circular Functions	20
Differentiation of Exponential and Logarithmic Functions	27
Examples	29

CHAPTER III.

Successive Differentiation	35
Maclaurin's Theorem	38
Examples	39
Series for the Length of the Circle	42
Series for the Computation of Logarithms	45
Exponential Series	47
Series dependent on the Circle	50
Multinomial Theorem	53
Examples	54

CHAPTER IV.

Taylor's Theorem	56
Examples	58
Approximation to the Root of an Equation	60
Value of $\frac{d^n u}{dx^n}$ investigated	62
Examples	64

CHAPTER V.

Failure of Taylor's Theorem—Explanation of the Term	66
Limits of Taylor's Theorem	70
Separation of Symbols of Operation from those of Quantity	75

CHAPTER VI.

PAGE
Vanishing Fractions 76
Examples 79

CHAPTER VII.

Maximа and Minima defined 83
Rules for determining Maxima or Minima 85
Examples and Problems 88
The Cells of Bees 101
Case of Maximum or Minimum when $\frac{du}{dx} = \infty$ 103

CHAPTER VIII.

Functions of two or more Variables 105
Expansion of $f(x+h,\ y+k)$ 106
Differentiation of Functions of two or more Variables 109
Examples 113
Homogeneous Functions 115
Implicit Functions 117
Elimination by means of Differentiation 121

CHAPTER IX.

Maxima and Minima of Functions of two Variables 126
Examples 127

CHAPTER X.

Equations to Curves 136
Equations to a Point 137
The Straight Line 138
The Circle 142
Transformation of Co-ordinates 148
Parabola, Ellipse, Hyperbola 145
The General Equation of the Second Degree 149
Cissoid, Conchoid, Witch, Logarithmic Curve, Quadratrix, Lemniscata, Cycloid, Trochoid, Epicycloid, Hypocycloid, Spirals, Involute of Circle 151

CHAPTER XI.

Tangents to Curves—Equation to Tangent 162
Equation to Normal 164
Values of Tangent, Sub-tangent, Normal, and Sub-normal ... 166
Asymptotes 170
Examples 171

CHAPTER XII.

PAGE
The Differentials of the Areas and Lengths of Curves 184
If A be the Area of a Curve, $\frac{dA}{dx} = y$ 185
If s be the Length, $\frac{ds}{dx} = \sqrt{1 + \frac{dy^2}{dx^2}}$ 186
If V be the Volume of a Solid of Revolution, $\frac{dV}{dx} = \pi y^2$ 187
If S be the Surface of the same Solid,
$$\frac{dS}{dx} = 2\pi y \sqrt{1 + \frac{dy^2}{dx^2}}$$ 188
Value of the Perpendicular on the Tangent in Spirals 190
Differential Coefficients of A and s in Spirals 191
Tangents and Asymptotes to Spirals 193
To find the Locus of y 195
Examples 196

CHAPTER XIII.

Singular Points in Curves 200
When a Curve is Concave or Convex 201
Points of Contrary Flexure 202
Multiple Points 207
Conjugate Points 211
Cusps 214
To trace a Curve from its Equation 217
Examples 223

CHAPTER XIV.

Curvature and Osculating Curves 226
Order of Contact 227
Osculating Circle—Radius of Curvature 228
Evolute and its Properties 230
Radius of Curvature, and Evolutes of Spirals 233
Examples 237

CHAPTER XV.

Envelopes to Curves 245
Caustics 248

CHAPTER XVI.

Change of the Independent Variable 256
Lagrange's Theorem 266

INTEGRAL CALCULUS.

CHAPTER I.

PAGE

DEFINITION of Integration 272
Integration of Monomials..... 273
Examples of Simple Integration 275

CHAPTER II.

Rational Fractions 278
Roots of the Denominator, (1) all different, (2) some equal, (3) some impossible 278
Integration of $\frac{1}{(x^2+1)^n}$ 285
Integration by parts 287
Integration of $\frac{1}{(x^n \pm 1)}$ 291
Integration of $\frac{x^r}{x^n \pm 1}$, and $\frac{x^r}{x^{2n} - 2x^n \cdot \cos\alpha + 1}$ 293
Examples..... 299

CHAPTER III.

Irrational Quantities 303
Integration of known Differential Coefficients 304
Binomial Differential Coefficients 310
Examples of Integration by parts 312
Bernoulli's Series 320
Examples..... 321

CHAPTER IV.

Integration of Logarithmic and Exponential Functions 326
Find $\int_t e^{-t^2}$ from $t = -\infty$ to $t = \infty$ 330
Examples..... 331

CHAPTER V.

PAGE
Integration of Circular Functions 333
Integration of $\frac{1}{a+b\cos x}$, $\frac{a'+b'\cos x}{(a+b\cos x)^m}$ 339
Integration of $(a+b\cos x)^m$ 341
Integration of $\log(1+n\cos x)$ 343
Examples 344

CHAPTER VI.

Methods of finding the Areas and Lengths of Curves, and the Volumes and Surfaces of Solids 347
Areas of Curves 349
Lengths of Curves 358
Volumes of Solids 364
Surfaces of Solids 370
Examples 372

CHAPTER VII.

Differential Equations, Order and Degree of 374
Equations of the first Order and first Degree 374
Equations of the first Order and of the n^{th} Degree 394
Equations of the second and higher Orders 400
Lagrange's Variation of Parameters 406
Simultaneous Differential Equations 410
Total Differential Equations 412
Partial Differential Equations 415

CHAPTER VIII.

Calculus of Variations 424
Formula for the Variation of $\int V dx$ 426
Maxima and Minima of Integral Formulæ 429
Isoperimetrical Problems 438

By the Rev. T. G. HALL, M.A.,

THE ELEMENTS OF ALGEBRA.

Post Octavo, 6*s.* 6*d.*

ELEMENTS OF DESCRIPTIVE GEOMETRY.

Post Octavo, 6*s.* 6*d.*

THE

DIFFERENTIAL CALCULUS.

CHAPTER I.

1. One quantity u is said to be a function of another x when the value of the magnitude of u depends upon the variation of x. Thus the area of a triangle is a function of the base, when the altitude remains unaltered, since the area will increase or decrease with the increase or decrease of the base.

And if $u = ax^2 \pm bx$, where a and b are constant quantities, and x a variable one, u is said to be a function of x, since if x changes, the value of u will be altered: this relation between u and x is usually expressed by writing $u = f(x)$ or $\phi(x)$, the symbols f and ϕ expressing the word function.

The quantities expressed by the letters a and b are omitted in the equation $u = f(x)$. Since, although they determine the particular kind of function, they remain unchanged, while x passes through every degree of magnitude.

The quantity x is called the independent variable, and u the dependent variable.

2. Functions are called explicit and implicit: u is an explicit function of x, when u is known in terms of x, as in the equation $u = ax^2 + bx$. An implicit function is when u and x are involved together, as in the equation $u^2x - aux + bx^2 = 0$. An implicit function is written $f(u, x)$ or $\phi(u, x) = 0$.

3. Functions are also divided into algebraical and transcendental.

Algebraical functions are those where u may be expressed in terms of x, by means of an equation consisting of a finite number of terms.

Thus $u = ax^m + bx^{m-1} + \&c. + qx^2 + rx + s$ where (m) is finite, is an *algebraical* function of x.

A *Transcendental* function is one where u is equal to an infinite series, the sum of which cannot be expressed by a limited number of terms.

Thus $u = \log(1+x)$, which

$$= \frac{1}{A}\left\{x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \text{\&c. to infinity}\right\},$$

and $u = \sin x = x - \frac{x^3}{2.3} + \frac{x^5}{2.3.4.5} - \text{\&c. to infinity}$,

are transcendental functions of x*.

4. Functions are also called continuous or discontinuous. A function is continuous, when it undergoes a gradual change; and it is discontinuous, when the change is not gradual, or when the function changes suddenly from one value to another very different value. And thus when the difference between $f(x)$ and $f(x+h)$ may, by the continued diminution of h, be made as small as we please, $f(x)$ is a continuous function; but when under the same circumstances $f(x+h)$ differs widely from $f(x)$, the latter is a discontinuous function. We may familiarly liken a continuous quantity, to a stream of water flowing equably and steadily through a tube, and a discontinuous one, to water falling interruptedly, or in drops, from a height.

5. The equation $u = f(x)$ expresses the relation between the function u and the *single* variable x, and the values of u solely depend upon the change that may take place in x: but if we have an equation between three unknown quantities, such as $u = ax^2y - bxy^2$, where x and y are independent of each other, i.e. not connected together by any other equation; then the value of u depends upon the change, both of x and y, and u is said to be a function of two variables; this is expressed by writing $u = f(x, y)$.

As an instance, we may again take the area of a triangle, the magnitude of which depends upon the rectangle of the base and the altitude, which lines are totally independent of each other.

It is obvious that there may be functions of three, four, or of n variables.

6. But to return to functions of one variable; let $u = f(x)$ express the relation between the function and its independent variable x.

Let x increase and become $x+h$, then the value of u

* $u = 1 + x + x^2 + x^3 + \text{\&c. to infinity}$ is an algebraical function of x, since the sum of the series is expressed by $\frac{1}{1-x}$.

will most probably be altered. Let the new value be represented by u_1, then $u_1 = f(x+h)$,

and $u = f(x)$, by hypothesis;

$$\therefore u_1 - u = f(x+h) - f(x).$$

Now $u_1 - u$, or the difference between the functions of $x+h$ and x, must depend upon h, and we shall first shew that it may be expressed by a series of the form

$$Ah + Bh^2 + Ch^3 + \&c.;$$

and $\therefore$ that $u_1 = u + Ah + Bh^2 + Ch^3 + \&c.$,

or that u_1 is equal to u + a series of terms involving positive and integral powers of h, which ascend from the simple power: the primary object of the Differential Calculus is to find the coefficients A, B, C, &c.

7. We will first shew that u_1 may be expressed by a series of the above form by a few particular examples:

(1) Let $u = x^3$;

$$\therefore u_1 = (x+h)^3 = x^3 + 3x^2h + 3xh^2 + h^3$$
$$= u + 3x^2h + 3xh^2 + h^3,$$

which is of the required form.

(2) Next, let $u = x^n$;

$$\therefore u_1 = (x+h)^n = x^n + nx^{n-1}h + n\frac{(n-1)}{2}x^{n-2}h^2 + \&c.$$

by the Binomial Theorem,

Or, putting u for x^n,

$$u_1 = u + nx^{n-1}h + n\frac{n-1}{2}x^{n-2}h^2 + \&c.,$$

a series with ascending powers of h.

(3) Let $u = Ax^m + Bx^n + Cx^p + \&c.$;

$$\therefore u_1 = A(x+h)^m + B(x+h)^n + C(x+h)^p + \&c.$$
$$= A\left(x^m + mx^{m-1}h + m\frac{(m-1)}{2}x^{m-2}h^2 + \&c.\right)$$
$$+ B\left(x^n + nx^{n-1}h + n\frac{(n-1)}{2}x^{n-2}h^2 + \&c.\right)$$
$$+ C\left(x^p + px^{p-1}h + p\frac{p-1}{2}x^{p-2}h^2 + \&c.\right)$$
$$+ \&c.$$

$$= Ax^m + Bx^n + Cx^p + \&c.$$
$$+ (mAx^{m-1} + nBx^{n-1} + \&c.)h$$
$$+ \left\{m\frac{(m-1)}{2}.Ax^{m-2} + n\frac{(n-1)}{2}Bx^{n-2} + \&c.\right\}h^2$$
$$+ \&c. \qquad + \&c.$$
$$= u + ph + qh^2 + \&c.$$

by writing u for its value, $Ax^m + Bx^n + \&c.$, and putting p, q, &c. for the coefficients of h, h^2, &c.

(4) It may also be shewn that a^{x+h}, $\log(x+h)$, $\sin(x+h)$, can be expanded into series of the form

$$u + Ah + Bh^2 + Ch^3, \&c.$$

but we proceed to demonstrate the following general Proposition.

8. Prop. If $u = f(x)$, and u_1 be the value of u when x becomes $x + h$, then shall

$$u_1 = u + Ah + Uh^2,$$

where u is the original function, and Uh^2, represents all the terms that follow Ah. It is however necessary to prove,

(1) That u_1 or $f(x+h)$ can only contain powers of h with positive indices. For if

$$u_1 = M + Ah^\alpha + Bh^{-\beta} + \&c. = M + Ah^\alpha + \frac{B}{h^\beta} + \&c.$$

when $h = 0$, u_1 instead of becoming $= u$, would be infinite.

(2) That none of the indices of h can be fractional; for if possible let

$$u_1 = M + Ph^{\frac{m}{n}} + R.$$

then $\because$ $h^{\frac{m}{n}}$ or $\sqrt[n]{h^m}$ has n different values, let h_1 and h_2 be two of them;

$$\therefore u_1 = M + Ph_1 + R; \text{ and } u_1 = M + Ph_2 + R;$$

$\therefore$ subtracting, $P(h_1 - h_2) = 0$; $\therefore P = 0$; or there is no term of the form $Ph^{\frac{m}{n}}$.

(3) That the first term of the expansion $= u$.

For let u_1 or $f(x+h) = M + Ah^\alpha + \&c.$ then let $h = 0$;

$\therefore f(x) = u = M$, or $M = u$; and $u_1 = u + Ah^\alpha + \&c.$

Let $\therefore$ u_1 or $f(x+h) = u + Ah^\alpha + Bh^\beta + Ch^\gamma + \&c.$,

where α is the least of the indices of h, and β the next in magnitude, and A, B, &c. are functions of x.

Now whether x become $x + h$, or h become $h + k$ or $2h$, u_1 will become $f(x + 2h)$, and the expansions on either supposition must be identical.

(1) Let h become $2h$, and let u_2 be the value of u_1;

$$\therefore\ u_2 = f(x + 2h)$$
$$= u + A(2h)^\alpha + B(2h)^\beta + \&c.$$
$$= u + 2^\alpha A h^\alpha + 2^\beta B h^\beta + \&c. \ldots\ldots\ldots (1).$$

(2) Let x become $x + h$, then u_1 is as before $f(x + 2h)$ or u_2; also let (u), (A), (B), &c. represent the values of u, A, B, &c., for these undergo change, $\because$ they are functions of x;

$$\therefore\ u_2 = (u) + (A)h^\alpha + (B)h^\beta + \&c.$$

But (u) is the same as u_1, for it is $f(x + h)$,

$$\therefore\ (u) = u + Ah^\alpha + Bh^\beta + \&c.$$

Also (A), (B), &c. being what A, B, &c. become by putting $x + h$ for x, must be of a similar form;

$$\therefore\ (A) = A + A_1 h^{\alpha_1} + A_2 h^{\beta_1} + \&c.$$
$$(B) = B + B_1 h^{\alpha_2} + B_2 h^{\beta_2} + \&c.$$

then multiplying (A) by h^α and (B) by h^β, and substituting,

$$u_2 = u + Ah^\alpha + Bh^\beta + Ch^\gamma + \&c.$$
$$+ Ah^\alpha + A_1 h^{\alpha+\alpha_1} + A_2 h^{\alpha+\beta_1} + \&c.$$
$$+ Bh^\beta + B_1 h^{\alpha_2+\beta} + \&c.$$
$$= u + 2Ah^\alpha + A_1 h^{\alpha+\alpha_1} + 2Bh^\beta + \&c. \ldots (2).$$

Equating the coefficients of h^α in series (1) and (2),

$$2A = 2^\alpha A;\quad 2 = 2^\alpha;\quad \therefore\ \alpha = 1,$$
$$\text{and } u_1 = f(x + h) = u + Ah + Bh^\beta + \&c.$$

whence it appears that the *second term of the expansion of* $f(x + h)$ *contains the first power of* h *only.*

Hence also $\alpha_1 = 1$ for $A_1 h^{\alpha_1}$ is the second term of the expression for A when x becomes $x + h$;

$$\therefore\ u_2 = u + 2Ah + A_1 h^2 + 2Bh^\beta + \&c. \text{ from (2)}$$
$$= u + 2Ah + 2^\beta . Bh^\beta + \&c. \ldots\ldots\ldots\ldots (1).$$

Now, since in series (2) a term is found involving h^2, some

corresponding term must be found in series (1); and as β is less than any index that follows it, β must $= 2$;

$$\therefore\ u_1 = f(x+h) = u + Ah + Bh^2 + Ch^\gamma + \&c.$$
$$= u + Ah + (B + Ch^{\gamma-2} + \&c.)h^2$$
$$= u + Ah + Uh^2.$$

Cor. From this it follows that $\beta_1 = 2$; and therefore $\alpha + \beta_1 = 3$; and therefore $A_2 h^{\alpha+\beta_1} = A_2 h^3$; which being a term in series (2), there must be a corresponding term in series (1); whence $\gamma = 3$, and similarly the next index $= 4$, and so on;

$$\therefore\ u_1 = f(x+h) = u + Ah + Bh^2 + Ch^3 + Dh^4 + \&c.$$

it will afterwards be shewn, by a Theorem, called Taylor's Theorem, that the coefficients, A, B, C, &c. have a dependence on each other.

9. The second term of the expansion, or Ah, is called the differential of u: differential being the diminutive of difference; for Ah is the first term of the difference between u_1 and u, and is consequently a part only of the difference: but the difference and differential differ the less, the less h is, and in cases of approximation, the latter is sometimes taken for the former.

Instead of writing differential at full length, the letter d is used, thus du is put for differential of u, and thus $du = Ah$: but as then h is called the differential of x, therefore for symmetry of notation dx is put for h, and thus $du = A\,dx$.

A is called the *first differential coefficient*, and is expressed by the symbol $\frac{du}{dx}$, when $u = f(x)$.

Hence we define a *differential* to be *the second term of the* expansion of $f(x+h)$, and the differential coefficient to be the *coefficient of the first power of h.*

The process by which A, or $\frac{du}{dx}$ is found is called *differentiation.*

From these definitions we see that the differential of u, is the product of A into the differential of x; or calling the first quantity δu, and the second δx, we have

$$\delta u = A\delta x;\quad \therefore \frac{\delta u}{\delta x} = A = \frac{du}{dx},$$

or the ratio of the differentials of u and x is equal to the ratio of the differential coefficient to unity.

The letter δ is here used only to avoid confounding the differentials with the differential coefficient, but in general we make use of the letter d.

10. Again, since $u_1 = u + Ah + Uh^2$,

$$\therefore \frac{u_1 - u}{h} = A + Uh;$$

but $u_1 - u$ is the increment of u, and h is the increment of x; therefore the ratio of the increment of the function, to the increment of x, $= A + Uh$; and as h decreases, this ratio tends to A as its limit, and when h vanishes actually $= A$.

That is, A or $\frac{du}{dx}$ is the *limit of the ratio of the increment of the function to that of the variable upon which it depends.*

Cor. Hence also, the ratio of the differentials of u and x, is equal to the limit of the ratio of the increments of u and x.

11. Hence we have a method of finding the differential coefficient which is frequently very convenient. Expand $f(x + h)$, subtract $f(x)$, divide both sides by h, then make $h = 0$; and the term or terms remaining of the expansion will be equal to the coefficient required.

12. We have seen that if u be any function of x, and x become $x + h$,

$$f(x + h) = u + \frac{du}{dx} h + Uh^2.$$

Similarly, if z, v, &c. be functions of x, then they will respectively become when x is made $x + h$,

$$z + \frac{dz}{dx} h + Zh^2, \text{ and } v + \frac{dv}{dx} h + Vh^2,$$

where Zh^2 and Vh^2 represent all the terms after the first two.

13. Thus it appears, that in order to find the differential or differential coefficient, we have merely to put $x + h$ for x, and expand $f(x + h)$ according to the powers of h, and the term corresponding to Ah will give us at once both of the objects of our enquiry. But such a direct process would always be tedious, and often almost impracticable. We therefore proceed to investigate rules which will not only greatly diminish the labour of *differentiation*, but render it a simple

algebraical operation; but we will first apply the general process to the function

$$u = \frac{a+x}{b+x};$$

$$\therefore u_1 = \frac{a+x+h}{b+x+h} = \frac{\dfrac{a+x}{b+x} + \dfrac{h}{b+x}}{1+\dfrac{h}{b+x}}$$

$$= \left(\frac{a+x}{b+x} + \frac{h}{b+x}\right) \cdot \frac{1}{1+\dfrac{h}{b+x}}$$

$$= \left(\frac{a+x}{b+x} + \frac{h}{b+x}\right) \cdot \left\{1 - \frac{h}{b+x} + \frac{h^2}{(b+x)^2} - \&c.\right\}$$

$$= \frac{a+x}{b+x} + h\left\{\frac{1}{b+x} - \frac{a+x}{(b+x)^2}\right\} + ph^2 + \&c.;$$

$$\therefore \frac{du}{dx} = \frac{1}{b+x} - \frac{a+x}{(b+x)^2} = \frac{b-a}{(b+x)^2}.$$

Again, since $u = \dfrac{a+x}{b+x}$, we shall have by the same process

$$u_1 = u + h \cdot \left\{\frac{b-a}{(b+x)^2}\right\} + ph^2 + qh^3 + \&c.$$

$$\text{and } \frac{u_1 - u}{h} = \frac{b-a}{(b+x)^2} + ph + qh^2 + \&c.;$$

and by making $h = 0$, as in Art. 11,

$$\frac{du}{dx} = \frac{b-a}{(b+x)^2}.$$

RULES FOR FINDING THE DIFFERENTIAL COEFFICIENT.

14. We repeat the definition of Art. 9, that if $u = f(x)$; $\dfrac{du}{dx}$ is the coefficient of the first power of h in the expansion of u_1, or of $f(x+h)$.

Let $u = ax$, a being a constant quantity;

$$\therefore u_1 = a(x+h) = ax + ah = u + ah;$$

$$\therefore \frac{du}{dx} = a \text{ or } \frac{d(ax)}{dx} = a.$$

Cor. If $u = x$; $\therefore \frac{du}{dx} = 1$, $\because a = 1$.

15. Let $u = ax \pm b$, where a and b are constant;

$$\therefore u_1 = a(x+h) \pm b = ax \pm b + ah = u + ah;$$

$$\therefore \frac{du}{dx} = a, \text{ that is, } \frac{d(ax \pm b)}{dx} = a.$$

But by the preceding Article, $\frac{d(ax)}{dx} = a$;

$$\therefore \frac{d(ax \pm b)}{dx} = \frac{d(ax)}{dx};$$

that is, constant quantities connected with a variable one by the signs $\pm$ disappear in differentiation.

16. Let $u = ax^m$. Then,

$$u_1 = a(x+h)^m = a.(x^m + mx^{m-1}h + \&c.)$$

$$= ax^m + max^{m-1}.h + \&c.;$$

$$\therefore \frac{du}{dx} = max^{m-1},$$

or to find the differential coefficient of ax^m, *multiply by the index and then diminish the index by unity.*

Ex. $u = 5x^7$; $\therefore \frac{du}{dx} = 35x^6$.

17. Let $u = az$ where z is a function of x;

therefore if x become $x + h$,

z becomes, $z + \frac{dz}{dx}h + Zh^2$;

$$\therefore u_1 = az + a\frac{dz}{dx}h + aZh^2;$$

$$\therefore \frac{du}{dx} \text{ or } \frac{d(az)}{dx} = a.\frac{dz}{dx}.$$

18. If $u = az + b$, a and b being constant quantities,

$$\text{then } \frac{du}{dx} = a.\frac{dz}{dx},$$

$$\therefore \frac{d(ax+b)}{dx} = \frac{a dx}{dx} = \frac{d(ax)}{dx}.$$

19. Let $u = x + v + w + \&c.$ where x, v, w are each functions of x;

$$\therefore u + \frac{du}{dx}h + \&c. = x + \frac{dx}{dx}h + v + \frac{dv}{dx}h + w + \frac{dw}{dx}h + \&c.;$$

$$\therefore \frac{du}{dx} = \frac{dx}{dx} + \frac{dv}{dx} + \frac{dw}{dx} + \&c.$$

$$\text{or } \frac{d.(x + v + w + \&c.)}{dx} = \frac{dx}{dx} + \frac{dv}{dx} + \frac{dw}{dx} + \&c.$$

Or the differential coefficient of the sum of any functions equals the sum of the differential coefficients of each function taken separately.

20. To find the differential coefficient of the product of two functions.

Let $u = xv$;

$$\therefore u_1 = (x + \frac{dx}{dx}h + Zh^2)(v + \frac{dv}{dx}h + Vh^2)$$

$$= xv + \left(x\frac{dv}{dx} + v.\frac{dx}{dx}\right)h + Bh^2 + \&c.,$$

$$\text{where } B = Zv + Vx + \frac{dx}{dx}.\frac{dv}{dx};$$

$$\therefore \frac{du}{dx} = x\frac{dv}{dx} + v.\frac{dx}{dx};$$

or the differential coefficient of the product of two functions equals the sum of the products of each function into the differential coefficient of the other.

21. To find the differential coefficient of the quotient of two functions.

Let $u = \frac{x}{v}$. Here $\frac{du}{dx}$ may be found by substituting the values which u, x, and v have when x becomes $x + h$; but it may be readily deduced from the preceding Article.

$$\because u = \frac{x}{v}; \quad \therefore vu = x; \quad v\frac{du}{dx} + u\frac{dv}{dx} = \frac{dx}{dx}.$$

$$\therefore \frac{du}{dx} = \frac{1}{v}\cdot\frac{dz}{dx} - \frac{u}{v}\cdot\frac{dv}{dx}$$

$$= \frac{1}{v}\cdot\frac{dz}{dx} - \frac{z}{v^2}\cdot\frac{dv}{dx}$$

$$= \frac{v\cdot\dfrac{dz}{dx} - z\cdot\dfrac{dv}{dx}}{v^2}.$$

A simple expression, the form of which is more easily remembered than the enunciation.

22. Let $u = zvw$, writing vw for v in Art. 20;

$$\therefore \frac{du}{dx} = z\cdot\frac{d(vw)}{dx} + vw\frac{dz}{dx}.$$

$$\text{But } \frac{d.(vw)}{dx} = v\cdot\frac{dw}{dx} + w\cdot\frac{dv}{dx};$$

$$\therefore \frac{du}{dx} = zv\cdot\frac{dw}{dx} + zw\cdot\frac{dv}{dx} + vw\cdot\frac{dz}{dx}.$$

Similarly may the differential coefficient be found for the product of n functions, and it will be equal to the sum of the n products of the differential coefficient of each of the functions multiplied by the remaining $n-1$ functions. Thus,

$$\frac{d.\{z.v.w.s\ldots(n)\}}{dx} = v.w.s\ldots(n-1)\frac{dz}{dx} + zws\ldots(n-1)\cdot\frac{dv}{dx}.$$

$$+ zvs\ldots(n-1)\frac{dw}{dx} + \&c.$$

23. Lemma. If u be a function of z, and z be a function of x, then

$$\frac{du}{dx} = \left(\frac{du}{dz}\right)\left(\frac{dz}{dx}\right).$$

For if δu, δz, δx be the corresponding differentials of u, z, and x;

$$\text{then } \because \frac{a}{c} = \frac{a}{b}\times\frac{b}{c}; \quad \therefore \frac{\delta u}{\delta x} = \frac{\delta u}{\delta z}\cdot\frac{\delta z}{\delta x}.$$

$$\text{But } \frac{\delta u}{\delta x} = \frac{du}{dx}; \quad \frac{\delta u}{\delta z} = \frac{du}{dz}; \quad \frac{\delta z}{\delta x} = \frac{dz}{dx};$$

$$\therefore \frac{du}{dx} = \left(\frac{du}{dz}\right)\left(\frac{dz}{dx}\right);$$

an important theorem, of which we shall hereafter give another demonstration.

24. Let $u = z^n$, z being $= f(x)$; find $\frac{du}{dx}$.

$$\frac{du}{dz} = nz^{n-1}; \quad \therefore \frac{du}{dx} = \frac{du}{dz} \cdot \frac{dz}{dx} = nz^{n-1} \cdot \frac{dz}{dx};$$

or, to find the differential coefficient of z^n, *multiply by the index, diminish the index by unity, and then multiply by the differential coefficient of* z.

Ex. If $u = (a^2 + x^2)^n$ then $z = a^2 + x^2$ and $\frac{dz}{dx} = 2x$;

$$\therefore \frac{du}{dx} = n(a^2 + x^2)^{n-1} \cdot 2x.$$

25. The rule for finding the differential coefficient of z^n is perfectly general, but when $n = \frac{1}{2}$ it has a value which it is useful to remember. Thus

$$\frac{d(\sqrt{z})}{dx} = \tfrac{1}{2} z^{\frac{1}{2}-1} \frac{dz}{dx} = \frac{\frac{dz}{dx}}{2\sqrt{z}};$$

whence this rule. To find the differential coefficient of the square root of any quantity, divide the differential coefficient of the quantity under the square root, by twice the square root of the quantity itself.

Ex. Let $u = \sqrt{a + bx + cx^2}$;

$$\therefore z = a + bx + cx^2; \quad \therefore \frac{dz}{dx} = b + 2cx;$$

$$\text{and } \frac{du}{dx} = \frac{b + 2cx}{2\sqrt{a + bx + cx^2}}.$$

EXAMPLES.

(1) $u = 3x^{\frac{7}{3}}$; $\therefore \frac{du}{dx} = 3 \cdot \frac{7}{3} \cdot x^{\frac{7}{3}-1} = 7x^{\frac{4}{3}}$.

(2) $u = x^3 + x^2 + x + 1$;

$$\frac{du}{dx} = 3x^2 + 2x + 1.$$

(3) $u = (x+a).(x+b)$;

$$\frac{du}{dx} = (x+a)\frac{d(x+b)}{dx} + (x+b).\frac{d(x+a)}{dx}. \text{ Art. (20)}$$

$$= x + a + x + b = 2x + (a+b).$$

(4) $u = x(1+x^2)(1+x^3)$;

$$\frac{du}{dx} = (1+x^2)(1+x^3)\frac{dx}{dx} + x(1+x^3).\frac{d(1+x^2)}{dx} + x(1+x^2).\frac{d(1+x^3)}{dx}$$

$$= (1+x^2)(1+x^3) + x(1+x^3).2x + x(1+x^2)3x^2$$

$$= 1 + 3x^2 + 4x^3 + 6x^5.$$

(5) $u = \frac{a^n}{x^n} = a^n x^{-n}$; $\frac{du}{dx} = -na^n x^{-n-1} = -\frac{na^n}{x^{n+1}}$;

$\therefore$ if $u = \frac{a}{x}$, $\frac{du}{dx} = -\frac{a}{x^2}$.

(6) $u = \frac{x+a}{x+b}$; See (Art. 13)

$$\therefore \frac{du}{dx} = \frac{(x+b).\frac{d(x+a)}{dx} - (x+a).\frac{d(x+b)}{dx}}{(x+b)^2}$$

$$= \frac{x+b-(x+a)}{(x+b)^2} = \frac{b-a}{(x+b)^2}.$$

(7) $u = \frac{x^m}{(x+1)^m}$;

$$\therefore \frac{du}{dx} = \frac{(x+1)^m . mx^{m-1} - x^m . m . \overline{x+1}^{m-1}}{(x+1)^{2m}}$$

$$= \frac{(x+1).mx^{m-1} - mx^m}{\overline{x+1}^{m+1}} = \frac{mx^{m-1}}{\overline{x+1}^{m+1}}.$$

(8) $u = \sqrt{1+x^2}$;

$$\frac{du}{dx} = \frac{2x}{2\sqrt{1+x^2}} = \frac{x}{\sqrt{1+x^2}}. \text{ Art. (25).}$$

(9) $u = \sqrt{x+\sqrt{1+x^2}}$;

$$\frac{du}{dx} = \tfrac{1}{2}(x+\sqrt{1+x^2})^{-\frac{1}{2}}.\frac{d.(x+\sqrt{1+x^2})}{dx},$$

and $\dfrac{d(x+\sqrt{1+x^2})}{dx} = 1 + \dfrac{x}{\sqrt{1+x^2}} = \dfrac{x+\sqrt{1+x^2}}{\sqrt{1+x^2}}$;

$$\therefore \frac{du}{dx} = \frac{1}{2} \cdot \frac{\sqrt{x+\sqrt{1+x^2}}}{\sqrt{1+x^2}}.$$

(10) $u = (2ax+x^2)^m$,

$$\frac{du}{dx} = m(2ax+x^2)^{m-1} \cdot \frac{d(2ax+x^2)}{dx}$$

$$= 2m \cdot (2ax+x^2)^{m-1} \cdot (a+x).$$

(11) $u = \dfrac{\sqrt{a^2+x^2}}{\sqrt{a^2-x^2}}$; $\therefore u^2 = \dfrac{a^2+x^2}{a^2-x^2}$;

$$\therefore 2u\frac{du}{dx} = \frac{2x(a^2-x^2)+2x(a^2+x^2)}{(a^2-x^2)^2} = \frac{4a^2x}{(a^2-x^2)^2};$$

$$\therefore \frac{du}{dx} = \frac{2a^2x}{(a^2-x^2)^2} \frac{\sqrt{a^2-x^2}}{\sqrt{a^2+x^2}} = \frac{2a^2x}{(a^2-x^2)^{\frac{3}{2}}\sqrt{a^2+x^2}}.$$

(12) $u = \dfrac{\sqrt{1+x}+\sqrt{1-x}}{\sqrt{1+x}-\sqrt{1-x}} = \dfrac{(\sqrt{1+x}+\sqrt{1-x})^2}{2x}$

$$= \frac{1+\sqrt{1-x^2}}{x};$$

$$\therefore \frac{du}{dx} = \frac{\dfrac{-x^2}{\sqrt{1-x^2}} - (1+\sqrt{1-x^2})}{x^2} = -\frac{x^2+\sqrt{1-x^2}+1-x^2}{x^2\sqrt{1-x^2}}$$

$$= -\frac{1+\sqrt{1-x^2}}{x^2\sqrt{1-x^2}}.$$

(13) $u = x(1+x^2) \cdot \sqrt{1-x^2} = (x+x^3)\sqrt{1-x^2}$,

$$\frac{du}{dx} = (1+3x^2)\sqrt{1-x^2} - \frac{x^2+x^4}{\sqrt{1-x^2}}$$

$$= \frac{1+3x^2-x^2-3x^4-x^2-x^4}{\sqrt{1-x^2}}$$

$$= \frac{1+x^2-4x^4}{\sqrt{1-x^2}}.$$

(14) $u = x^3(1+x^4)$; $\dfrac{du}{dx} = 2x(1+3x^4)$.

(15) $u = (1+2x^2)(1+4x^3)$; $\dfrac{du}{dx} = 4x(1+3x+10x^3)$.

(16) $u = 2x(1+x)^{\frac{5}{2}}$; $\dfrac{du}{dx} = (2+7x)(1+x)^{\frac{3}{2}}$.

(17) $u = (1+x)^4(1+x^2)^2$;

$$\frac{du}{dx} = 4(1+x)^3(1+x^2)\{1+x+2x^2\}.$$

(18) $u = (a+x)(b+x)(c+x)$;

$$\frac{du}{dx} = 3x^2 + 2(a+b+c)x + ab + ac + bc.$$

(19) $u = (1-2x)(1-3x)(1-4x)$;

$$\therefore \frac{du}{dx} = -(9 - 52x + 72x^2).$$

(20) $u = (1+x^2)^3(1+x^3)^2$

$$\frac{du}{dx} = 6x(1+x^2)^2(1+x^3)(1+x+2x^3).$$

(21) $u = (1+x^m)^n(1+x^n)^m$.

$$\therefore \frac{du}{dx} = mn(1+x^m)^{n-1}(1+x^n)^{m-1}\{x^{m-1} + x^{n-1} + 2x^{m+n-1}\}.$$

(22) $u = (1+x)\sqrt{1-x^2}$;

$$\frac{du}{dx} = \frac{1-x-2x^2}{\sqrt{1-x^2}};$$

(23) $u = \dfrac{x^4-1}{x^4+1}$; $\dfrac{du}{dx} = \dfrac{8x^3}{(x^4+1)^2}$.

(24) $u = \dfrac{x}{\sqrt{1+x^2}}$; $\dfrac{du}{dx} = \dfrac{1}{(1+x^2)^{\frac{3}{2}}}$.

(25) $u = \dfrac{x^2-2}{3}\sqrt{1+x^2}$; $\dfrac{du}{dx} = \dfrac{x^3}{\sqrt{1+x^2}}$.

(26) $u = x^{\frac{1}{3}}\sqrt{x^{\frac{1}{2}}+1}$; $\dfrac{du}{dx} = \dfrac{7x^{\frac{1}{2}}+4}{12\sqrt[3]{x^2}\sqrt{x^{\frac{1}{2}}+1}}$.

(27) $u = \dfrac{x^4 - x^2 + 1}{x^4 + x^2 + 1}$; $\dfrac{du}{dx} = \dfrac{4x(x^4 - 1)}{(x^4 + x^2 + 1)^2}$.

(28) $u = \dfrac{x^3}{\sqrt{1 + x^6}}$; $\dfrac{du}{dx} = \dfrac{3x^3}{(1 + x^6)^{\frac{3}{2}}}$.

(29) $u = \dfrac{\sqrt{1 + x^3} - 1}{\sqrt{1 + x^2} + 1}$; $\dfrac{du}{dx} = \dfrac{2(\sqrt{1 + x^3} - 1)^2}{x^3\sqrt{1 + x^3}}$.

(30) $u = \dfrac{(x + 1)^{\frac{3}{2}}}{\sqrt{x - 1}}$; $\dfrac{du}{dx} = \dfrac{(x - 2)\sqrt{x + 1}}{(x - 1)^{\frac{3}{2}}}$.

(31) $u = \sqrt{\dfrac{1 - \sqrt{x}}{1 + \sqrt{x}}}$; $\dfrac{du}{dx} = -\dfrac{1}{2(1 - \sqrt{x})\sqrt{x - x^2}}$.

(32) $u = \dfrac{1 + 2x}{\sqrt{1 + x + x^2}}$; $\dfrac{du}{dx} = \dfrac{3}{2(1 + x + x^2)^{\frac{3}{2}}}$.

(33) $u = \dfrac{\sqrt{1 - x + x^2}}{\sqrt{1 + x + x^2}}$; $\dfrac{du}{dx} = \dfrac{x^2 - 1}{(1 - x + x^2)^{\frac{1}{2}}(1 + x + x^2)^{\frac{3}{2}}}$.

(34) $u = \sqrt[4]{\left(a - \dfrac{b}{\sqrt{x}} + \sqrt[3]{(c^2 - x^2)^2}\right)^3}$;

$$\frac{du}{dx} = \frac{\dfrac{3b}{2x\sqrt{x}} - \dfrac{4x}{\sqrt[3]{c^2 - x^2}}}{4\sqrt[4]{a - \dfrac{b}{\sqrt{x}} + \sqrt[3]{(c^2 - x^2)^2}}}.$$

(35) Let $u^2x - u + x^2 - a^2 = 0$; find $\dfrac{du}{dx}$, this is an implicit function; put $v = u^2x$, and differentiate;

$$\therefore \frac{dv}{dx} - \frac{du}{dx} + 2x = 0: \text{ but } \frac{dv}{dx} = 2u_a\frac{du}{dx} + u^2 \quad (20);$$

$$\therefore 2ux\frac{du}{dx} + u^2 - \frac{du}{dx} + 2x = 0;$$

$$\therefore \frac{du}{dx}(2ux - 1) = -(u^2 + 2x) = -\frac{u + x^2 + a^2}{x},$$

$$\therefore \frac{du}{dx} = -\frac{u + x^2 + a^2}{2ux^2 - x}.$$

(36) $u^3 - 3ux^2 + x^3 = 0$; find $\frac{du}{dx}$;

$$\therefore 3u^2\frac{du}{dx} - 3x^2\frac{du}{dx} - 6ux + 3x^2 = 0;$$

$$\therefore \frac{du}{dx} = \frac{2ux - x^2}{u^2 - x^2} = \frac{u}{x}.$$

(37) $2ux + au^2 - bx^2 = 0$; $\frac{du}{dx} = \frac{u}{x}$.

(38) $au^4 + c^3ux - bx^4 = 0$; $\frac{du}{dx} = \frac{u}{x}\left(\frac{4au^3 + 3c^3x}{4au^3 + c^3x}\right)$.

(39) $ux = (a + u)\sqrt{b^2 - u^2}$; $\frac{du}{dx} = -\frac{u^3x}{(a + u)(ab^2 + u^3)}$.

(40) $u = \sqrt{a + x + \sqrt{a + x + \sqrt{a + x}}}$ &c. *in infin.*;

$$\therefore \frac{du}{dx} = \frac{1}{\sqrt{1 + 4a + 4x}}.$$

26. In the preceding examples, differential coefficients have only been obtained; but by the definitions, the differentials may be found by multiplying the differential coefficients by the increment of the independent variable.

Thus to find the differential of the product of two functions z and v, if $u = zv$, then

$$\frac{du}{dx} = z\frac{dv}{dx} + v\frac{dz}{dx};$$ let δx = increment of x;

$$\therefore d(zv) = z\frac{dv}{dx}\delta x + v\frac{dz}{dx}\delta x.$$

But $\frac{dv}{dx}\delta x$ = differential of $v = dv$,

and $\frac{dz}{dx}\delta x$ = differential of $z = dz$;

$$\therefore d(zv) = zdv + vdz;$$

and in the same manner,

$$d.\left(\frac{z}{v}\right) = \frac{vdz - zdv}{v^2},$$

and $d.(z^n) = nz^{n-1}dz$.

These expressions are often very convenient, and we see that the differential coefficients may be derived from them by dividing the differentials by dx.

27. It will now be natural for the student to enquire what is the object to be attained, by finding the differential coefficient, but it will be difficult at present to give a completely satisfactory answer to the enquiry, without introducing subjects with which he can have no acquaintance. Lacroix says: "Il serait fort difficile d'expliquer clairement la nature du Calcul différentiel à ceux qui n'en ont pas les premières notions." Yet perhaps he may be told, that if y be the ordinate and x the abscissa of a curve, $\frac{dy}{dx}$ is the trigonometrical tangent of the angle, which the tangent makes with the axis of x; that if u be the area of the same curve, $\frac{du}{dx} = y$; an equation by which hereafter the area of the curve may be found. Again, if s represent the space described by a point in a time t, that $\frac{ds}{dt}$ represents the velocity (v) with which at the end of that time the point is moving, and $\frac{dv}{dt}$, the force which either accelerates or retards the point's motion. And again, if $u = f(x)$ be an equation admitting of maximum or minimum values, that $\frac{du}{dx}$ will furnish an equation, by which the values of x that make u a maximum or minimum may be found. And lastly, if $u = f(x) = 0$ be an algebraic equation of which the roots are a, b, c, &c., then $\frac{du}{dx} = 0$ will give the limiting equation, the knowledge of the roots of which is so useful in determining the roots of the original equation.

28. We shall conclude this Chapter by a few simple applications.

(1) The radius of a circular plate of metal is 12 inches; find the increase of area when the radius is increased .001 inch.

If u = area of a circle, radius = x;

$$\therefore u = \pi x^2; \text{ and } du = 2\pi x\, dx.$$

Make $x = 12$; $dx = .001$; then du = increase of area;

$\therefore$ $du = 3.1416 \times 24 \times .001 = .0753984$ of a square inch.

(2) A cube of metal of the same thickness is similarly increased; find the cubical expansion.

$u = x^3$; $\therefore$ $du = 3x^2dx = 3 \times 144 \times .001 = .432$ cubic inch.

Cor. Divide by u; $\therefore \frac{du}{u} = \frac{3dx}{x}$. Now $\frac{du}{u}$ is called the cubical, and $\frac{dx}{x}$ the linear expansion; hence the cubical is three times the linear expansion.

(3) As an instance of finding the ratio of infinitely small quantities. Upon AB describe a semicircle, draw a chord AP; draw PN perpendicular to AB; then prove that $AP = PN$ ultimately; i.e. at the moment when the arc AP vanishes.

Make $AN = x$, $NB = 2a$;

$$\therefore AP = \sqrt{2ax},\ PN = \sqrt{2ax - x^2};$$

$$\therefore \frac{AP}{PN} = \frac{\sqrt{2ax}}{\sqrt{2ax - x^2}} = \frac{\sqrt{2a}}{\sqrt{2a - x}};$$ and if $x = 0$,

$$\frac{AP}{PN} = \frac{\sqrt{2a}}{\sqrt{2a}} = 1;$$ or $AP = PN$ ultimately.

In the next Chapter we shall find the differential coefficients of the trigonometrical ratios, of exponentials, and of logarithms.

CHAPTER II.

DIFFERENTIATION OF ANGULAR, EXPONENTIAL, AND LOGARITHMIC FUNCTIONS.

---

29. To find the differential coefficient of u, when

$u = \sin x$, $\cos x$, $\tan x$, $\sec x$, &c.

The following Proposition must first be proved.

If h be an angle, $\frac{\sin h}{h}$, or $\frac{\tan h}{h}$ = unity, when $h = 0$.

It is proved, that $h > \sin h, < \tan h$ (*Trig.* Art. 53,)

or h lies between $\sin h$ and $\tan h$,

or $\sin h$, h, and $\tan h$ are in order of magnitude;

$\therefore$ $\tan h - \sin h$, is $> h - \sin h$, or $> \tan h - h$.

If therefore $\tan h - \sin h$ ever $= 0$, or $\frac{\tan h}{\sin h} = 1$; *a fortiori* will $h - \sin h = 0$, and $\tan h - h = 0$; or $\frac{\sin h}{h} = \frac{\tan h}{h} = 1$,

Now $\frac{\sin h}{\tan h} = \frac{\cos h}{1} = \frac{1}{1} = 1$, when $h = 0$;

$\therefore$ $\frac{\sin h}{h}$ and $\frac{\tan h}{h}$ also respectively $= 1$, if $h = 0$.

30. Let $u = \sin x$; find $\frac{du}{dx}$.

For x, put $x + h$; $\therefore$ u becomes $u + \frac{du}{dx} + Uh^2$,

and $u + \frac{du}{dx}h + Uh^2 = \sin(x + h)$,

and $u \qquad\qquad\qquad = \sin x$;

$$\therefore \frac{du}{dx}h + Uh^2 = \sin(x + h) - \sin x$$

$$= 2\cos\left(x + \frac{h}{2}\right).\sin\frac{h}{2}\text{*}.$$

* Since $\sin A - \sin B = 2\cos.\frac{A+B}{2}.\sin\frac{A-B}{2}$.

$$\therefore \frac{du}{dx} + Uh = \cos\left(x + \frac{h}{2}\right) \cdot \frac{\sin\frac{h}{2}}{\frac{h}{2}};$$

and making $h = 0$; $\therefore \dfrac{\sin\frac{h}{2}}{\frac{h}{2}} = 1$;

$$\therefore \frac{du}{dx} = \cos x, \text{ or } \frac{d \cdot \sin x}{dx} = \cos x.$$

31. $u = \cos x$; find $\dfrac{du}{dx}$;

$$\therefore u + \frac{du}{dx}h + Uh^2 = \cos(x + h);$$

$$\therefore \frac{du}{dx}h + Uh^2 = \cos(x + h) - \cos x$$

$$= -2\sin\left(x + \frac{h}{2}\right)\sin\frac{h}{2};$$

$$\therefore \frac{du}{dx} + Uh = -\sin\left(x + \frac{h}{2}\right)\frac{\sin\frac{h}{2}}{\frac{h}{2}};$$

$\therefore$ making $h = 0$, $\dfrac{du}{dx} = \dfrac{d \cdot \cos x}{dx} = -\sin x$.

32. $u = \tan x$; find $\dfrac{du}{dx}$;

$$\therefore u + \frac{du}{dx}h + Uh^2 = \tan(x + h);$$

$$\therefore \frac{du}{dx}h + Uh^2 = \tan(x + h) - \tan x$$

$$= \frac{\tan h\,(1 + \tan^2 x)}{1 - \tan x \cdot \tan h};$$

$$\therefore \frac{du}{dx} + Uh = \frac{\tan h}{h} \cdot \frac{(1 + \tan^2 x)}{1 - \tan x \cdot \tan h};$$

make $h = 0$; $\therefore \tan h = 0$, and $\dfrac{\tan h}{h} = 1$;

$$\therefore \frac{du}{dx} = \frac{d.\tan x}{dx} = 1 + \tan^2 x = \sec^2 x = \frac{1}{\cos^2 x}.$$

33. $u = \sec x = \dfrac{1}{\cos x}$; find $\dfrac{du}{dx}$;

$$\therefore \frac{du}{dx} = \frac{-\dfrac{d.\cos x}{dx}}{(\cos x)^2} = \frac{\sin x}{(\cos x)^2} = \frac{\sin x}{\cos x}.\frac{1}{\cos x},$$

$$\text{or } \frac{d.\sec x}{dx} = \tan x.\sec x.$$

34. $u = V.\sin x = 1 - \cos x$;

$$\therefore \frac{du}{dx} = -\frac{d.\cos x}{dx} = \sin x.$$

35. $u = \operatorname{cotan} x = \dfrac{\cos x}{\sin x}$.

$$\frac{du}{dx} = \frac{\sin x.\dfrac{d.\cos x}{dx} - \cos x.\dfrac{d\sin x}{dx}}{(\sin x)^2} = -\frac{(\sin x)^2 + (\cos x)^2}{(\sin x)^2}$$

$$= -\frac{1}{(\sin x)^2} = -(\operatorname{cosec} x)^2 = -(1 + \cot^2 x).$$

36. $u = \operatorname{cosec} x = \dfrac{1}{\sin x}$;

$$\therefore \frac{du}{dx} = \frac{-\dfrac{d\sin x}{dx}}{(\sin x)^2} = \frac{-\cos x}{(\sin x)^2} = -\cot x.\operatorname{cosec} x.$$

37. Hence collecting the results,

If $u = \sin x$, $\quad \dfrac{du}{dx} = \cos x$,

$u = \cos x$, $\quad \dfrac{du}{dx} = -\sin x$,

$u = \tan x$, $\quad \dfrac{du}{dx} = 1 + \tan^2 x = \dfrac{1}{\cos^2 x}$,

$u = \sec x$, $\quad \dfrac{du}{dx} = \sec x.\tan x$,

$u = v.\sin x$; $\therefore \dfrac{du}{dx} = \sin x$,

$$u = \cot x, \qquad \frac{du}{dx} = -(1 + \cot^2 x) = -\frac{1}{\sin^2 x},$$

$$u = \operatorname{cosec} x, \qquad \frac{du}{dx} = -\operatorname{cosec} x \cdot \cot x.$$

38. Next, let $u = \sin z$, where $z = f(x)$.

$$\text{Then } \frac{du}{dx} = \frac{du}{dz} \cdot \frac{dz}{dx}.$$

$$\text{But } \frac{du}{dz} = \cos z; \quad \therefore \frac{du}{dx} = \cos z \frac{dz}{dx}.$$

39. Let $u = \cos z$; find $\frac{du}{dx}$.

$$\frac{du}{dz} = -\sin z; \quad \therefore \frac{du}{dx} = -\sin z \cdot \frac{dz}{dx}.$$

Ex. Let $u = \sin 3x$; $\quad \therefore \frac{du}{dx} = 3 \cos 3x.$

Let $u = \cos(ax + b)$; $\quad \therefore \frac{du}{dx} = -a \cdot \sin(ax + b).$

40. Let $u = \tan z$; $\quad \therefore \frac{du}{dz} = 1 + \tan^2 z$;

$$\therefore \frac{du}{dx} = (1 + \tan^2 z) \frac{dz}{dx}.$$

41. And if $u = \sec z$; $\quad \frac{du}{dx} = \sec z \cdot \tan z \cdot \frac{dz}{dx}$,

$u = v \sin z$; $\quad \frac{du}{dx} = \sin z \cdot \frac{dz}{dx}$,

$u = \cot z$; $\quad \frac{du}{dx} = -(1 + \cot^2 z) \frac{dz}{dx}$,

$u = \operatorname{cosec} z$; $\quad \frac{du}{dx} = -\operatorname{cosec} z \cdot \cot z \cdot \frac{dz}{dx}$.

42. To find the differential coefficients of the angle in terms of the sine, cosine, tangent, &c.

Before we do this it will be necessary to shew that if z be a function of x, or if $u = f(x)$, and consequently x a function of u (since it is a matter of convention which

of the two is the independent variable), or as it is written $f^{-1}(u)$, where f^{-1} is called the inverse function,

$$\frac{du}{dx} = \frac{1}{\frac{dx}{du}}.$$

Let δu, δx, be the differentials of u and x.

Then since $\frac{a}{b} = \frac{1}{\frac{b}{a}}$; $\therefore \frac{\delta u}{\delta x} = \frac{1}{\frac{\delta x}{\delta u}}$.

But since the ratio of the differentials is equal to the ratio of the differential coefficient to unity;

$$\therefore \frac{\delta u}{\delta x} = \frac{du}{dx}; \quad \frac{\delta x}{\delta u} = \frac{dx}{du}. \quad \text{And } \therefore \frac{du}{dx} = \frac{1}{\frac{dx}{du}}.$$

43. Hence*, if $u = \sin^{-1}x$, $\cos^{-1}x$, $\tan^{-1}x$, &c. find $\frac{du}{dx}$.

(1) $u = \sin^{-1}x$; $\therefore x = \sin u$;

$$\therefore \frac{dx}{du} = \cos u = \sqrt{1 - \sin^2 u} = \sqrt{1 - x^2};$$

$$\therefore \frac{du}{dx} = \frac{1}{\frac{dx}{du}} = \frac{1}{\sqrt{1 - x^2}}.$$

(2) $u = \cos^{-1}x$, or $x = \cos u$; $\therefore \frac{dx}{du} = -\sin u$;

$$\therefore \frac{du}{dx} = -\frac{1}{\sin u} = -\frac{1}{\sqrt{1 - x^2}}.$$

(3) $u = \tan^{-1}x$; $\therefore x = \tan u$; $\therefore \frac{dx}{du} = (1 + \tan^2 u)$;

$$\therefore \frac{du}{dx} = \frac{1}{1 + \tan^2 u} = \frac{1}{1 + x^2}.$$

(4) $u = \sec^{-1}x$; $\therefore x = \sec u$; $\therefore \frac{dx}{du} = \sec u . \tan u$;

$$\therefore \frac{du}{dx} = \frac{1}{\sec u \tan u} = \frac{1}{x\sqrt{x^2 - 1}}.$$

* By $u = \sin^{-1}x$ is meant, u is an angle whose sine is x. Similarly, $u = \tan^{-1}x$ is an angle u of which the tangent is x; these are called inverse functions. Thus, if $u = \log x$, then $u = \log^{-1}x$ expresses that u is a number of which the logarithm is x.

(5) $u = \cot^{-1}x$; $\therefore\ x = \cot u$; $\therefore\ \dfrac{dx}{du} = -(1+\cot^2 u)$;

$$\therefore\ \frac{du}{dx} = -\frac{1}{1+\cot^2 u} = -\frac{1}{1+x^2}.$$

(6) $u = \operatorname{cosec}^{-1}x$; $\therefore\ x = \operatorname{cosec} u$; $\therefore\ \dfrac{dx}{du} = -\operatorname{cosec} u.\cot u$;

$$\therefore\ \frac{du}{dx} = \frac{-1}{\operatorname{cosec} u \cot u} = -\frac{1}{x\sqrt{x^2-1}}.$$

(7) $u = v.\sin^{-1}x$; $\therefore\ x = v\sin.u$;

$$\therefore\ \frac{dx}{du} = \sin u = \sqrt{1-\cos^2 u} = \sqrt{(1-\cos u)(1+\cos u)}.$$

But $1-\cos u = x$; $1+\cos u = 2-x$,

$$\therefore\ \frac{dx}{du} = \sqrt{2x-x^2} \text{ and } \frac{du}{dx} = \frac{1}{\sqrt{2x-x^2}}.$$

Hence, recapitulating;

$$\therefore\ \frac{d.\sin^{-1}x}{dx} = \frac{1}{\sqrt{1-x^2}},$$

$$\frac{d.\cos^{-1}x}{dx} = \frac{-1}{\sqrt{1-x^2}},$$

$$\frac{d.\tan^{-1}x}{dx} = \frac{1}{1+x^2},$$

$$\frac{d.\sec^{-1}x}{dx} = \frac{1}{x\sqrt{x^2-1}},$$

$$\frac{d.\cot^{-1}x}{dx} = \frac{-1}{1+x^2},$$

$$\frac{d.\operatorname{cosec}^{-1}x}{dx} = \frac{-1}{x\sqrt{x^2-1}},$$

$$\frac{d.v\sin^{-1}x}{dx} = \frac{1}{\sqrt{2x-x^2}}.$$

44. Again, if $u = \sin^{-1}\dfrac{x}{a}$; $\therefore\ \dfrac{x}{a} = \sin u$;

$$\therefore\ \frac{dx}{du} = a\cos u = a\sqrt{1-\frac{x^2}{a^2}} = \sqrt{a^2-x^2};$$

$$\therefore \frac{du}{dx} = \frac{1}{\frac{dx}{du}} = \frac{1}{\sqrt{a^2 - x^2}}.$$

Similarly, if $u = \cos^{-1}\frac{x}{a}$, $\frac{du}{dx} = -\frac{1}{\sqrt{a^2 - x^2}}$.

If $u = \tan^{-1}\frac{x}{a}$; $\therefore \frac{x}{a} = \tan u$,

and $\frac{dx}{du} = a(1 + \tan^2 u) = a\left(1 + \frac{x^2}{a^2}\right) = \frac{a^2 + x^2}{a}$;

$$\therefore \frac{du}{dx} = \frac{a}{a^2 + x^2}.$$

Or $$\frac{d.\left(\sin^{-1}\frac{x}{a}\right)}{dx} = \frac{1}{\sqrt{a^2 - x^2}},$$

$$\frac{d.\cos^{-1}\left(\frac{x}{a}\right)}{dx} = \frac{-1}{\sqrt{a^2 - x^2}},$$

$$\frac{d.\left(\tan^{-1}\frac{x}{a}\right)}{dx} = \frac{a}{a^2 + x^2},$$

and similarly, $$\frac{d.\left(\sec^{-1}\frac{x}{a}\right)}{dx} = \frac{a}{x\sqrt{x^2 - a^2}}.$$

45. Also, if $u = \sin^{-1}z$, where $z = f(x)$, to find $\frac{du}{dx}$.

$$\frac{du}{dz} = \frac{1}{\sqrt{1 - z^2}}; \quad \therefore \frac{du}{dx} = \frac{du}{dz}\cdot\frac{dz}{dx} = \frac{\frac{dz}{dx}}{\sqrt{1 - z^2}}.$$

Next let $u = \cos^{-1}z$; $\therefore \frac{du}{dz} = \frac{-1}{\sqrt{1 - z^2}}$,

and $\therefore \frac{du}{dx} = \frac{-\frac{dz}{dx}}{\sqrt{1 - z^2}}$.

$$\text{Let } u = \tan^{-1} x; \quad \therefore \frac{du}{dx} = \frac{1}{1+x^2};$$

$$\therefore \frac{du}{dx} = \frac{\frac{dx}{dx}}{1+x^2},$$

and in the same manner for the other circular functions.

46. And as in the preceding Chapter we find the differentials from the differential coefficients: thus,

$$d.(\sin x) = \cos x \, . \, dx,$$

$$d.(\cos x) = -\sin x \, . \, dx,$$

$$d.(\tan x) = (1 + \tan^2 x)\, dx,$$

$$\text{and } d.(\sin^{-1} x) = \frac{dx}{\sqrt{1-x^2}},$$

$$d.(\tan^{-1} x) = \frac{dx}{1+x^2},$$

$$d.(\sec^{-1} x) = \frac{dx}{x\sqrt{x^2-1}}.$$

Ex. Find that angle (x) which increases twice as fast as its sine.

$$\text{Let } u = \sin x; \quad du = \cos x \, . \, dx.$$

$$\text{But } du = \tfrac{1}{2} dx; \quad \therefore \cos x = \tfrac{1}{2}; \quad \therefore x = 60^0.$$

EXPONENTIAL AND LOGARITHMIC FUNCTIONS.

47. Let $u = a^x$, which in general expresses the relation between a number u and its logarithm x, find $\frac{du}{dx}$.

We must first prove that,

$$a^x = 1 + Ax + Bx^2 + \&c.$$

$$\text{where } A = (a-1) - \tfrac{1}{2}(a-1)^2 + \tfrac{1}{3}(a-1)^3 - \&c.$$

$$\text{Now } a^x = (1 + a - 1)^x = (1+b)^x, \text{ if } b = a - 1.$$

$$\text{and } (1+b)^x = 1 + xb + \frac{x(x-1)}{2} b^2 + \frac{x(x-1)(x-2)}{2.3} b^3 + \&c.$$

$$= 1 + xb + \left(\frac{x^2}{2} - \frac{x}{2}\right) b^2 + \left(\frac{x^3}{6} - \frac{x^2}{2} + \frac{x}{3}\right) b^3 + \&c.$$

$$= 1 + \left(b - \frac{b^2}{2} + \frac{b^3}{3} - \&c.\right) x + \left(\frac{b^2}{2} - \frac{b^3}{2} + \&c.\right) x^2 + \&c.$$

$$= 1 + Ax + Bx^2 + \&c.$$

where $A = b - \frac{b^2}{2} + \frac{b^3}{3} - \&c. = (a-1) - \frac{1}{2}(a-1)^2 + \&c.$

and B is also a function of $(a-1)$;

$$\therefore u + \frac{du}{dx}h + Uh^2 = a^{x+h} = a^x . a^h$$

$$= a^x . \{1 + Ah + Bh^2 + \&c.\};$$

$$\therefore \frac{du}{dx} = A . a^x; \quad \text{and } du = A . a^x . dx.$$

Cor. There is a value e of a which makes $A = 1$,

or $(e-1) - \frac{1}{2}(e-1)^2 + \frac{1}{3}(e-1)^3 - \&c. = 1$;

$$\therefore \frac{d . e^x}{dx} = e^x; \quad \text{and } d . e^x = e^x dx;$$

e is found to be $= 2.71828$, &c. and is the base of the Napierian system of logarithms.

48. Next let $u = \log x$; $\therefore x = a^u$; $\therefore \frac{dx}{du} = Aa^u = A . x$;

$$\therefore \frac{du}{dx} = \frac{1}{\frac{dx}{du}} = \frac{1}{A} . \frac{1}{x}.$$

If the base be (e), $A = 1$ and $\frac{du}{dx} = \frac{1}{x}$, or $d . \log x = \frac{dx}{x}$.

49. Again, if $u = a^z$, find $\frac{du}{dx}$.

$$\frac{du}{dz} = Aa^z, \quad \therefore \frac{du}{dx} = \frac{du}{dz} . \frac{dz}{dx} = A . a^z . \frac{dz}{dx}.$$

Cor. If $A = 1$, $\therefore a = e$, $\therefore \frac{d . e^x}{dx} = e^x . \frac{dz}{dx}$.

50. And if $u = \log(x)$, find $\frac{du}{dx}$.

$$\frac{du}{dz} = \frac{1}{A}\cdot\frac{1}{z}; \quad \therefore \frac{du}{dx} = \frac{du}{dz}\cdot\frac{dz}{dx} = \frac{1}{A}\cdot\frac{1}{z}\cdot\frac{dz}{dx}.$$

$$\text{If } A = 1, \frac{d.(\log z)}{dx} = \frac{\frac{dz}{dx}}{z}; \text{ and } d.(\log z) = \frac{dz}{z}.$$

From the former of which equations, we obtain this rule:

The differential coefficient of the logarithm of any function is equal to the differential coefficient of the function, divided by the function itself.

Observe in future, whenever *log* is used, the Napierian logarithm is meant.

EXAMPLES.

(1) $u = (\sin x)^n$; $\frac{du}{dx} = n(\sin x)^{n-1}\cos x$.

(2) $u = \sin nx$; $\frac{du}{dx} = n\cos nx$.

(3) $u = (\tan x)^3$; $\frac{du}{dx} = 3\tan^2 x \sec^2 x$.

(4) $u = \sin 3x \cdot \cos 2x$,

$$\frac{du}{dx} = 3\cos 3x\cos 2x - 2\sin 3x\cdot\sin 2x$$
$$= \cos 3x\cos 2x + 2(\cos 3x\cos 2x - \sin 3x\sin 2x)$$
$$= \cos 3x\cos 2x + 2\cos 5x.$$

(5) $u = \sin(\cos x) = \sin z$, if $z = \cos x$;

$$\therefore \frac{du}{dx} = \frac{du}{dz}\cdot\frac{dz}{dx} = \cos z\cdot(-\sin x) = -\sin x\cos(\cos x).$$

(6) $u = \sin^{-1}\frac{x}{\sqrt{1+x^2}} = \sin^{-1} z$, if $z = \frac{x}{\sqrt{1+x^2}}$,

$$\frac{du}{dx} = \frac{\frac{dz}{dx}}{\sqrt{1-z^2}},$$

$$\frac{dz}{dx} = \frac{\sqrt{1+x^2} - \frac{x^2}{\sqrt{1+x^2}}}{1+x^2} = \frac{1}{(1+x^2)^{\frac{3}{2}}},$$

$$\sqrt{1-z^2} = \sqrt{1-\frac{x^2}{1+x^2}} = \frac{1}{\sqrt{1+x^2}},$$

$$\therefore \frac{du}{dx} = \frac{1}{1+x^2}.$$

(7) $u = \log(x + \sqrt{1+x^2}) = \log z,$

$$\therefore \frac{du}{dx} = \frac{\frac{dz}{dx}}{z}, \text{ and } z = x + \sqrt{1+x^2};$$

$$\therefore \frac{dz}{dx} = 1 + \frac{x}{\sqrt{1+x^2}} = \frac{\sqrt{1+x^2}+x}{\sqrt{1+x^2}} = \frac{z}{\sqrt{1+x^2}};$$

$$\therefore \frac{du}{dx} = \frac{1}{\sqrt{1+x^2}}.$$

(8) $u = \log(x + 1 + \sqrt{2x+x^2}); \quad \therefore \frac{du}{dx} = \frac{1}{\sqrt{2x+x^2}}.$

(9) $u = \log \frac{\sqrt{x^2+1} - x}{\sqrt{x^2+1} + x}; \quad \frac{du}{dx} = -\frac{2}{\sqrt{x^2+1}}.$

(10) $u = \log \frac{x}{\sqrt{x^2+1}+x} = \frac{du}{dx} = \frac{1}{x} - \frac{1}{\sqrt{x^2+1}}.$

(11) $u = \log \frac{x}{\sqrt{x^2+1}+1}; \quad \frac{du}{dx} = \frac{1}{x\sqrt{x^2+1}}.$

(12) $u = (\log x)^n; \quad \therefore \frac{du}{dx} = n \cdot (\log x)^{n-1} \cdot \frac{1}{x}.$

(13) $u = \log(\log x) = \log z;$

$$\therefore \frac{du}{dx} = \frac{du}{dz} \cdot \frac{dz}{dx} = \frac{1}{z} \cdot \frac{1}{x} = \frac{1}{x \log x}.$$

(14) $u = x^{f(x)} = x^z$, where $z = f(x)$.

$$\log u = z \log x; \quad \therefore \frac{du}{dx} \cdot \frac{1}{u} = \frac{dz}{dx} \log x + z \cdot \frac{1}{x};$$

$$\therefore \frac{du}{dx} = x^z \left\{ \frac{z}{x} + \frac{dz}{dx} . \log x \right\}.$$

If $z = x$, $\frac{du}{dx} = x^x \{1 + \log x\} = x^x \log (ex)$.

(15) $u = z^v$, z and v being functions of x.

$$\log u = v \log z; \quad \therefore \frac{du}{dx} . \frac{1}{u} = \frac{dv}{dx} . \log z + v . \frac{dz}{dx} . \frac{1}{z};$$

$$\therefore \frac{du}{dx} = z^v \left\{ \frac{dv}{dx} \log z + \frac{v}{z} . \frac{dz}{dx} \right\}.$$

Let $z = \sin x$, and $v = \cos x$;

$$\therefore \frac{du}{dx} = \overline{\sin x}^{\cos x} \left\{ - \sin x \log \sin x + \frac{\cos^2 x}{\sin x} \right\}.$$

(16) $u = e^{e^x} = e^z$, if $z = e^x$;

$$\therefore \frac{du}{dx} = e^z . \frac{dz}{dx} = e^z . e^x = e^{e^x} e^x.$$

(17) $u = z^{v^y}$, where z, v, and y are functions of x.

Let $v^y = v_1$; $\therefore u = z^{v_1}$,

$$\text{and } \frac{du}{dx} = z^{v_1} \left\{ \log z . \frac{dv_1}{dx} + \frac{v_1}{z} . \frac{dz}{dx} \right\}.$$

But $\because v_1 = v^y$; $\therefore \frac{dv_1}{dx} = v^y \left\{ \log v . \frac{dy}{dx} + \frac{y}{v} . \frac{dv}{dx} \right\}$;

$$\therefore \frac{du}{dx} = z^{v^y} \left\{ v^y . \log z \left(\log v . \frac{dy}{dx} + \frac{y}{v} . \frac{dv}{dx} \right) + \frac{v^y}{z} . \frac{dz}{dx} \right\}$$

$$= z^{v^y} . v^y \left\{ \log z . \log v . \frac{dy}{dx} + \frac{y}{v} \log z . \frac{dv}{dx} + \frac{1}{z} . \frac{dz}{dx} \right\}.$$

(18) $u = \log \tan x$;

$$\therefore \frac{du}{dx} = \frac{1 + \tan^2 x}{\tan x} = \frac{\sec^2 x}{\tan x} = \frac{1}{\sin x \cos x} = \frac{2}{\sin 2x}.$$

(19) $u = \log \sqrt{\frac{1 + \sin x}{1 - \sin x}}$

$$= \tfrac{1}{2} \{ \log (1 + \sin x) - \log (1 - \sin x) \};$$

$$\therefore \frac{du}{dx} = \tfrac{1}{2}\left(\frac{\cos x}{1+\sin x} + \frac{\cos x}{1-\sin x}\right) = \frac{\cos x}{1-\sin^2 x} = \frac{1}{\cos x}.$$

(20) $u = \sin^{-1}(3x - 4x^3)$;

$$\therefore x = \sin\frac{u}{3}; \quad u = 3\sin^{-1}x;$$

$$\therefore \frac{du}{dx} = \frac{3}{\sqrt{1-x^2}}.$$

(21) $u = \log(\cos x + \sqrt{-1}\sin x)$;

$$\therefore \frac{du}{dx} = \frac{-\sin x + \sqrt{-1}\cos x}{\cos x + \sqrt{-1}\sin x} = \sqrt{-1}\,\frac{\cos x + \sqrt{-1}\sin x}{\cos x + \sqrt{-1}\sin x}$$

$$= \sqrt{-1}.$$

(22) $u = \dfrac{1}{\sqrt{a^2-b^2}} \cdot \cos^{-1}\left(\dfrac{b + a\cos x}{a + b\cos x}\right)$.

$$\text{Let } z = \frac{b + a\cos x}{a + b\cos x};$$

$$\therefore \frac{du}{dx} = \frac{1}{\sqrt{a^2-b^2}} \cdot \frac{-\dfrac{dz}{dx}}{\sqrt{1-z^2}}.$$

$$\text{But } -\frac{dz}{dx} = \frac{a\sin x\,.\,(a + b\cos x) - b\sin x\,(b + a\cos x)}{(a + b\cos x)^2}$$

$$= \frac{(a^2-b^2)\,.\,\sin x}{(a + b\cos x)^2},$$

$$1 - z^2 = \frac{(a + b\cos x)^2 - (b + a\cos x)^2}{(a + b\cos x)^2}$$

$$= \frac{(a^2-b^2) - (a^2-b^2)\cos^2 x}{(a + b\cos x)^2} = \frac{(a^2-b^2)\sin^2 x}{(a + b\cos x)^2};$$

$$\therefore \sqrt{1-z^2} = \frac{\sqrt{a^2-b^2}\,.\,\sin x}{a + b\cos x};$$

$$\therefore \frac{-\dfrac{dz}{dx}}{\sqrt{1-z^2}} = \frac{\sqrt{a^2-b^2}}{a + b\,.\,\cos x};$$

$$\therefore \frac{du}{dx} = \frac{1}{\sqrt{a^2 - b^2}} \cdot \frac{-\frac{dz}{dx}}{\sqrt{1 - z^2}} = \frac{1}{a + b \cdot \cos x}.$$

(23) $u = e^x(x^4 - 4x^3 + 12x^2 - 24x + 24)$; $\frac{du}{dx} = x^4 \cdot e^x$.

(24) $u = \sin^3 x \cos x$; $\frac{du}{dx} = \sin^2 x (3 - 4 \sin^2 x)$.

(25) $u = \frac{\cos x}{\cos 2x}$; $\frac{du}{dx} = \frac{\sin x (2 + \cos 2x)}{(\cos 2x)^2}$.

(26) $u = e^x \cdot \cos x$; $\frac{du}{dx} = e^x (\cos x - \sin x)$.

(27) $u = \cos^{-1}(4x^3 - 3x)$; $\frac{du}{dx} = \frac{-3}{\sqrt{1 - x^2}}$.

(28) $u = \tan^{-1} \cdot \frac{2x}{1 - x^2}$; $\frac{du}{dx} = \frac{2}{1 + x^2}$.

(29) $u = e^{\sin x} \cdot \cos x$; $\frac{du}{dx} = e^{\sin x}(1 - \sin x - \sin^2 x)$.

(30) $u = \log \sqrt{\frac{1 - \cos x}{1 + \cos x}}$; $\frac{du}{dx} = \frac{1}{\sin x}$.

(31) $u = \sin^{-1}(2x - 1)$; $\frac{du}{dx} = \frac{1}{\sqrt{x - x^2}}$.

(32) $u = (\sin x)^x$; $\frac{du}{dx} = (\sin x)^x \{\log \sin x + x \cot x\}$.

(33) $u = \frac{x^4 (\log x)^2}{4} - \frac{x^4 \log x}{8} + \frac{x^4}{32}$; $\frac{du}{dx} = x^3 (\log x)^2$.

(34) $u = \tan^{-1}(\sqrt{1 + x^2} - x)$; $\frac{du}{dx} = -\frac{1}{2} \cdot \frac{1}{1 + x^2}$.

(35) $u = \log \sqrt{\sin x} + \log \sqrt{\cos x}$; $\frac{du}{dx} = \cot 2x$.

(36) $u = \sqrt{1 - x^2} + \sin^{-1} x$; $\frac{du}{dx} = \sqrt{\frac{1 - x}{1 + x}}$.

(37) $u = \frac{1}{\sqrt{2}} \sin^{-1} \cdot \frac{x\sqrt{2}}{1 + x^2}$; $\frac{du}{dx} = \frac{1 - x^2}{1 + x^2} \cdot \frac{1}{\sqrt{1 + x^4}}$.

(38) $u = x^{\frac{1}{x}}$; $\dfrac{du}{dx} = \dfrac{x^{\frac{1}{x}}}{x^2} \log\left(\dfrac{e}{x}\right)$.

(39) $u = \log \dfrac{(x+2)^2}{\sqrt{x+1}.(x+3)^{\frac{1}{2}}}$; $\dfrac{du}{dx} = \dfrac{x}{x^3+6x^2+11x+6}$.

(40) $u = \log \sqrt[4]{\dfrac{1+x}{1-x}} + \frac{1}{2}\tan^{-1}x$; $\dfrac{du}{dx} = \dfrac{1}{1-x^4}$.

(41) $u = \dfrac{e^{ax}(a\sin x - \cos x)}{a^2+1}$; $\dfrac{du}{dx} = e^{ax}.\sin x$.

(42) $u = a\log\left(\dfrac{a+\sqrt{a^2-x^2}}{x}\right) - \sqrt{a^2-x^2}$; $\dfrac{du}{dx} = \dfrac{-\sqrt{a^2-x^2}}{x}$.

(43) $u = a(\sin x - \cos x)$; $\dfrac{du}{dx} = \sqrt{2a^2-u^2}$.

(44) $u = \dfrac{e^x+e^{-x}}{e^x-e^{-x}}$; $\dfrac{du}{dx} = 1-u^2$.

(45) $u = xe^{\tan^{-1}x}$; $\dfrac{du}{dx} = e^{\tan^{-1}x}\left(\dfrac{1+x+x^2}{1+x^2}\right)$.

(46) $u = \sin(\log x)$; $\dfrac{du}{dx} = \dfrac{1}{x}\cos(\log x)$.

(47) $u = a(z+\sin z)$; $x = a(1-\cos z)$;

$$\frac{du}{dx} = \frac{\sqrt{2ax-x^2}}{x}.$$

(48) $u = e^x\sqrt{\dfrac{1+x}{1-x}}$; $\dfrac{du}{dx} = e^x.\dfrac{2-x^2}{(1-x)^{\frac{1}{2}}\sqrt{1+x}}$.

(49) $u = -\frac{1}{3}\dfrac{\sec x}{(\sin x)^3} - \frac{8}{3}\cot 2x$; $\dfrac{du}{dx} = \dfrac{1}{(\sin x)^4(\cos x)^2}$.

(50) $u = \sin^{-1}(2x\sqrt{1-x})^2$; $\dfrac{du}{dx} = \dfrac{2}{\sqrt{1-x^2}}$.

CHAPTER III.

SUCCESSIVE DIFFERENTIATION. MACLAURIN'S THEOREM.

---

51. If $u = f(x)$; the differential coefficient $\frac{du}{dx}$ may also be a function of x, and therefore be capable of being differentiated.

For suppose that $\frac{du}{dx} = p$, and for x put $x + h$,

Then $\frac{du}{dx} + \frac{d.\left(\frac{du}{dx}\right)}{dx} h + \&c. = p + \frac{dp}{dx} h + \&c.$ (by writing $\frac{du}{dx}$ for u, in $u + \frac{du}{dx} h + Uh^2$);

$$\therefore \frac{d.\left(\frac{du}{dx}\right)}{dx} = \frac{dp}{dx}.$$

In the same manner if $\frac{dp}{dx}$ be a function of x and be put equal to q, we shall have

$$\frac{dq}{dx} = \frac{d.\left(\frac{dp}{dx}\right)}{dx} = \frac{d}{dx}.\left(\frac{d.\left(\frac{du}{dx}\right)}{dx}\right);$$

and so on for other differential coefficients.

This process is called successive differentiation, and $\frac{du}{dx}$, $\frac{dp}{dx}$, $\frac{dq}{dx}$, &c. are called the first, second, third, &c. differential coefficients.

A more convenient notation than that above is used, the reason for which may be derived from the consideration of differentials; and here we may remark that dx as well as h, is always considered to be invariable, when $u = f(x)$: this being observed;

Since $\frac{du}{dx} = p$; $\therefore du = p dx$;

$$\therefore\ d(du) = dp\,dx\text{: but } dp = q\,dx;$$

$$\therefore\ d(du) = q\,dx^2.$$

But because $d(du)$ is the symbol for shewing that u has been twice differentiated, and since d is the symbol of differentiation; therefore d^2u will fitly express the fact of u being thus twice operated upon;

$$\therefore\ d^2u = q\,dx^2.$$

Similarly $d(d^2u) = d^3u = dq\,dx^2 = r\,dx^3$,

$d(d^3u) = d^4u = dr\,dx^3 = s\,dx^4$,

&c.

Hence therefore it follows, that

$$\frac{dp}{dx} = q = \frac{d^2u}{dx^2},$$

$$\frac{dq}{dx} = r = \frac{d^3u}{dx^3},$$

$$\frac{dr}{dx} = s = \frac{d^4u}{dx^4};$$

and the n^{th} differential coefficient is $\frac{d^nu}{dx^n}$.

Observe $\frac{d^2u}{dx^2}$, $\frac{d^3u}{dx^3}$, &c. are most commonly read, second du by dx squared, third du by dx cubed; but sometimes, d two u, by dx squared, d three u, by dx cubed, &c.

Ex. 1. Let $u = x^4 + x^3 + x^2 + x + 1$,

$$\frac{du}{dx} = 4x^3 + 3x^2 + 2x + 1,$$

$$\frac{d^2u}{dx^2} = 3 \cdot 4x^2 + 2 \cdot 3x + 2,$$

$$\frac{d^3u}{dx^3} = 2 \cdot 3 \cdot 4x + 2 \cdot 3,$$

$$\frac{d^4u}{dx^4} = 2 \cdot 3 \cdot 4,$$

$$\frac{d^5u}{dx^5} = 0.$$

Ex. 2. Let $u = \frac{1}{x} = x^{-1}$.

$$\frac{du}{dx} = -x^{-2} = -\frac{1}{x^2},$$

$$\frac{d^2u}{dx^2} = 2x^{-3} = \frac{2}{x^3},$$

$$\frac{d^3u}{dx^3} = -2.3x^{-4} = -\frac{2.3}{x^4},$$

$$\frac{d^4u}{dx^4} = 2.3.4x^{-5} = \frac{2.3.4}{x^5},$$

$$\vdots \qquad \vdots \qquad \vdots$$

$$\frac{d^nu}{dx^n} = (-1)^n 2.3.4.5\ldots n.x^{-(n+1)}.$$

Ex. 3. Let $u = \sin(ax + b)$; find $\frac{d^4u}{dx^4}$.

$$\therefore \frac{du}{dx} = a\cos(ax + b),$$

$$\frac{d^2u}{dx^2} = -a^2\sin(ax + b),$$

$$\frac{d^3u}{dx^3} = -a^3\cos(ax + b),$$

$$\frac{d^4u}{dx^4} = a^4\sin(ax + b) = a^4u.$$

Ex. 4. Let $u = e^{ax}$; find $\frac{d^nu}{dx^n}$.

$$\therefore \frac{du}{dx} = ae^{ax}; \quad \frac{d^2u}{dx^2} = a^2e^{ax} = a^2u,$$

$$\frac{d^3u}{dx^3} = a^2\frac{du}{dx} = a^3e^{ax} = a^3u,$$

$$\frac{d^4u}{dx^4} = a^3\frac{du}{dx} = a^4e^{ax} = a^4u;$$

$$\therefore \frac{d^nu}{dx^n} = a^ne^{ax} = a^nu.$$

Ex. 5. Let $u = xv$; to find d^2u, d^3u, &c.

$$\because du = xdv + vdx \ldots\ldots\ldots(1);$$

$$\therefore d^2u = d(xdv) + d(vdx).$$

But from (1), $d(zdv) = zd^2v + dzdv$,

$$d(vdz) = vd^2z + dvdz;$$

$$\therefore\ d^2u = zd^2v + 2dzdv + vd^2z;$$

$$\therefore\ d^3u = d(zd^2v) + 2.d(dzdv) + d(vd^2z).$$

But $d(z.d^2v) = zd^3v + dzd^2v$,

$$2d(dzdv) = 2d^2zdv + 2dzd^2v,$$

$$d.(vd^2z) = vd^3z + dvd^2z;$$

$$\therefore\ d^3u = zd^3v + 3dzd^2v + 3dvd^2z + vd^3z,$$

and $d^4u = zd^4v + 4dzd^3v + 6d^2zd^2v + 4d^3zdv + vd^4z.$

Since the law of the numerical coefficients is apparently that of the coefficients of $(a + b)^n$;

$$\therefore\ d^nu = zd^nv + n.dzd^{n-1}v + n.\frac{(n-1)}{2}d^2zd^{n-2}v + \&c.;$$

$$\therefore\ \frac{d^nu}{dx^n} = z.\frac{d^nv}{dx^n} + n\frac{dz}{dx}.\frac{d^{n-1}v}{dx^{n-1}} + n.\frac{(n-1)}{2}.\frac{d^2z}{dx^2}.\frac{d^{n-2}v}{dx^{n-2}} + \&c.$$

a theorem due to Leibnitz, and which may be used to find the differential coefficient of the product of two functions.

Proof of the law of the coefficients in Leibnitz's Theorem. Assume that,

$$d^nu = zd^nv + ndzd^{n-1}v + n\frac{n-1}{2}.d^2zd^{n-2}v + \&c.;$$

$$\therefore\ d^{n+1}u = zd^{n+1}v + dzd^nv + n.(dzd^nv + d^2zd^{n-1}v)$$

$$+ n\frac{n-1}{2}.(d^2zd^{n-1}v + d^3z.d^{n-2}v) + \&c.$$

$$= zd^{n+1}v + (n+1).dzd^nv + \frac{n.(n+1)}{2}.d^2z.d^{n-1}v + \&c.,$$

which shews that if the theorem be true for n, it is true for $n + 1$, and it has been shewn to be true for $n = 2$, and $n = 3$; it is $\therefore$ true when $n = 4$; and $\therefore$ when n is any integer.

EXPANSION OF FUNCTIONS.

52. If $u = f(x)$ can be expanded into a series of the form

$$u = A + Bx + Cx^2 + Dx^3 + Ex^4, \&c.$$

where A, B, C, &c. are constant, to find these coefficients.

This is Maclaurin's or Stirling's Theorem.

Since $u = A + Bx + Cx^2 + Dx^3 + Ex^4 + \&c.$

Then by successive differentiation we have

$$\frac{du}{dx} = B + 2Cx + 3Dx^2 + 4E.x^3 + \&c.$$

$$\frac{d^2u}{dx^2} = 2C + 2.3Dx + 3.4E.x^2 + \&c.$$

$$\frac{d^3u}{dx^3} = 2.3.D + 2.3.4E.x + \&c.$$

$$\frac{d^4u}{dx^4} = 2.3.4E + \&c.$$

$$\&c. = \&c.$$

Make $x = 0$ in these several equations, and let U_0, U_1, U_2, U_3, &c. represent the values of u, $\frac{du}{dx}$, $\frac{d^2u}{dx^2}$, &c. on that supposition;

$$\therefore U_0 = A; \quad U_1 = B; \quad U_2 = 2C; \quad \therefore C = \frac{U_2}{1.2},$$

$$U_3 = 2.3D; \quad \therefore D = U_3\frac{1}{2.3},$$

$$E = U_4.\frac{1}{2.3.4}, \quad \&c. = \&c.;$$

$$\therefore u = U_0 + U_1x + U_2.\frac{x^2}{1.2} + U_3.\frac{x^3}{2.3} + U_4.\frac{x^4}{2.3.4} + \&c.$$

Cor. The general term is obviously $U_n\frac{x^n}{1.2.3...n}$.

EXAMPLES OF THE EXPANSION OF FUNCTIONS.

(1) Let $u = (x + a)^4$; $\quad \therefore U_0 = a^4$; if $x = 0$;

$$\frac{du}{dx} = 4(x + a)^3; \quad \therefore U_1 = 4a^3,$$

$$\frac{d^2u}{dx^2} = 4.3(x + a)^2; \quad \therefore U_2 = 3.4a^2,$$

$$\frac{d^3u}{dx^3} = 2.3.4(x + a); \quad \therefore U_3 = 2.3.4a,$$

$$\frac{d^4u}{dx^4} = 2.3.4; \qquad \therefore\ U_4 = 2.3.4,$$

$$\frac{d^5u}{dx^5} = 0; \quad \therefore\ U_5 = 0, \text{ and } U_6,\ U_7, \text{ \&c. each } = 0;$$

$$\therefore\ u = (x+a)^4 = a^4 + 4a^3x + \frac{3.4}{1.2}a^2x^2 + \frac{2.3.4}{1.2.3}ax^3 + \frac{2.3.4}{2.3.4}x^4$$

$$= a^4 + 4a^3x + 6a^2x^2 + 4ax^3 + x^4.$$

(2) Expand $(a + bx + cx^2)^n$.

$$u = (a + bx + cx^2)^n; \quad \therefore\ U_0 = a^n,$$

$$\frac{du}{dx} = n.(a + bx + cx^2)^{n-1}(b + 2cx); \quad \therefore\ U_1 = nba^{n-1},$$

$$\frac{d^2u}{dx^2} = n(n-1)(a+bx+cx^2)^{n-2}(b+2cx)^2 + 2cn.(a+bx+cx^2)^{n-1};$$

$$\therefore\ U_2 = n.(n-1)a^{n-2}b^2 + n.2c.a^{n-1},$$

$$\frac{d^3u}{dx^3} = n.(n-1).(n-2)(a+bx+cx^2)^{n-3}(b+2cx)^3$$

$$+ 2n.(n-1)(a+bx+cx^2)^{n-2}2c.(b+2cx)$$

$$+ 2c.n(n-1).(a+bx+cx^2)^{n-2}(b+2cx);$$

$$\therefore\ U_3 = n(n-1)(n-2).a^{n-3}b^3 + 2.3n.(n-1)a^{n-2}bc,$$

&c. = &c.

$$\therefore\ (a+bx+cx^2)^n = a^n + na^{n-1}bx + \left\{n.\frac{(1)}{2}.a^{n-2}b^2 + na^{n-1}c\right\}x^2$$

$$+ \left\{\frac{n.(n-1)(n-2)}{1.2.3}a^{n-3}b^3 + n.(n-1).a^{n-2}bc\right\}x^3$$

+ &c.

(3) Expand $\sin x$ and $\cos x$ in terms of x.

If $u = \sin x$,	If $u = \cos x$,
then $\frac{du}{dx} = \cos x$,	$\frac{du}{dx} = -\sin x$,
$\frac{d^2u}{dx^2} = -\sin x$,	$\frac{d^2u}{dx^2} = -\cos x$,
$\frac{d^3u}{dx^3} = -\cos x$,	$\frac{d^3u}{dx^3} = \sin x$,
$\frac{d^4u}{dx^4} = +\sin x$,	$\frac{d^4u}{dx^4} = \cos x$,
&c. = &c.	&c. = &c.

After the 4th differentiation the values of the differential coefficients recur.

Now make $x = 0$, then in the series for sin x,

$U_0 = 0$, $U_1 = 1$, $U_2 = 0$, $U_3 = -1$, $U_4 = 0$, $U_5 = +1$, &c.

and for cos x,

$$U_0 = 1, \quad U_1 = 0, \quad U_2 = -1, \quad U_3 = 0, \quad U_4 = 1, \text{ \&c.}$$

$$\therefore \sin x = x - \frac{x^3}{2.3} + \frac{x^5}{2.3.4.5} - \text{\&c.}$$

$$\text{and } \cos x = 1 - \frac{x^2}{1.2} + \frac{x^4}{2.3.4} - \text{\&c.}$$

Cor. The series for cos x may be derived from that of sin x by differentiating the latter.

(4) Similarly, if $u = \tan x$, we may find tan x in terms of x.

But more readily in the following manner*.

Let $u = \tan x = a_1 x + a_3 x^3 + a_5 x^5 + a_7 x^7 + \text{\&c.}$;

$$\therefore \frac{du}{dx} = 1 + \tan^2 x = a_1 + 3a_3 x^2 + 5a_5 x^4 + 7a_7 x^6 + \text{\&c.}$$

But $1 + \tan^2 x = 1 + (a_1 x + a_3 x^3 + a_5 x^5 + \text{\&c.})^2$

$$= 1 + a_1^2 x^2 + 2a_1 a_3 x^4 + (a_3^2 + 2a_1 a_5) x^6 + \text{\&c.};$$

therefore equating coefficients of the like powers of x,

$$a_1 = 1, \quad 3a_3 = a_1^2; \quad \therefore a_3 = \frac{1}{3},$$

$$5a_5 = 2a_1 a_3 = \frac{2}{3}; \quad \therefore a_5 = \frac{2}{3.5},$$

$$7a_7 = a_3^2 + 2a_1 a_5 = \frac{1}{9} + \frac{4}{3.5} = \frac{17}{5.9}; \quad \therefore a_7 = \frac{17}{5.7.9};$$

$$\therefore u = x + \frac{x^3}{1.3} + \frac{2x^5}{3.5} + \frac{17x^7}{5.7.9} + \text{\&c.}$$

* That tan x can only involve odd powers of (x) may be thus shewn:

Let $\tan x = a_1 x + b_2 x^2 + a_3 x^3 + b_4 x^4 + a_5 x^5 + \text{\&c.}$;

$\therefore \tan(-x) = -a_1 x + b_2 x^2 - a_3 x^3 + b_4 x^4 - a_5 x^5 + \text{\&c.}$;

$\therefore \tan x - \tan(-x) = 2a_1 x + 2a_3 x^3 + 2a_5 x^5 + \text{\&c.}$

But $\tan(-x) = -\tan(x)$; $\therefore \tan x - \tan(-x) = 2\tan x$;

$\therefore \tan x = a_1 x + a_3 x^3 + a_5 x^5 + \text{\&c.}$

(5) In a similar manner may $\sin x$ be found.

Let $\sin x = a_1 x + a_3 x^3 + a_5 x^5 + \&c.$;

$$\therefore \cos x = a_1 + 3a_3 x^2 + 5a_5 x^4 + \&c.;$$

$$\therefore -\sin x = 2.3a_3 x + 4.5a_5 x^3 + \&c.$$

$$= -a_1 x - a_3 x^3 - a_5 x^5 - \&c.;$$

$$\therefore a_3 = -\frac{a_1}{2.3}; \quad a_5 = -\frac{a_3}{4.5} = \frac{a_1}{2.3.4.5};$$

$$\therefore \sin x = a_1\left(x - \frac{x^3}{2.3} + \frac{x^5}{2.3.4.5} - \&c.\right),$$

where a_1 is to be found, but

$$\because \frac{\sin x}{x} = a_1\left(1 - \frac{x^2}{2.3} + \&c.\right),$$

and if $x = 0$; $\quad \frac{\sin x}{x} = 1$; $\quad \therefore a_1 = 1$;

$$\therefore \sin x = x - \frac{x^3}{2.3} + \frac{x^5}{2.3.4.5} - \&c.$$

(6) $u = \sin^{-1} x$, whence if $x = 0$, $U_0 = \sin^{-1} 0 = 0$,

and $\frac{du}{dx} = \frac{1}{\sqrt{1-x^2}} = (1-x^2)^{-\frac{1}{2}} = 1 + \frac{1}{2}x^2 + \frac{1.3}{2.4}x^4 + \frac{1.3.5}{2.4.6}x^6 + \&c.$

but from Maclaurin's Theorem,

$$\frac{du}{dx} = U_1 + 2U_2 . \frac{x}{1.2} + 3U_3 . \frac{x^2}{2.3} + 4U_4 . \frac{x^3}{2.3.4} + 5U_5 . \frac{x^4}{2.3.4.5} + \&c.$$

Equating coefficients of the same powers of x,

$$U_1 = 1, \quad U_2 = 0, \quad \frac{3U_3}{2.3} = \frac{1}{2}; \quad \therefore U_3 = 1, \quad U_4 = 0,$$

$$\frac{5U_5}{2.3.4.5} = \frac{1.3}{2.4}, \quad U_5 = 1.3^2, \quad U_6 = 0,$$

$$\frac{7U_7}{2.3.4.5.6.7} = \frac{1.3.5}{2.4.6}; \quad \therefore U_7 = 1^2 . 3^2 . 5^2;$$

$$\therefore \sin^{-1} x = x + \frac{x^3}{1.2.3} + \frac{1.3^2 x^5}{1.2.3.4.5} + \frac{1^2 . 3^2 . 5^2 . x^7}{1.2.3.4.5.6.7} + \&c.$$

$$= x + \frac{1}{1.2} . \frac{x^3}{3} + \frac{1.3}{2.4} . \frac{x^5}{5} + \frac{1.3.5}{2.4.6} . \frac{x^7}{7} + \&c.$$

the general term of which is obviously

$$\frac{1.3.5 \ldots\ldots (2n-3)}{2.4.6 \ldots\ldots (2n-2)} \cdot \frac{x^{2n-1}}{2n-1}.$$

By this series, the length of a circular arc may be found; thus, let $\sin^{-1}x = 30$; $\therefore\ x = \frac{1}{2}$;

$$\therefore\ 30^0 = \frac{1}{2} + \frac{1}{1.2} \cdot \frac{1}{3 \times 8} + \frac{1.3}{2.4} \cdot \frac{1}{32 \times 5} + \&c.$$

(7) The same series may be thus obtained without the use of Maclaurin's Theorem.

Let $u = \sin^{-1}x = a_1 x + a_3 x^3 + a_5 x^5 + a_7 x^7 + \&c.$;

for that $\sin^{-1}x$ cannot contain even powers of x, may be proved by the method used in the note. (Page 41).

Differentiating

$$\frac{du}{dx} = \frac{1}{\sqrt{1-x^2}} = a_1 + 3a_3 x^2 + 5a_5 x^4 + 7a_7 x^6 + \&c.$$

$$\text{But } \frac{1}{\sqrt{1-x^2}} = 1 + \frac{1}{2}x^2 + \frac{1.3}{2.4} \cdot x^4 + \frac{1.3.5}{2.4.6} x^6 + \&c.$$

equating coefficients,

$$a_1 = 1, \quad 3a_3 = \frac{1}{2}; \quad \therefore\ a_3 = \frac{1}{2} \cdot \frac{1}{3},$$

$$5a_5 = \frac{1.3}{2.4}; \quad \therefore\ a_5 = \frac{1.3}{2.4} \cdot \frac{1}{5},$$

$$7a_7 = \frac{1.3.5}{2.4.6}; \quad \therefore\ a_7 = \frac{1.3.5}{2.4.6} \cdot \frac{1}{7};$$

$$\therefore\ \sin^{-1}x = \frac{x}{1} + \frac{1}{2} \cdot \frac{x^3}{3} + \frac{1.3}{2.4} \cdot \frac{x^5}{5} + \frac{1.3.5}{2.4.6} \cdot \frac{x^7}{7} + \&c.$$

(8) $u = \tan^{-1}x$; $x = 0$, $U_0 = \tan^{-1}0 = 0$;

$$\therefore\ \frac{du}{dx} = \frac{1}{1+x^2} = 1 - x^2 + x^4 - x^6 + \&c.$$

$$= U_1 + 2U_2 \frac{x}{2} + \frac{3U_3 x^2}{1.2.3} + \frac{4U_4 x^3}{2.3.4} + \frac{5U_5 x^4}{2.3.4.5} + \&c.$$

$$\therefore\ U_1 = 1,\ U_2 = 0,\ \frac{U_3}{2} = -1;\quad \therefore\ U_3 = -2,\ U_4 = 0,$$

$$\frac{U_5}{2.3.4} = 1;\quad \therefore\ U_5 = 2.3.4;$$

$$\therefore\ u = x - \frac{2x^3}{2.3} + \frac{2.3.4x^5}{2.3.4.5} - \&c.$$

$$\text{or } \tan^{-1}x = x - \frac{x^3}{3} + \frac{x^5}{5} - \&c.$$

a series for the arc in terms of the tangent.

53. Hence may be found approximate expressions for the length of the arc of a circle.

(1⁰) Let $\tan^{-1}x = \frac{\pi}{4};\quad \therefore\ x = \tan\frac{\pi}{4} = 1,$

$$\text{and } \frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \&c.$$

(2⁰) Since $\frac{\pi}{4} = \tan^{-1}\frac{1}{2} + \tan^{-1}\frac{1}{3},$

$$\text{and } \tan^{-1}\frac{1}{2} = \frac{1}{2} - \frac{1}{3}.\frac{1}{2^3} + \frac{1}{5.2^5} - \&c.$$

$$\tan^{-1}\frac{1}{3} = \frac{1}{3} - \frac{1}{3}.\frac{1}{3^3} + \frac{1}{5.3^5} - \&c.;$$

$$\therefore\ \frac{\pi}{4} = \left(\frac{1}{2} + \frac{1}{3}\right) - \frac{1}{3}.\left(\frac{1}{2^3} + \frac{1}{3^3}\right) + \frac{1}{5}.\left(\frac{1}{2^5} + \frac{1}{3^5}\right) - \&c.$$

(3⁰) A very convergent series is given by Machin's formula,

$$\text{that } \frac{\pi}{4} = 4\tan^{-1}.\frac{1}{5} - \tan^{-1}.\frac{1}{239}.$$

To prove this, let $A = 4\tan^{-1}\frac{1}{5} = 4a.$

$$\text{Then } \tan A = \frac{4\tan a - 4\tan^3 a}{1 - 6\tan^2 a + \tan^4 a} = \frac{\frac{4}{5} - \frac{4}{125}}{1 - \frac{6}{25} + \frac{1}{625}}$$

$$= \frac{4(125 - 5)}{625 - 150 + 1} = \frac{4 \times 120}{476} = \frac{120}{119};\quad \therefore\ > 1,$$

$$\text{and } \tan(A-45^0) = \frac{\tan A - 1}{\tan A + 1} = \frac{\frac{120}{119} - 1}{\frac{120}{119} + 1} = \frac{1}{239};$$

$$\therefore A - 45^0 = \tan^{-1}\frac{1}{239};$$

$$\therefore 45^0 = 4\tan^{-1}\frac{1}{5} - \tan^{-1}\frac{1}{239}$$

$$= 4\left\{\frac{1}{5} - \frac{1}{3}\cdot\frac{1}{5^3} + \frac{1}{5}\cdot\frac{1}{5^5} - \&c.\right\}$$

$$- \left\{\frac{1}{239} - \frac{1}{3}\cdot\frac{1}{(239)^3} + \frac{1}{5}\cdot\frac{1}{(239)^5} - \&c.\right\}.$$

54. The logarithm of x cannot be found by Maclaurin's Theorem, since if $x = 0$, U_0, U_1, U_2, &c. become infinite: but $u = \log(1+x)$ may be so found.

Suppose the logarithms to be Napierian, where $A = 1$.

$$u = \log(1+x); \quad \therefore U_0 = \log(1) = 0,$$

$$\frac{du}{dx} = \frac{1}{1+x} = 1 - x + x^2 - x^3 + x^4 - x^5 + \&c.$$

But from the theorem,

$$\frac{du}{dx} = U_1 + U_2 x + \frac{U_3 x^2}{2} + \frac{U_4 x^3}{2.3} + \frac{U_5 x^4}{2.3.4} + \&c.$$

therefore equating coefficients,

$$U_1 = 1, \quad U_2 = -1, \quad U_3 = 2, \quad U_4 = -2.3, \quad U_5 = 2.3.4;$$

$$\therefore u = \log(1+x) = x - \frac{x^2}{2} + \frac{2x^3}{2.3} - \frac{2.3x^4}{2.3.4} + \frac{2.3.4x^5}{2.3.4.5} + \&c.$$

$$= x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \frac{x^5}{5} - \&c.$$

Cor. Had a been the base, then (Art. 48.)

$\frac{du}{dx} = \frac{1}{A}\cdot\frac{1}{x+1}$, and if $\log_a$ be a log in that system,

$$\log_a(1+x) = \frac{1}{A}\left(x - \frac{x^2}{2} + \frac{x^3}{3} - \&c.\right)$$

$$= \frac{1}{A}\log_e(1+x).$$

Hence, if we know the Nap. log. we may find the log to a base a by multiplying the Nap. log. by $\frac{1}{A}$. The factor $\frac{1}{A}$ called the *modulus* will be found afterwards.

55. As the series for $\log(1+x)$ does not converge, it is useless in actual computation; but from it a number of series may be derived which are rapidly convergent.

Let $-x$ be written for x in the series for $\log(1+x)$;

$$\therefore \log(1-x) = -x - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{4} - \frac{x^5}{5} - \&c.$$

subtract this from $\log(1+x)$, then since $\log a - \log b = \log \frac{a}{b}$,

$$\log\left(\frac{1+x}{1-x}\right) = 2\left\{x + \frac{x^3}{3} + \frac{x^5}{5} + \&c.\right\},$$

$$\text{for } \frac{1+x}{1-x} \text{ put } \frac{M}{N}; \quad \therefore x = \frac{M-N}{M+N},$$

$$\text{and } \log\frac{M}{N} = 2\left\{\frac{M-N}{M+N} + \tfrac{1}{3}\left(\frac{M-N}{M+N}\right)^3 + \&c.\right\}.$$

Suppose $M = N + z$; $\therefore M + N = 2N + z$;

$$\therefore \log(N+z) = \log N + 2\left\{\frac{z}{2N+z} + \tfrac{1}{3}\frac{z^3}{(2N+z)^3} + \tfrac{1}{5}\frac{z^5}{(2N+z)^5} + \&c.\right\},$$

and finally, if $z = 1$,

$$\log(N+1) = \log N + 2\left\{\frac{1}{2N+1} + \tfrac{1}{3}\frac{1}{(2N+1)^3} + \&c.\right\},$$

a formula from which logarithms may be calculated.

Thus, since $\log 1 = 0$,

$$\log 2 = 2\left\{\frac{1}{3} + \frac{1}{3}\cdot\frac{1}{3^3} + \frac{1}{5}\cdot\frac{1}{3^5} + \&c.\right\},$$

$$\log 3 = \log 2 + 2\left\{\frac{1}{5} + \frac{1}{3}\cdot\frac{1}{5^3} + \frac{1}{5}\cdot\frac{1}{5^5} + \&c.\right\},$$

$$\log 4 = \log 3 + 2\left\{\frac{1}{7} + \frac{1}{3}\cdot\frac{1}{7^3} + \frac{1}{5}\cdot\frac{1}{7^5} + \&c.\right.$$

&c. &c.

56. Expand a^x in ascending powers of x.

$$u = a^x, \qquad x = 0; \qquad \therefore U_0 = 1;$$

$$\frac{du}{dx} = A \cdot a^x \ldots\ldots\ldots\ldots \therefore U_1 = A,$$

$$\frac{d^2u}{dx^2} = A^2 a^x \ldots\ldots\ldots\ldots \therefore U_2 = A^2,$$

$$\frac{d^3u}{dx^3} = A^3 a^x \ldots\ldots\ldots\ldots \therefore U_3 = A^3,$$

$$\vdots$$

$$\frac{d^nu}{dx^n} = A^n \cdot a^x \ldots\ldots\ldots\ldots \therefore U_n = A^n;$$

$$\therefore a^x = 1 + Ax + \frac{A^2x^2}{1.2} + \frac{A^3x^3}{2.3} + \frac{A^4x^4}{2.3.4} + \&c.$$

Cor. 1. To find A let $x = \frac{1}{A}$, or $Ax = 1$;

$$\therefore a^{\frac{1}{A}} = 1 + 1 + \tfrac{1}{2} + \frac{1}{2.3} + \&c. = 2.71828, \&c. = e;$$

$$\therefore \frac{1}{A}\log a = \log e; \quad \therefore A = \frac{\log a}{\log e} = \log_e a.$$

Cor. 2. If $a = e$; $A = \log_e e = 1$;

and $$e^x = 1 + x + \frac{x^2}{1.2} + \frac{x^3}{2.3} + \&c.$$

a remarkable expansion which must be carefully remembered.

Cor. 3. Hence $\log_a(1+x) = \frac{1}{A} \cdot \log_e(1+x) = \frac{\log_e(1+x)}{\log_e a}$;

$$\therefore \log_e(1+x) = \log_a(1+x)\log_e a.$$

Cor. 4. To compute A.

$$\because e = a^{\frac{1}{A}}; \quad \therefore e^A = a; \quad e^{-A} = \frac{1}{a}; \quad \therefore -A = \log\left(\frac{1}{a}\right).$$

$$\text{But } \log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \&c.$$

$$\text{for } 1 + x \text{ put } \frac{1}{a}; \quad \therefore x = \frac{1}{a} - 1;$$

$$\therefore -A = \log\left(\frac{1}{a}\right) = \left(\frac{1}{a} - 1\right) - \tfrac{1}{2}\left(\frac{1}{a} - 1\right)^2 + \tfrac{1}{3}\left(\frac{1}{a} - 1\right)^3 - \&c.;$$

$$\text{or } A = \left(\frac{a-1}{a}\right) + \tfrac{1}{2}\left(\frac{a-1}{a}\right)^2 + \tfrac{1}{3}\left(\frac{a-1}{a}\right)^3 + \&c.$$

Let $a = 10$.

$$A = (.9) + \tfrac{1}{2}(.9)^2 + \tfrac{1}{3}(.9)^3 + \&c. = 2.302585.....$$

and $\dfrac{1}{A} = .43429448.$

This is the number by which the Napierian logarithms are multiplied to obtain those to a base 10, or Briggs' logarithms.

57. In the expansion for e^x

$$\text{or } e^x = 1 + x + \frac{x^2}{1.2} + \frac{x^3}{2.3} + \frac{x^4}{2.3.4} + \frac{x^5}{2.3.4.5} + \&c.;$$

put successively for x, $x\sqrt{-1}$, and $-x\sqrt{-1}$;

$$\therefore e^{x\sqrt{-1}} = 1 + x\sqrt{-1} - \frac{x^2}{2} - \frac{x^3\sqrt{-1}}{2.3} + \frac{x^4}{2.3.4} + \frac{x^5\sqrt{-1}}{2.3.4.5} - \&c.$$

$$e^{-x\sqrt{-1}} = 1 - x\sqrt{-1} - \frac{x^2}{2} + \frac{x^3\sqrt{-1}}{2.3} + \frac{x^4}{2.3.4} - \frac{x^5\sqrt{-1}}{2.3.4.5} - \&c.$$

Therefore first by addition and then by subtraction,

$$e^{x\sqrt{-1}} + e^{-x\sqrt{-1}} = 2\left\{1 - \frac{x^2}{2} + \frac{x^4}{2.3.4} - \&c.\right\} = 2\cos x.$$

$$e^{x\sqrt{-1}} - e^{-x\sqrt{-1}} = 2\sqrt{-1}\left\{x - \frac{x^3}{2.3} + \&c.\right\} = 2\sqrt{-1}\sin x.$$

Again adding and dividing by 2,

$$e^{x\sqrt{-1}} = \cos x + \sqrt{-1}\sin x.$$

Also by subtraction and dividing by 2,

$$e^{-x\sqrt{-1}} = \cos x - \sqrt{-1}\sin x.$$

Cor. 1. Hence $\cos x = \dfrac{e^{x\sqrt{-1}} + e^{-x\sqrt{-1}}}{2}$,

$$\text{and } \sin x = \frac{e^{x\sqrt{-1}} - e^{-x\sqrt{-1}}}{2\sqrt{-1}};$$

$$\therefore \tan x = \frac{1}{\sqrt{-1}} \frac{e^{x\sqrt{-1}} - e^{-x\sqrt{-1}}}{e^{x\sqrt{-1}} + e^{-x\sqrt{-1}}} = \frac{1}{\sqrt{-1}}\left(\frac{e^{2x\sqrt{-1}} - 1}{e^{2x\sqrt{-1}} + 1}\right).$$

Cor. 2. These equations have been proved independently of the value of x, we may therefore put mx for x;

$$\therefore \cos mx + \sqrt{-1}\sin mx = e^{mx\sqrt{-1}} = \overline{e^{x\sqrt{-1}}}^{m}$$

$$= (\cos x + \sqrt{-1}\sin x)^m,$$

the formula of De Moivre.

58. We have seen, that,

$$\log(1+u) = u - \frac{u^2}{2} + \frac{u^3}{3} - \frac{u^4}{4} + \&c.;$$

$$\therefore \log\left(1+\frac{1}{u}\right) = u^{-1} - \frac{u^{-2}}{2} + \frac{u^{-3}}{3} - \frac{u^{-4}}{4} + \&c.;$$

$$\therefore \log\left\{\left(\frac{1+u}{1+\frac{1}{u}}\right)\right\} = \log u = (u - u^{-1}) - \tfrac{1}{2}(u^2 - u^{-2}) + \&c.$$

For u write $e^{x\sqrt{-1}}$; $\therefore \log u = x\sqrt{-1}$;

$$\therefore x\sqrt{-1} = (e^{x\sqrt{-1}} - e^{-x\sqrt{-1}}) - \tfrac{1}{2}(e^{2x\sqrt{-1}} - e^{-2x\sqrt{-1}}) + \&c.$$

$$= 2\sqrt{-1}\left\{\sin x - \tfrac{1}{2}\sin 2x + \tfrac{1}{3}\sin 3x - \&c.\right\};$$

$$\therefore \frac{x}{2} = \sin x - \tfrac{1}{2}\sin 2x + \tfrac{1}{3}\sin 3x - \&c.;$$

therefore, differentiating,

$$\tfrac{1}{2} = \cos x - \cos 2x + \cos 3x - \cos 4x + \&c.$$

59. By division, $\dfrac{e^{x\sqrt{-1}}}{e^{-x\sqrt{-1}}} = e^{2x\sqrt{-1}} = \dfrac{\cos x + \sqrt{-1}\sin x}{\cos x - \sqrt{-1}\sin x}$

$$= \frac{1+\sqrt{-1}\tan x}{1-\sqrt{-1}\tan x}.$$

$$2x\sqrt{-1} = \log(1+\sqrt{-1}\tan x) - \log(1-\sqrt{-1}\tan x).$$

But $\log(1+u) - \log(1-u) = 2\left\{u + \dfrac{u^3}{3} + \dfrac{u^5}{5} + \&c.\right\}$;

$$\therefore 2x\sqrt{-1} = 2\left\{\sqrt{-1}\tan x + \tfrac{1}{3}(\sqrt{-1}\tan x)^3 + (\sqrt{-1}\tan x)^5 + \&c.\right\}$$

$$= 2\sqrt{-1}\left\{\tan x - \tfrac{1}{3}\tan^3 x + \tfrac{1}{5}\tan^5 x - \&c.\right\};$$

$$\therefore x = \tan x - \tfrac{1}{3}\tan^3 x + \tfrac{1}{5}\tan^5 x - \&c.,$$

a result which has been obtained before.

60. Let $u^2 - ux - a^2 = 0$; find u in terms of x: this is an implicit function, and if

$x = 0$; u becomes U_0; $\therefore (U_0)^2 - a^2 = 0$; $U_0 = \pm a$.

Differentiating, $2u\dfrac{du}{dx} - x\dfrac{du}{dx} - u = 0$;

make $x = 0$; $\therefore u = \pm a$; $\therefore 2aU_1 - a = 0$; $U_1 = \frac{1}{2}$.

Differentiating a second time,

$$(2u - x)\frac{d^2u}{dx^2} + 2\frac{du^2}{dx^2} - 2\frac{du}{dx} = 0;$$

$$\therefore \pm 2aU_2 + 2.\frac{1}{4} - 1 = 0; \quad \therefore U_2 = \pm\frac{1}{4a};$$

$$\therefore (2u - x)\frac{d^3u}{dx^3} + \left(2\frac{du}{dx} - 1\right)\frac{d^2u}{dx^2} + \left(4\frac{du}{dx} - 2\right)\frac{d^2u}{dx^2} = 0;$$

$$\text{or } (2u - x)\frac{d^3u}{dx^3} + 3\left(2\frac{du}{dx} - 1\right)\frac{d^2u}{dx^2} = 0;$$

$$\text{if } x = 0; \quad \because 2U_1 - 1 = 0; \quad \therefore U_3 = 0;$$

$$\therefore u = \pm a + \tfrac{1}{2}x \pm \frac{1}{4a}\frac{x^2}{1.2} + \&c.$$

61. From the expressions for $\sin x$ and $\cos x$ some series may be deduced; which, although not strictly examples of Maclaurin's Theorem, may find a place here.

$$\text{Since } \sin x = x - \frac{x^3}{2.3} + \frac{x^5}{2.3.4.5} - \&c.,$$

and that $\sin x$ vanishes whenever $x =$, 0, $\pm\pi$, $\pm 2\pi$, $\pm 3\pi$, &c.; $\therefore x$, $(\pi^2 - x^2)$, $(2^2\pi^2 - x^2)$, $(3^2\pi^2 - x^2)$, &c. are factors of the equation $\sin x = 0$; and therefore

$$\sin = Ax.(\pi^2 - x^2)(2^2\pi^2 - x^2)(3^2\pi^2 - x^2)(4^2\pi^2 - x^2), \&c.$$

$$= k.x.\left(1 - \frac{x^2}{\pi^2}\right)\left(1 - \frac{x^2}{2^2\pi^2}\right)\left(1 - \frac{x^2}{3^2\pi^2}\right), \&c.$$

$$\text{where } k = A.\pi^2 \times 2^2\pi^2 \times 3^2\pi^2, \&c.;$$

$$\therefore \frac{\sin x}{x} = k\left\{\left(1 - \frac{x^2}{\pi^2}\right)\left(1 - \frac{x^2}{2^2\pi^2}\right)\left(1 - \frac{x^2}{3^2\pi^2}\right)\right\}, \&c.$$

Let $x = 0$; $\therefore \frac{\sin x}{x} = 1$, and the right-hand side of the equation is reduced to k; $\therefore k = 1$;

$$\therefore \sin x = x\left(1 - \frac{x^2}{\pi^2}\right)\left(1 - \frac{x^2}{2^2\pi^2}\right)\left(1 - \frac{x^2}{3^2\pi^2}\right), \text{ &c.}$$

$$= x\left\{1 - \frac{x^2}{\pi^2}\left(\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \text{&c.}\right) + Bx^4 - \text{&c.}\right\}.$$

But $\sin x = x\left\{1 - \frac{x^2}{2.3} + \frac{x^4}{2.3.4.5} - \text{&c.}\right\}$;

therefore, equating coefficients of like powers,

$$\frac{1}{\pi^2}\cdot\left\{\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \text{&c.}\right\} = \frac{1}{6}; \quad \therefore \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \text{&c.} = \frac{\pi^2}{6}.$$

62. Also, since $\cos x = 1 - \frac{x^2}{2} + \frac{x^4}{2.3.4} - \text{&c.}$ vanishes, when

$$x = \pm\frac{\pi}{2}, \quad \pm\frac{3\pi}{2}, \quad \pm\frac{5\pi}{2}, \text{ &c.};$$

$$\therefore \cos x = A\left(\frac{\pi^2}{2^2} - x^2\right)\cdot\left(\frac{3^2\pi^2}{2^2} - x^2\right)\cdot\left(\frac{5^2\pi^2}{2^2} - x^2\right), \text{ &c}$$

$$= k\left(1 - \frac{2^2x^2}{\pi^2}\right)\left(1 - \frac{2^2x^2}{3^2\pi^2}\right)\left(1 - \frac{2^2x^2}{3^2\pi^2}\right), \text{ &c.}$$

whence making $x = 0$, $k = 1$;

$$\therefore \cos x = \left(1 - \frac{2^2x^2}{\pi^2}\right)\left(1 - \frac{2^2x^2}{3^2\pi^2}\right)\left(1 - \frac{2^2x^2}{5^2\pi^2}\right), \text{ &c.}$$

$$= 1 - \frac{2^2x^2}{\pi^2}\left\{\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \text{&c.}\right\} + Bx^4 - \text{&c.}$$

But $\cos x = 1 - \frac{x^2}{1.2} + \frac{x^4}{2.3.4} - \text{&c.}$;

$$\therefore \frac{2^2}{\pi^2}\left\{\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \text{&c.}\right\} = \frac{1}{2}; \quad \therefore \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \text{&c.} = \frac{\pi^2}{8}.$$

63. Again, since

$$\sin x = x\cdot\left(1 - \frac{x^2}{\pi^2}\right)\left(1 - \frac{x^2}{2^2\pi^2}\right)\left(1 - \frac{x^2}{3^2\pi^2}\right), \text{ &c.};$$

$$\therefore \log\sin x = \log x + \log\left(1 - \frac{x^2}{\pi^2}\right) + \log\left(1 - \frac{x^2}{2^2\pi^2}\right) + \text{&c.};$$

$$\therefore \frac{\cos x}{\sin x} = \frac{1}{x} - \frac{\frac{2x}{\pi^2}}{1-\frac{x^2}{\pi^2}} - \frac{\frac{2x}{2^2\pi^2}}{1-\frac{x^2}{2^2\pi^2}} - \frac{\frac{2x}{3^2\pi^2}}{1-\frac{x^2}{3^2\pi^2}}$$

$$= \frac{1}{x} - \frac{\frac{2x}{\pi^2}}{1-\frac{x^2}{\pi^2}} - \frac{\frac{2x}{\pi^2}}{2^2-\frac{x^2}{\pi^2}} - \frac{\frac{2x}{\pi^2}}{3^2-\frac{x^2}{\pi^2}} - \&c.$$

(1⁰) Let $\frac{x^2}{\pi^2} = \theta^2$; $\therefore x = \pi\theta$, and $\frac{2x}{\pi^2} = \frac{2\theta}{\pi}$;

$$\therefore \frac{1}{\tan\pi\theta} = \frac{1}{\pi\theta} - \frac{2\theta}{\pi}\left\{\frac{1}{1^2-\theta^2} + \frac{1}{2^2-\theta^2} + \frac{1}{3^2-\theta^2} + \&c.\right\};$$

$$\therefore \frac{1}{1^2-\theta^2} + \frac{1}{2^2-\theta^2} + \frac{1}{3^2-\theta^2} + \&c. = \frac{1}{2\theta^2} - \frac{\pi}{2\theta\tan\pi\theta}.$$

(2⁰) Let $\frac{x^2}{\pi^2} = -\theta^2$; $\therefore x = \pi\theta\sqrt{-1}$, and $\frac{2x}{\pi^2} = \frac{2\theta\sqrt{-1}}{\pi}$,

and $\frac{\cos x}{\sin x} = \frac{1+e^{-2x\sqrt{-1}}}{1-e^{-2x\sqrt{-1}}}\sqrt{-1} = \frac{1+e^{+2\pi\theta}}{1-e^{+2\pi\theta}}\sqrt{-1}$;

$$\therefore \frac{2\theta\sqrt{-1}}{\pi}\left\{\frac{1}{1^2+\theta^2} + \frac{1}{2^2+\theta^2} + \frac{1}{3^2+\theta^2} + \&c.\right\}$$

$$= \frac{1}{\pi\theta\sqrt{-1}} - \frac{1+e^{2\pi\theta}}{1-e^{2\pi\theta}}\sqrt{-1};$$

$$\therefore \frac{1}{1^2+\theta^2} + \frac{1}{2^2+\theta^2} + \&c. = \frac{\pi}{2}\cdot\frac{e^{2\pi\theta}+1}{e^{2\pi\theta}-1} - \frac{1}{2\theta^2}.$$

64. Again, $\cos x = \left(1-\frac{2^2x^2}{\pi^2}\right)\left(1-\frac{2^2x^2}{3^2\pi^2}\right)\left(1-\frac{2^2x^2}{5^2\pi^2}\right)$ &c.;

$$\therefore \log\cos x = \log\left(1-\frac{2^2x^2}{\pi^2}\right) + \log\left(1-\frac{2^2x^2}{3^2\pi^2}\right) + \&c.;$$

$$\therefore \frac{\sin x}{\cos x} = \frac{2x\cdot\frac{2^2}{\pi^2}}{1-\frac{2^2x^2}{\pi^2}} + \frac{2x\cdot\frac{2^2}{3^2\pi^2}}{1-\frac{2^2x^2}{3^2\pi^2}} + \&c.$$

$$= 2x\left\{\frac{\frac{2^2}{\pi^2}}{1-\frac{2^2x^2}{\pi^2}} + \frac{\frac{2^2}{\pi^2}}{3^2-\frac{2^2x^2}{\pi^2}} + \&c.\right\}.$$

$$(1^0)\quad \text{Let } \frac{2^2x^2}{\pi^2} = \theta^2; \quad \therefore\ x = \frac{\pi\theta}{2}; \quad 2x \cdot \frac{2^2}{\pi^2} = \frac{4\theta}{\pi};$$

$$\therefore \tan\frac{\pi\theta}{2} = \frac{4\theta}{\pi} \cdot \left\{\frac{1}{1^2-\theta^2} + \frac{1}{3^2-\theta^2} + \frac{1}{5^2-\theta^2} + \&c.\right\};$$

$$\therefore \frac{1}{1^2-\theta^2} + \frac{1}{3^2-\theta^2} + \frac{1}{5^2-\theta^2} + \&c. = \frac{\pi}{4\theta}\tan\frac{\pi\theta}{2}.$$

$$(2^0)\quad \text{Let } \frac{2^2x^2}{\pi^2} = -\theta^2; \quad \therefore\ 2x \cdot \frac{2^2}{\pi^2} = \frac{4\theta\sqrt{-1}}{\pi};$$

$$\frac{\sin x}{\cos x} = \frac{1}{\sqrt{-1}}\frac{1-e^{-2x\sqrt{-1}}}{1+e^{-2x\sqrt{-1}}} = \frac{1}{\sqrt{-1}}\frac{1-e^{\pi\theta}}{1+e^{\pi\theta}} = \sqrt{-1}\cdot\frac{e^{\pi\theta}-1}{e^{\pi\theta}+1};$$

$$\therefore \frac{1}{1^2+\theta^2} + \frac{1}{3^2+\theta^2} + \frac{1}{5^2+\theta^2} + \&c. = \frac{\pi}{4\theta}\cdot\frac{e^{\pi\theta}-1}{e^{\pi\theta}+1};$$

Other similar series may be readily deduced.

65. From the expression

$$e^x = 1 + x + \frac{x^2}{1.2} + \frac{x^3}{2.3} + \&c.$$

Lagrange in the Calcul. des Fonctions has derived an expression for the general term of the polynomial $(a+b+c+d+\&c.)^m$.

Thus for x put $(a+b+c+d+\&c.)\,x$;

$$\therefore e^{(a+b+c+d+\&c.)x} = 1 + (a+b+c+\&c.)\,x + \frac{(a+b+c+d+\&c.)^2x^2}{1.2} + \&c.$$

$$+ \frac{(a+b+c+d+\&c.)^m \cdot x^m}{1\ 2\ 3 \ldots\ldots m} + \&c.$$

But $e^{(a+b+c+\&c.)x} = e^{ax} \times e^{bx} \times e^{cx} + \&c.$

$$= \left(1 + ax + \frac{a^2x^2}{1.2} + \frac{a^3x^3}{2.3} + \&c.\right)$$

$$\times \left(1 + bx + \frac{b^2x^2}{1.2} + \frac{b^3x^3}{2.3} + \&c.\right)$$

$$\times \left(1 + cx + \frac{c^2x^2}{1.2} + \frac{c^3x^3}{2.3} + \&c.\right)$$

&c.

Now the m^{th} term of this expansion will be the product of

$$\frac{a^p \cdot x^p}{1 \cdot 2 \ldots p} \times \frac{b^q \cdot x^q}{1 \cdot 2 \ldots q} \times \frac{c^r \cdot x^r}{1 \cdot 2 \ldots r} \times \&c.,$$

$$\text{where } p + q + r + \&c. \ldots\ldots = m,$$

whence $(a + b + c + d + \&c.)^m$ will consist of terms included under the general expression

$$\frac{1 \cdot 2 \cdot 3 \ldots m \times a^p \cdot b^q \cdot c^r \ldots}{1 \cdot 2 \ldots p \times 1 \cdot 2 \ldots q \times 1 \cdot 2 \cdot 3 \ldots r \times \&c.},$$

subject to the condition that $p + q + r + \&c. = m$.

EXAMPLES.

(1) If $u = \dfrac{1}{1 + x^2}$; $\dfrac{d^4u}{dx^4} = 24 \cdot \left\{\dfrac{1 - 10x^2 + 5x^4}{(1 + x^2)^5}\right\}$.

(2) If $u = \dfrac{x}{\sqrt{1 - x^2}}$; $\dfrac{d^3u}{dx^3} = \dfrac{3 \cdot (1 + 4x^2)}{(1 - x^2)^{\frac{7}{2}}}$.

(3) If $u = \sin x$; $\dfrac{d^nu}{dx^n} = \sin\left(x + n\dfrac{\pi}{2}\right)$.

For $\dfrac{du}{dx} = \cos x = \sin\left(\dfrac{\pi}{2} - x\right) = \sin\left(x + \dfrac{\pi}{2}\right)$;

$\dfrac{d^2u}{dx^2} = \cos\left(x + \dfrac{\pi}{2}\right) = \sin\left(\dfrac{\pi}{2} + x + \dfrac{\pi}{2}\right) = \sin\left(x + 2\dfrac{\pi}{2}\right)$.

(4) If $u = \cos x$; $\dfrac{d^nu}{dx^n} = \cos\left(x + n\dfrac{\pi}{2}\right)$.

(5) If $u = x^n \cdot e^x$; $\dfrac{d^4u}{dx^4} = e^x\{x^n + 4nx^{n-1} + 6n(n-1)x^{n-2}$

$+ 4n \cdot (n-1) \cdot (n-2)x^{n-3} + n(n-1)(n-2)(n-3) \cdot x^{n-4}\}$.

(6) If $u = e^x \sin x$, $\dfrac{d^2u}{dx^2} = 2e^x \cos x$,

$$\frac{d^4u}{dx^4} = -4u, \quad \frac{d^8u}{dx^8} = 4^2u, \quad \frac{d^{12}u}{dx^{12}} = -4^3u.$$

(7) Shew that

$$\sin(a + bx) = \sin a + bx \cos a - \frac{b^2x^2}{2} \cdot \sin a - \frac{b^3x^3}{2 \cdot 3} \cos a + \&c.$$

$$\cos(a + bx) = \cos a - bx \sin a - \frac{b^2x^2}{2} \cdot \cos a + \frac{b^3x^3}{2.3} \sin a + \&c.$$

$$\log(a + bx) = \log a + \frac{bx}{a} - \frac{b^2x^2}{2a^2} + \frac{b^3x^3}{3a^3} - \&c.$$

(8) And $\sec x = 1 + \frac{x^2}{2} + \frac{5x^4}{2.3.4} + \&c.$

$$(\cos x)^3 = 1 - \frac{3x^2}{2} + \frac{7x^4}{8} - \&c.$$

$$(\tan x)^4 = x^4 + \frac{4}{3}x^6 + \frac{6}{5}x^8 + \&c.$$

$$e^{\sin x} = 1 + x + \frac{x^2}{2} - \frac{3x^4}{2.3.4} - \frac{8x^5}{2.3.4.5} - \&c.$$

(9) Shew that $\sqrt{e^x + 1}$

$$= \sqrt{2}\left(1 + \frac{1}{2^2} \cdot \frac{x}{1} + \frac{3}{2^4} \cdot \frac{x^2}{1.2} + \frac{7}{2^6} \frac{x^3}{2.3} + \&c.\right)$$

(10) Expand $\sin(a + bx + cx^2)$, and $\log(a + bx + cx^2)$ according to the powers of x.

(11) If $\cos(m) - \cos(m + y) = x$, shew that

$$y = \frac{x}{\sin m} - \tfrac{1}{2} \cot m \cdot \left(\frac{x}{\sin m}\right)^2 + \&c.$$

(12) If $\sin u = m \sin x$, prove that

$$u = mx + \frac{m(m^2 - 1)x^3}{2.3} + \frac{m(9m^4 - 10m^2 + 1)x^5}{2.3.4.5} + \&c.$$

(13) If $u^3 - ux = 1$; $u = 1 + \frac{x}{3} - \frac{x^3}{81} + \frac{x^4}{243} - \&c.$

(14) $\mathrm{Log}\left(\frac{x}{x-1}\right) = \frac{1}{x-1} - \frac{1}{2}\frac{1}{(x-1)^2} + \frac{1}{3}\frac{1}{(x-1)^3} - \&c.$

(15) Shew that $\frac{x}{e^x - 1} = 1 - \frac{1}{2}x + \frac{1}{6}\frac{x^2}{1.2.} - \frac{1}{30}\frac{x^4}{2.3.4} - \&c.$

(16) If a_n and b_n respectively represent the coefficients of x^n in the expansions of $u = f(x)$, and $\log u$: then shall

$$n a_n = b_1 a_{n-1} + 2b_2 a_{n-2} + 3b_3 a_{n-3} + \&c. + n b_n a_0.$$

CHAPTER IV.

TAYLOR'S THEOREM.

66. If $u = f(x)$, and $u_1 = f(x + h)$, then by Taylor's Theorem, so called from its inventor,

$$u_1 = u + \frac{du}{dx}h + \frac{d^2u}{dx^2}\frac{h^2}{1.2} + \frac{d^3u}{dx^3}\frac{h^3}{2.3} + \&c. + \frac{d^nu}{dx^n}\cdot\frac{h^n}{2.3\ldots n} + \&c.$$

The proof of this theorem may be made to depend upon the proposition, that if $u_1 = f(x + h)$; $\dfrac{du_1}{dx} = \dfrac{du_1}{dh}$,

or the coefficient of h is the same in the expansion $f(x + 2h)$, whether in $f(x + h)$, x become $x + h$, or h become $h + h$.

Let $x + h = x_1$; $\therefore u_1 = f(x_1)$ and $\dfrac{du_1}{dx_1} = \dfrac{d.f(x_1)}{dx_1}$.

But $\dfrac{du_1}{dx_1} = \dfrac{du_1}{dx}\cdot\dfrac{dx}{dx_1}$ = also $\dfrac{du_1}{dh}\cdot\dfrac{dh}{dx_1}$.

But h constant, $\dfrac{dx}{dx_1} = 1$; x constant, $\dfrac{dh}{dx_1} = 1$;

$$\therefore \frac{du_1}{dx} = \frac{du_1}{dh}.$$

Hence also $\dfrac{d\left(\dfrac{du_1}{dx}\right)}{dx} = \dfrac{d.\left(\dfrac{du_1}{dh}\right)}{dh}$; or $\dfrac{d^2u_1}{dx^2} = \dfrac{d^2u_1}{dh^2}$;

$$\text{and finally } \frac{d^nu_1}{dx^n} = \frac{d^nu_1}{dh^n}.$$

$$\text{Let } \therefore u_1 = f(x + h) = u + \frac{du}{dx}.h + Ph^\alpha + Qh^\beta + Rh^\gamma + \&c.$$

P, Q, R, &c. being functions of x, and the indices of h in order of magnitude, beginning with the least;

$$\therefore \frac{du_1}{dx} = \frac{du}{dx} + \frac{d^2u}{dx^2}h + \frac{dP}{dx}h^\alpha + \frac{dQ}{dx}h^\beta + \frac{dR}{dx}h^\gamma + \&c.,$$

$$\text{and } \frac{du_1}{dh} = \frac{du}{dx} + \alpha Ph^{\alpha-1} + \beta Qh^{\beta-1} + \gamma Rh^{\gamma-1} + \&c.;$$

$\therefore$ taking away the common term $\frac{du}{dx}$, and dividing by h,

$$\frac{d^2u}{dx^2} + \frac{dP}{dx}h^{\alpha-1} + \frac{dQ}{dx}h^{\beta-1} + \&c.....$$

$$= \alpha P h^{\alpha-2} + \beta Q h^{\beta-2} + \gamma R h^{\gamma-2} + \&c.$$

Now since in the upper series there is a term $\frac{d^2u}{dx^2}$ independent of h, there must also be an equal and corresponding term in the lower series; which will be the first, since α is the least index;

$$\therefore \frac{d^2u}{dx^2} = \alpha P h^{\alpha-2}; \quad \therefore \alpha - 2 = 0; \text{ and } \alpha = 2;$$

$$\therefore 2P = \frac{d^2u}{dx^2}; \text{ and } P = \frac{d^2u}{dx^2}\cdot\frac{1}{1.2}.$$

Similarly, $\frac{dP}{dx}h^{\alpha-1}$, or $\frac{dP}{dx}h$ must $= \beta Q h^{\beta-2}$;

$$\therefore \beta - 2 = 1; \quad \beta = 3; \quad \text{and } Q = \tfrac{1}{3}\frac{dP}{dx} = \frac{d^3u}{dx^3}\cdot\frac{1}{2.3}.$$

$$\text{Again, } \frac{dQ}{dx}h^{\beta-1} = \frac{dQ}{dx}h^2 = \gamma R h^{\gamma-2};$$

$$\therefore \gamma - 2 = 2; \quad \gamma = 4; \quad \therefore R = \tfrac{1}{4}\frac{dQ}{dx} = \frac{d^4u}{dx^4}\cdot\frac{1}{2.3.4};$$

$$\text{and } \therefore u_1 = u + \frac{du}{dx}h + \frac{d^2u}{dx^2}\frac{h^2}{1.2} + \frac{d^3u}{dx^3}\frac{h^3}{2.3} + \&c.$$

$$+ \frac{d^nu}{dx^n}\frac{h^n}{1.2.3....n} + \&c.,$$

a theorem which will give the expansion of $f(x+h)$ in all cases, if x remain indeterminate.

67. In the preceding proof the indices of h after the first have been considered as unknown; but if we extend the expansion of $f(x+h)$, as in Cor. 1. Art. 8, it will be found that they are the natural numbers, and we may assume that

$$u_1 = u + \frac{du}{dx}h + Ph^2 + Qh^3 + Rh^4 + \&c.$$

$$\therefore \frac{du_1}{dx} = \frac{du}{dx} + \frac{d^2u}{dx^2}h + \frac{dP}{dx}h^2 + \frac{dQ}{dx}h^3 + \&c.$$

$$\frac{du_1}{dh} = \frac{du}{dx} + 2Ph + 3Qh^2 + 4Rh^3 + \&c.$$

whence equating the coefficients of the same powers of h,

$$2P = \frac{d^2u}{dx^2}; \quad \therefore P = \tfrac{1}{2} \cdot \frac{d^2u}{dx^2},$$

$$3Q = \frac{dP}{dx} = \tfrac{1}{2} \cdot \frac{d^3u}{dx^3}; \quad \therefore Q = \frac{1}{2.3} \cdot \frac{d^3u}{dx^3},$$

$$4R = \frac{dQ}{dx} = \frac{1}{2.3} \cdot \frac{d^4u}{dx^4}; \quad \therefore R = \frac{1}{2.3.4} \cdot \frac{d^4u}{dx^4};$$

$$\therefore u_1 = u + \frac{du}{dx}h + \frac{d^2u}{dx^2}\frac{h^2}{1.2} + \frac{d^3u}{dx^3}\frac{h^3}{2.3} + \&c.$$

Cor. We may now deduce the theorem of Maclaurin which we have proved by an independent process in the preceding chapter.

For by making $x = 0$, u_1 becomes $f(h)$ and u, $\frac{du}{dx}$, $\frac{d^2u}{dx^2}$, $\frac{d^3u}{dx^3}$, &c. become U_0, U_1, U_2, U_3, &c.;

$$\therefore f(h) = U_0 + U_1h + U_2\frac{h^2}{1.2} + U_3\frac{h^3}{2.3} + \&c.$$

or putting x for h, in which case u may be put for $f(x)$,

$$u = U_0 + U_1x + U_2\frac{x^2}{1.2} + U_3\frac{x^3}{2.3} + \&c.$$

the theorem required.

EXAMPLES.

68. To expand $\sin(x+h)$, $\cos(x+h)$, $\log(x+h)$ and $(x+h)^n$, by Taylor's Theorem,

$$u_1 = u + \frac{du}{dx}h + \frac{d^2u}{dx^2}\frac{h^2}{1.2} + \frac{d^3u}{dx^3}\frac{h^3}{2.3} + \&c.$$

(1) $u = \sin x$;

$$\therefore \frac{du}{dx} = \cos x, \quad \frac{d^2u}{dx^2} = -\sin x, \quad \frac{d^3u}{dx^3} = -\cos x, \quad \frac{d^4u}{dx^4} = \sin x,$$

after which the values recur;

$$\therefore\ u_1 = \sin(x+h) = \sin x + \cos x\,.h - \sin x\,\frac{h^2}{1.2} - \cos x\,\frac{h^3}{2.3}$$

$$+ \sin x\,\frac{h^4}{2.3.4} + \cos x\,\frac{h^5}{2.3.4.5} - \&c.$$

(2) $u = \cos x$;

$$\frac{du}{dx} = -\sin x,\ \frac{d^2u}{dx^2} = -\cos x,\ \frac{d^3u}{dx^3} = \sin x,\ \frac{d^4u}{dx^4} = \cos x,$$

after which the values recur;

$$\therefore\ u_1 = \cos(x+h) = \cos x - \sin x\,.\,\frac{h}{1} - \cos x\,\frac{h^2}{1.2}$$

$$+ \sin x\,.\,\frac{h^3}{2.3} + \cos x\,.\,\frac{h^4}{2.3.4} - \&c.$$

Cor. If in the two expansions we make $x = 0$, we have

$$\sin h = h - \frac{h^3}{2.3} + \frac{h^5}{2.3.4.5} - \&c.$$

$$\cos h = 1 - \frac{h^2}{1.2} + \frac{h^4}{2.3.4} - \&c.$$

(3) $u = \log(x)$;

$$\therefore\ \frac{du}{dx} = \frac{1}{x} = x^{-1},\ \frac{d^2u}{dx^2} = -x^{-2},\ \frac{d^3u}{dx^3} = 2x^{-3},\ \frac{d^4u}{dx^4} = -2.3x^{-4};$$

$$\therefore\ u_1 = \log(x+h) = \log x + \frac{h}{x} - \tfrac{1}{2}\cdot\frac{h^2}{x^2} + \tfrac{1}{3}\cdot\frac{h^3}{x^3} - \tfrac{1}{4}\cdot\frac{h^4}{x^4} + \&c.$$

Let $x = 1$; $\therefore \log x = 0$;

$$\therefore\ \log(1+h) = h - \tfrac{1}{2}h^2 + \tfrac{1}{3}h^3 - \tfrac{1}{4}h^4 + \tfrac{1}{5}h^5 - \&c.$$

(4) $u = x^n$;

$$\therefore\ \frac{du}{dx} = nx^{n-1},\ \frac{d^2u}{dx^2} = n(n-1)x^{n-2},\ \frac{d^3u}{dx^3} = n\,.\,(n-1)\,(n-2)\,.x^{n-3};$$

$$\therefore\ u_1 = (x+h)^n = x^n + n\,.\,x^{n-1}h + \frac{n\,.\,(n-1)}{1.2}\,.\,x^{n-2}h^2$$

$$+ \frac{n\,(n-1)\,(n-2)}{1.2.3}\,.\,x^{n-3}h^3 + \&c.$$

69. The following Proposition which is used in some

demonstrations of the parallelogram of forces, is a good application of the Theorem. Given that

$$f(x).f(h) = f(x+h) + f(x-h),$$

find the form of $f(x)$.

Let u be put for $f(x)$;

$$\therefore\ u.f(h) = 2\left\{u + \frac{d^2u}{dx^2}\,\frac{h^2}{1.2} + \frac{d^4u}{dx^4}\,\frac{h^4}{2.3.4} + \&c.\right\};$$

$$\therefore\ f(h) = 2\left\{1 + \frac{1}{u}\,\frac{d^2u}{dx^2}\,\frac{h^2}{1.2} + \frac{1}{u}\,\frac{d^4u}{dx^4}\,\frac{h^4}{2.3.4} + \&c.\right\}.$$

Now since h is independent of x, the coefficients $\frac{1}{u}.\frac{d^2u}{dx^2}$, $\frac{1}{u}\,\frac{d^4u}{dx^4}$, &c., which cannot contain h, must be constant.

Let $\frac{1}{u}\,\frac{d^2u}{dx^2} = -a^2$; $\therefore\ \frac{d^2u}{dx^2} = -a^2u$; $\frac{d^4u}{dx^4} = -a^2\frac{d^2u}{dx^2} = a^4u$.

Hence $f(h) = 2\left\{1 - \frac{a^2h^2}{1.2} + \frac{a^4h^4}{2.3.4} - \&c.\right\} = 2\cos ah$,

and $\therefore\ f(x) = 2\cos ax$; and $f(x \pm h) = 2\cos(ax \pm ah)$,

which may be verified by the formula

$$2\cos A\,.\,2\cos B = 2\cos(A+B) + 2\cos(A-B).$$

70. Taylor's Theorem may be used to approximate to the roots of equations.

Let $X = 0$ be an equation, of which x is one of the roots, and a an approximate value of x, so that $x = a + h$, h being a very small quantity, hence since $X = 0$ is a function of x;

$$\therefore\ X = 0 = f(a+h) = f(a) + \frac{d.f(a)}{da}h + \frac{d^2.f(a)}{da^2}\,\frac{h^2}{1.2} + \&c.$$

but since h is assumed very small, we may neglect the terms after the second, and so obtain an approximate value of h;

$$\therefore\ 0 = f(a) + \frac{d.f(a)}{da}h;\ \ \therefore\ h = \frac{-f(a)}{\frac{d.f(a)}{da}} = \frac{-f(a)}{p},$$

$$\text{and}\ \ x = a - \frac{f(a)}{p}.$$

If this value of x be not sufficiently near the true one, let it be put $= a_1$, and the process repeated: we shall at length arrive at results more and more near the true one.

Ex. 1. $x^3 - 3x + 1 = 0$. By trial 1.5 is found to be near one of the roots.

$$f(a) = a^3 - 3a + 1 = (1.5)^3 - 3 \times (1.5) + 1 = -.125,$$

$$\frac{d.f(a)}{da} = 3a^2 - 3 = 6.75 - 3 = 3.75;$$

$$\therefore\ h = \frac{.125}{3.75} = .033;\quad \therefore\ x = 1.5 + .033 = 1.533.$$

Ex. 2. $x^x = 100$. Since $3^3 = 27$ and $4^4 = 256$; x lies between 3 and 4; let $a = 3.5$.

$$\text{Now } x \log x - \log 100 = 0 = u;\quad \therefore\ 1 + \log x = \frac{du}{dx};$$

$$\therefore\ f(a) = 3.5 \log (3.5) - \log 100;\quad \frac{d.f(a)}{da} = 1 + \log (3.5).$$

$$\text{But } \log_e 100 = 4.60517;\quad \log_e 3.5 = 1.25276;$$

$$\therefore f(a) = 3.5 \times 1.25276 - 4.60517 = -.22051;\quad \frac{d.f(a)}{da} = 2.25276;$$

$$\therefore\ h = \frac{.22051}{2.25276} = .09832;\quad \text{and } x = a + h = 3.59832;$$

a more exact value may be obtained by putting 3.59832 for a.

The Napierian logarithms are obtained from the common logarithms by dividing each logarithm by the number .43429.

$$\text{Thus } \log_e 100 = \frac{2}{.43429} = 4.60517.$$

71. Transform the equation $x^n - px^{n-1} + qx^{n-2} - \&c. = 0$, into one whose roots shall be diminished by a constant quantity z.

Let $x = z + y$; $X = f(z + y)$, and let $Z = f(z)$;

$$\therefore\ X = Z + \frac{dZ}{dz}y + \frac{d^2Z}{dz^2} \cdot \frac{y^2}{1.2} + \frac{d^3Z}{dz^3} \cdot \frac{y^3}{2.3} + \&c. = 0,$$

Or if Z_1, Z_2, Z_3, &c.... Z_n, be put for the differential coefficients, the transformed equation becomes

$$Z + Z_1 y + \frac{Z_2 y^2}{1.2} + \frac{Z_3 y^3}{2.3} + \&c. + \frac{Z_{n-1} y^{n-1}}{1.2...(n-1)} + \frac{Z_n y^n}{2.3...n} = 0,$$

where Z is the value of X, when z is put for x;

$$\therefore\ Z = z^n - pz^{n-1} + qz^{n-2} - \&c.$$

$$\text{and } Z_1 = nz^{n-1} - (n-1)pz^{n-2} + (n-2)qz^{n-3} - \&c.$$

$$\vdots$$

$$Z_{n-1} = n(n-1)(n-2)\ldots 3\,.\,2z - (n-1)\,.\,(n-2)\ldots 2\,.\,p,$$

$$\text{and } Z_n = n(n-1)(n-2)\ldots 3\,.\,2;$$

therefore by substitution, the transformed equation will become, after writing the terms in an inverse order,

$$y^n + (nz - p)y^{n-1} + \&c. + Z = 0.$$

Cor. This transformed equation may be used to take away any particular term of an equation, by putting any of the coefficients Z_1, Z_2, &c. $= 0$, and substituting in the others the value of z derived from it.

Ex. Take away the second term from the equation

$$3x^3 + 15x^2 + 25x - 3 = 0.$$

The transformed equation is, when $x = z + y$,

$$Z + Z_1 y + \frac{Z_2 y^2}{1\,.\,2} + \frac{Z_3 y^3}{2\,.\,3} = 0, \quad (\text{for } Z_4 = 0),$$

$$Z = 3z^3 + 15z^2 + 25z - 3,$$

$$Z_1 = 9z^2 + 30z + 25,$$

$$Z_2 = 18z + 30,$$

$$Z_3 = 18, \text{ and } Z_4 = 0.$$

$$\text{But } Z_2 = 0; \quad \therefore\ z = -\frac{30}{18} = \frac{-5}{3};$$

$$Z_1 = 25 - 50 + 25 = 0,$$

$$Z = -\frac{125}{9} + \frac{125}{3} - \frac{225}{3} - 3 = \frac{-152}{9};$$

$$\therefore\ -\frac{152}{9} + \frac{18y^3}{6} = 0; \quad \therefore\ y^3 - \frac{152}{27} = 0.$$

72. Let $u = (a + bx + cx^2)^r$. Find $\frac{d^n u}{dx^n}$.

Since the coefficient of h^n in Taylor's Theorem is $\frac{d^n u}{dx^n}$ divided by $1\,.\,2\ldots n$, if we expand

$$\{a + b(x+h) + c(x+h)^2\}^r,$$

and collect the terms which are multiplied by h^n, these when multiplied by $1\,.\,2\,.\,3\ldots n$ will give $\frac{d^n u}{dx^n}$;

$$\therefore \{a+b(x+h)+c(x+h)^2\}^r = \{a+bx+cx^2+(b+2cx)h+ch^2\}^r.$$

Let $a+bx+cx^2=p$, and $b+2cx=q$;

$$\therefore u_1=(p+qh+ch^2)^r=p^r\left\{1+\frac{q}{p}h+\frac{4pc}{4p^2}h^2\right\}^r.$$

But $4pc=4ac+4bcx+4c^2x^2=(b+2cx)^2+4ac-b^2$

$$=q^2+e^2, \quad \text{if } e^2=4ac-b^2;$$

$$\therefore u_1=p^r\left\{1+\frac{q}{p}h+\frac{q^2}{4p^2}h^2+\frac{e^2h^2}{4p^2}\right\}^r=p^r\left\{\left(1+\frac{q}{2p}h\right)^2+\frac{e^2h^2}{4p^2}\right\}^r$$

$$=p^r\{(1+q_1h)^2+e_1^2h^2\}^r$$

$$=p^r\{(1+q_1h)^{2r}+r.(1+q_1h)^{2r-2}.e_1^2h^2$$

$$+r.\frac{(r-1)}{2}(1+q_1h)^{2r-4}e_1^4h^4+\&c.\}$$

Then writing down the coefficient of h^n in $(1+q_1h)^{2r}$,

$$\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots h^{n-2}\ldots(1+q_1h)^{2r-2},$$

$$\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots h^{n-4}\ldots(1+q_1h)^{2r-4},$$

$$\&c.\ldots\ldots\&c.$$

we shall have by addition the coefficient of h^n, which multiplied by $1.2.3\ldots n$, will give $\frac{d^nu}{dx^n}$.

Now by the Binomial Theorem, the coefficient of

$$h^n \text{ in } (1+q_1h)^{2r} = \frac{2r.(2r-1).(2r-2)\ldots(2r-n+1)}{1\ .\ 2\ .\ 3\ \ldots\ n}q_1^n\ldots(1),$$

$$\text{of } h^{n-2} \text{ in } (1+q_1h)^{2r-2} = \frac{(2r-2.(2r-3)\ldots(2r-n+1)}{1\ .\ 2\ \ldots\ldots\ (n-2)}q_1^{n-2}\ldots(2),$$

$$\text{of } h^{n-4} \text{ in } (1+q_1h)^{2r-4} = \frac{(2r-4).(2r-5)\ldots(2r-n+1)}{1\ .\ 2\ \ldots\ldots\ (n-4)}q_1^{n-4}\ldots(3);$$

therefore, substituting $\frac{q}{2p}$ for q_1, $\frac{e^2}{4p^2}$ for e_1^2, and multiplying (2) by $r\times e_1^2$, and (3) by $\frac{r.(r-1)}{1.2}e_1^4$, &c. and the sum of (1), (2), (3), &c. by $1.2.3\ldots n.p^r$,

$$\frac{d^nu}{dx^n}=1.2.3\ldots np^r\left\{\frac{2r.(2r-1).(2r-2)\ldots(2r-n+1)}{1\ .\ 2\ .\ 3\ \ldots\ldots\ n}\frac{q^n}{2^np^n}\right.$$

$$\left.+\frac{r.(2r-2)\ldots(2r-n+1)}{1\ .\ 2\ \ldots\ldots(n-2)}.\frac{q^n}{2^np^n}.\frac{e^2}{q^2}+\&c.\right\}$$

$$=2r.(2r-1).(2r-2)\ldots(2r-n+1)\left(\frac{q}{2}\right)^n.p^{r-n}\left\{1+r.\frac{n.(n-1)}{2r.(2r-1)}.\frac{e^2}{q^2}\right.$$
$$+\frac{r.(r-1)}{1.2}.\frac{n.(n-1)\ldots(n-3)}{2r.(2r-1)\ldots(2r-3)}.\frac{e^4}{q^4}$$
$$\left.+\frac{r.(r-1)(r-2)}{1\ .\ 2\ .\ 3}.\frac{n.(n-1)\ldots(n-5)}{2r\ \ldots\ldots\ (2r-5)}.\frac{e^6}{q^6}+\&c.\right\}$$

Ex. 1. Let $u=\dfrac{1}{\sqrt{1-x^2}}$, the example given by Euler.

Here $r=-\frac{1}{2}$, $a=1$, $b=0$, $c=-1$, $p=1-x^2$, $q=-2x$, $e^2=-4$;

$$\therefore\ \frac{d^n.(1-x^2)^{-\frac{1}{2}}}{dx^n}=\frac{1.2\ldots n\ .\ x^n}{(1-x^2)^{n+\frac{1}{2}}}.\left\{1+\tfrac{1}{2}.\frac{n.(n-1)}{1.2}.\frac{1}{x^2}\right.$$
$$\left.+\frac{1.3}{2.4}.\frac{n.(n-1)\ldots(n-3)}{1\ .\ 2\ \ \ldots\ \ 4}.\frac{1}{x^4}+\frac{1.3.5}{2.4.6}.\frac{n.(n-1)\ldots(n-5)}{1\ .\ 2\ \ldots\ 6}.\frac{1}{x^6}+\&c.\right\},$$

the law of which is obvious.

Ex. 2. Let $u=\sqrt{1-x^2}$.

Here $r=\frac{1}{2}$; $2r-1=0$; $\therefore$ some of the coefficients $=\dfrac{0}{0}$.

The example may thus be put in a proper form.

Since $u=\sqrt{1-x^2}$; $\quad\therefore\ \dfrac{du}{dx}=\dfrac{-x}{\sqrt{1-x^2}}$;

$$\frac{d^2u}{dx^2}=-\frac{1}{(1-x^2)^{\frac{3}{2}}},\quad\text{or}\quad\frac{d^2.(\sqrt{1-x^2})}{dx^2}=-(1-x^2)^{-\frac{3}{2}};$$
$$\therefore\ \frac{d^n.\sqrt{1-x^2}}{dx^n}=-\frac{d^{n-2}}{dx^{n-2}}.(1-x^2)^{-\frac{3}{2}};$$

and writing $n-2$ for n, and $-\dfrac{3}{2}$ for r, we shall have the required term.

EXAMPLES.

(1) $\mathrm{Tan}(x+h)=\tan x+h.\sec^2x+h^2.\tan x.\sec^2x+\&c.$

(2) $\mathrm{Sin}^{-1}(x+h)=\sin^{-1}x+\dfrac{h}{\sqrt{1-x^2}}+\dfrac{h^2x}{2(1-x^2)^{\frac{3}{2}}}$
$$+\frac{(1+2x^2)h^3}{2.3(1-x^2)^{\frac{5}{2}}}+\&c.$$

(3) Prove that if $u = f(x)$,

$$f\left(\frac{x}{2}\right) = u - \frac{du}{dx}\frac{x}{2} + \frac{d^2u}{dx^2}\cdot\frac{x^2}{2.2^2} - \frac{d^3u}{dx^3}\cdot\frac{x^3}{2.3.2^3} + \&c.$$

$$f\left(\frac{x}{1+x}\right) = u - \frac{du}{dx}\cdot\frac{x^2}{1+x} + \frac{d^2u}{dx^2}\cdot\frac{x^4}{2(1+x)^2} - \frac{d^3u}{dx^3}\cdot\frac{x^6}{2.3(1+x)^3} + \&c.$$

(4) Approximate to a root of the equations

(1) $x^3 - 12x - 28 = 0$. Ans. $x = 4.302$.

(2) $x^4 + x - 3 = 0$. Ans. $x = 1.165$.

(5) If $u = f(x)$, and if when $x = a$; $u = b$; then

$$u = b - \frac{du}{dx}(x-a) + \frac{d^2u}{dx^2}\cdot\frac{(x-a)^2}{2} - \frac{d^3u}{dx^3}\cdot\frac{(x-a)^3}{2.3} + \&c.$$

In the expansion of $f(x+h)$, put for h, $x-a$, and in $f(x+h)$, for $x+h$, put a, and the above theorem due to Euler is readily found.

(6) Find the n^{th} differential coefficient of $\sqrt{1+x+x^2}$.

By following the method given in Art. 72, we have, if

$$u = \sqrt{1+x+x^2}; \quad q = 2x+1; \quad u_1 = \sqrt{(u^2+qh)+h^2};$$

whence by expanding, the coefficient of h^n may be found. Or we may apply the formula after differentiating twice, since

$$\frac{d^2u}{dx^2} = \frac{3}{4}(1+x+x^2)^{-\frac{3}{2}}.$$

CHAPTER V.

FAILURE OF TAYLOR'S THEOREM; LIMITS OF THE SAME THEOREM.

---

73. By the Theorem of Taylor we are enabled to expand the $f(x+h)$ into a series of the form

$$f(x)+ph+qh^2+rh^3+\&c.$$

where the powers of h are integral and ascend.

Indeed we have proved, (Art. 8), that so long as x retains its general value, the expansion of $f(x+h)$ cannot contain fractional powers of h. As this proposition is an important one, we here give the reasoning commonly made use of to establish its truth.

Assume $f(x+h)=u+Ph^{\frac{m}{n}}+R,$

where R represents the sum of all the other terms.

Then since $x+h$ enters $f(x+h)$ in the same manner as x enters $f(x)$, it is plain that both functions (undeveloped) have the same number of values, and that the developement of $f(x+h)$ ought to contain no more than $f(x)$ or $f(x+h)$ does.

But if particular values be given to x, which will make P neither infinite nor evanescent; then to each value of x there will correspond n values of $Ph^{\frac{m}{n}}$, since $h^{\frac{m}{n}}$ has n different values; and consequently the expanded function will contain n times as many values as the unexpanded one; and therefore n times as many as $f(x)$, which is manifestly contradictory.

74. If then such a value as a given to x in $f(x+h)$ makes the unexpanded function $f(a+h)$ contain fractional powers of h, we cannot expect that Taylor's Theorem will give the required developement. Now the hypothesis that $x=a$ introduces a fractional index of h into $f(x+h)$, supposes that in the original function there must have been some such term as $(x-a)^{\frac{m}{n}}$, which becomes $(x-a+h)^{\frac{m}{n}}$ in u_1, or $h^{\frac{m}{n}}$ when $x=a$. In such a case it is clear that some of the differential coefficients will become infinite, when $x=a$.

As an illustration, let us suppose that

$$u = b + (x-a)^{\frac{m}{n}};$$

$$\therefore \frac{du}{dx} = \frac{m}{n}(x-a)^{\frac{m}{n}-1};$$

$$\frac{d^2u}{dx^2} = \frac{m}{n}\left(\frac{m}{n}-1\right)(x-a)^{\frac{m}{n}-2};$$

$$\therefore u_1 = b + (x-a)^{\frac{m}{n}} + \frac{m}{n}.(x-a)^{\frac{m}{n}-1}h$$

$$+\frac{m}{n}.\left(\frac{m}{n}-1\right).(x-a)^{\frac{m}{n}-2}\frac{h^2}{1.2} + \&c.$$

$$+\frac{m}{n}.\left(\frac{m}{n}-1\right)\ldots\left(\frac{m}{n}-p+1\right)(x-a)^{\frac{m}{n}-p}\frac{h^p}{1.2\ldots p} + \&c.$$

where if $\frac{m}{n} < p$ and $> p-1$, the $(p+1)^{\text{th}}$ term and all that follow it will become infinite when $x = a$.

This circumstance of the differential coefficients becoming infinite when $x = a$ is called the *Failure* of Taylor's Theorem, an improper phrase, since it should be taken as an index that the function cannot be expanded according to the integral powers of h.

75. Again, as the general expansion of $f(x+h)$ can never contain negative powers of h, for if $f(x+h)$ could

$$= A + Bh^{-\alpha} + \&c.$$

if $h = 0$, $f(x+h)$ instead of becoming $f(x)$, would be infinite, we may be led to expect that if $x = a$ introduces into the unexpanded function $f(x+h)$ a term involving $h^{-\alpha}$, the expansion by Taylor's Theorem will indicate some absurdity. Now it is clear that to have such a term dependent on $h^{-\alpha}$, we must originally have had such a term as $\frac{M}{(x-a)^\alpha}$; for putting $x+h$ for x,

$$\frac{M}{|x-a|^\alpha} \text{ becomes } \frac{M}{(x+h-a)^\alpha} = \frac{M}{h^\alpha},$$

when $x = a$. M not being supposed to vanish when $x = a$.

Here all the differential coefficients of $\frac{1}{(x-a)^\alpha}$ are infinite when $x = a$.

76. The theorem therefore fails whenever $x = a$ makes some surd disappear from $u = f(x)$, and therefore introduces into $u_1 = f(x + h)$, a term involving a fractional power of h; or when $x = a$ renders the original function infinite.

As an example of the first case, let $u = b + \sqrt{x - a}$;

$$\therefore\ u_1 = b + \sqrt{x + h - a}$$

$$= b + \sqrt{x - a} + \tfrac{1}{2} \cdot \frac{1}{\sqrt{x - a}} h - \tfrac{1}{4} \cdot \frac{1}{(x - a)^{\frac{3}{2}}} \frac{h^2}{1 \cdot 2} + \&c.$$

Make $x = a$; $\therefore\ u = b$, $u_1 = b + \sqrt{h}$,

and the expanded function contains infinite terms.

As an example of the second case, let $u = \dfrac{1}{x - a}$;

$$\therefore\ u_1 = \frac{1}{x - a + h} = \frac{1}{x - a} - \frac{h}{(x - a)^2} + \frac{h^2}{(x - a)^3} - \&c.$$

where $u = \infty$, $u_1 = \dfrac{1}{h}$, and the terms of the expanded function are infinite, when $x = a$.

77. Should however $f(a + h)$ contain, when expanded, integral powers of h as far as the $(n - 1)^{\text{th}}$, and afterwards fractional powers, the first (n) coefficients may be found by means of Taylor's Theorem.

Let $f(a + h) = A + Bh + Ch^2 + \&c. + Nh^{n-1} + Ph^{\alpha} + \&c.$ where α is a fraction between $n - 1$ and n.

Now since the coefficients A, B, C, N, do not contain h we may obtain their values by Maclaurin's Theorem, by finding the differential coefficients of $f(a + h)$ with respect to h, and then making $h = 0$; thus

$$\frac{d.f(a+h)}{dh} = B + 2Ch + \&c. + (n - 1)Nh^{n-2} + \alpha Ph^{\alpha-1} + \&c.$$

$$\frac{d^2.f(a+h)}{dh^2} = 2C + \&c. + (n-1)(n-2)Nh^{n-3} + \alpha(\alpha-1)Ph^{\alpha-2} + \&c.$$

$$\vdots$$

$$\frac{d^{n-1}.f(a+h)}{dh^{n-1}} = \ldots(n-1)(n-2)\ldots2.1N + \alpha.(\alpha-1)\ldots(\alpha-n+2)Ph^{\alpha-n+1},$$

$$\frac{d^n.f(a+h)}{dh^n} = \alpha(\alpha-1)(\alpha-2)\ldots(\alpha-n+1).Ph^{\alpha-n} + \&c.$$

Now if $h = 0$, since $a > n-1$, but $< n$, the terms involving P will vanish from the first $(n-1)$ equations, and the $(n-1)$ differential coefficients will be found.

But since $a - n$ is negative, then when $h = 0$

$$\frac{d^n . f(a+h)}{dh^n} = \frac{a(a-1)(a-2)\ldots(a-n+1) . P}{h^{n-a}} \text{ is infinite.}$$

78. Again, should the substitution of $x = a$ introduce negative powers of h, all the differential coefficients will be infinite. This, as it has been observed, is the case, when $u = f(x)$ contains a term $\frac{A}{(x-a)^m}$, for then if x become $x + h$,

$$\frac{A}{(x-a)^m} = \frac{A}{(x+h-a)^m} = \frac{A}{h^m} \text{ when } x = a.$$

Let then $f(a+h) = Ah^{-m} + \&c.$;

$$\therefore \frac{d . f(a+h)}{dh} = \frac{-mA}{h^{m+1}} + \&c.,$$

$$\vdots \qquad \vdots$$

$$\text{and } \frac{d^n . f(a+h)}{dh^n} = \frac{-m(m+1)(m+2)\ldots(m+n-1) . A}{h^{m+n}},$$

which it is manifest becomes infinite if $h = 0$.

79. From this it is obvious that if the n^{th} differential coefficient become infinite when $x = a$, the true expansion contains a fractional power of h lying between $(n-1)$ and (n); and that if $x = a$ makes $f(x) = \infty$, the true expansion contains negative powers of h.

Ex. 1. Let $u = f(x) = b + (x-a)^{\frac{3}{2}}$; find $f(a+h)$.

$$\frac{du}{dx} = \tfrac{3}{2}(x-a)^{\frac{1}{2}},$$

$$\frac{d^2u}{dx^2} = \tfrac{3}{2} \cdot \tfrac{1}{2} \frac{1}{(x-a)^{\frac{1}{2}}} \text{ which } = \infty \text{ if } x = a;$$

$\therefore$ the fractional index of h is $< 2 > 1$.

But $u_1 = b + (x+h-a)^{\frac{3}{2}} = b + h^{\frac{3}{2}}$ when $x = a$,

and $\frac{3}{2}$ lies between 2 and 1.

Ex. 2. If $u = bx^m + c(x-a)^{\frac{p}{q}}$.

$$\frac{du}{dx} = m \cdot bx^{m-1} + \frac{cp}{q} \cdot (x-a)^{\frac{p}{q}-1},$$

$$\vdots$$

$$\frac{d^n u}{dx^n} = m(m-1)\ldots(m-n+1)\,bx^{m-n}$$

$$+ c\frac{p}{q} \cdot \left(\frac{p}{q}-1\right) \ldots \left(\frac{p}{q}-n+1\right) \cdot (x-a)^{\frac{p}{q}-n},$$

and let $\frac{p}{q} < n$ but $> n-1$. Then $\frac{d^n u}{dx^n}$ is the first differential coefficient which becomes infinite, and there ought in the true expansion to be a term involving $h^{\frac{p}{q}}$, which there is; for by putting $x+h$ for x, and afterwards writing a for x, we have $f(a+h) = b \cdot (a+h)^m + ch^{\frac{p}{q}}$.

If $m < n$, the values of the differential coefficients until we come to the n^{th}, will disappear when $x = a$.

80. In functions of this description recourse must be had to the common algebraical methods, first writing $x+h$ for x, and then putting a for x.

Thus, suppose $u = 2ax + a\sqrt{x^2 - a^2}$;

$$\therefore f(a+h) = 2a(a+h) + a\sqrt{(a+h)^2 - a^2}$$

$$= 2a(a+h) + a\sqrt{2ah + h^2}$$

$$= 2a(a+h) + a\sqrt{2ah} \cdot \left(1 + \frac{h}{2a}\right)^{\frac{1}{2}},$$

and $\left(1 + \frac{h}{2a}\right)^{\frac{1}{2}}$ is to be expanded by the Binomial Theorem.

81. The Limits of Taylor's Theorem.

If $f(x+h)$ be expanded by Taylor's Theorem, and we stop at the n^{th} term, the sum of the first n terms may differ widely from the true value of $f(x+h)$; it is therefore necessary to calculate the amount or limit of the error which arises from neglecting the remaining terms of the series.

Our present object is to ascertain these limits; but the following proposition must precede the investigation.

82. PROP. If $u = f(x) = 0$ when $x = 0$, then u and $\frac{du}{dx}$ will have the same sign while x increases from 0 to a, if a be positive; but contrary signs, if a be negative; $\frac{du}{dx}$ being supposed neither to change its sign, nor to become infinite, while x increases from 0 to a.

Let a be divided into n equal parts, each $= h$, or $a = nh$.

Then since $f(x+h) = f(x) + \frac{du}{dx}h + Ph^2$ (1);

if U_1 and P_1 be the values of $\frac{du}{dx}$ and P, when $x = 0$,

$$f(h) = U_1 h + P_1 h^2.$$

Now if $\left.\begin{matrix} U_2, & U_3, & U_4 \ldots U_n \\ P_2, & P_3, & P_4 \ldots P_n \end{matrix}\right\}$ be the values of $\frac{du}{dx}$ and P, when h, $2h$, $3h \ldots (n-1)h$ are put for x, we have from (1),

$$f(h+h) - f(h) = U_2 h + P_2 h^2,$$
$$f(2h+h) - f(h+h) = U_3 h + P_3 h^2,$$
$$\vdots \qquad\qquad \vdots$$
$$f\{(n-1)h + h\} - f\{(n-2)h + h\} = U_n h + P_n h^2;$$

whence, by addition,

$f(nh)$ or $f(a) = (U_1 + U_2 + U_3 + \&c. + U_n)h + (P_1 + P_2 + \&c. + P_n)h^2$;

and by diminishing h, the first term $(U_1 + U_2 + U_3 + \&c. + U_n)h$ may be rendered greater than the second, and therefore the algebraical sign of $f(a)$ will depend only on the first term.

Also $f(h)$ will have the same sign as U_1, which is $\frac{du}{dx}$ when $x = 0$; or, since $\frac{du}{dx}$ does not change its sign, $f(h)$ will have the same sign as $\frac{du}{dx}$. Also $f(2h) - f(h)$ will have the same sign as U_2, which is the value of $\frac{du}{dx}$ when $x = h = \frac{a}{n}$; and therefore the same sign as $\frac{du}{dx}$.

And therefore $f(a)$ which has the same sign as the sum of the products $(U_1 + U_2 + U_3 + \&c. + U_n)\frac{a}{n}$ will have the

same sign as $\frac{du}{dx}$, if a be positive, but the contrary sign if a be negative.

83. This proposition being premised, let us assume that the true value of $f(x+h)$ or u_1 lies between the values

$$u+\frac{du}{dx}h+\frac{d^2u}{dx^2}\frac{h^2}{1.2}+\&c.+\frac{d^nu}{dx^n}\cdot\frac{h^n}{1.2...n}+\frac{mh^{n+1}}{1.2.3...(n+1)},$$

and $u+\frac{du}{dx}h+\frac{d^2u}{dx^2}\frac{h^2}{1.2}+\&c.+\frac{d^nu}{dx^n}\cdot\frac{h^n}{1.2...n}+\frac{Mh^{n+1}}{1.2...(n+1)}$,

where $\frac{mh^{n+1}}{2.3...(n+1)}$ and $\frac{Mh^{n+1}}{2.3...(n+1)}$ are the least and greatest values of which the remaining part of Taylor's Theorem is capable of;

$$\therefore u_1-\left(u+\frac{du}{dx}h+\frac{d^2u}{dx^2}\frac{h^2}{1.2}+\&c.+\frac{d^nu}{dx^n}\frac{h^n}{1.2...n}\right)>\frac{mh^{n+1}}{1.2...(n+1)}$$
$$<\frac{Mh^{n+1}}{1.2.3...(n+1)};$$

$$\text{or } u_1-\left(u+\frac{du}{dx}h+\&c.+\frac{d^nu}{dx^n}\frac{h^n}{1...n}\right)-\frac{mh^{n+1}}{1.2...(n+1)}>0,$$

$$\text{and } \frac{Mh^{n+1}}{1.2.3...(n+1)}-u_1+u+\frac{du}{dx}h+\&c.+\frac{d^nu}{dx^n}\frac{h^n}{1...n}>0.$$

Now both these quantities vanish when $h=0$, since then $u_1=u$; therefore, their first differential coefficients will have the same sign: hence differentiating with respect to h,

$$\frac{du_1}{dh}-\left\{\frac{du}{dx}+\frac{d^2u}{dx^2}\cdot h+\&c.+\frac{d^nu}{dx^n}\cdot\frac{h^{n-1}}{1...(n-1)}\right\}-\frac{mh^n}{1.2...n}>0,$$

$$\text{and } \frac{Mh^n}{1...n}-\frac{du_1}{dh}+\frac{du}{dx}+\frac{d^2u}{dx^2}h+\&c.+\frac{d^nu}{dx^n}\frac{h^{n-1}}{1...(n-1)}>0.$$

Again, considering these expressions as functions of h which vanish when $h=0$, for then $\frac{du_1}{dh}=\frac{du}{dx}$; their first differential coefficients will have the same sign as the functions have, or be both greater than zero; whence again differentiating, we have

$$\frac{d^2u_1}{dh^2}-\left\{\frac{d^2u}{dx^2}+\&c.+\frac{d^{n-2}u}{dx^{n-2}}\frac{h^{n-2}}{1.2...(n-2)}\right\}-\frac{mh^{n-1}}{1.2...(n-1)}>0,$$

and $\dfrac{Mh^{n-1}}{1.2...n-1} - \dfrac{d^2u_1}{dh^2} + \dfrac{d^2u}{dx^2} + \&c. + \dfrac{d^{n-2}u}{dx^{n-2}} \cdot \dfrac{h^{n-2}}{1.2...(n-2)} > 0,$

which are both functions of h, which vanish when $h = 0$, since then, $\dfrac{d^2u_1}{dh^2} = \dfrac{d^2u}{dx^2}$, and if this process be continued $(n+1)$ times, we shall at length obtain

$$\frac{d^{n+1}u_1}{dh^{n+1}} - m > 0, \text{ and } M - \frac{d^{n+1}u_1}{dh^{n+1}} > 0;$$

$$\text{or } M - \frac{d^{n+1}u_1}{dx^{n+1}} > 0, \text{ and } \frac{d^{n+1}u_1}{dx^{n+1}} - m > 0;$$

a condition which can be satisfied by taking M equal to the greatest value of the $(n+1)^{\text{th}}$ differential coefficient, and m equal to the least value;

$$\text{or } M = \frac{d^{n+1}f(x+h)}{dx^{n+1}}, \text{ and } m = \frac{d^{n+1}f(x)}{dx^{n+1}},$$

and therefore the true value of u_1 lies between

$$u + \frac{du}{dx}h + \frac{d^2u}{dx^2}\frac{h^2}{1.2} + \&c. + \frac{d^{n+1}f(x)}{dx^{n+1}} \cdot \frac{h^{n+}}{1.2.3\,(n+1)},$$

$$\text{and } u + \frac{du}{dx}h + \frac{d^2u}{dx^2}\frac{h^2}{1.2} + \&c. + \frac{d^{n+1}f(x+h)}{dx^{n+1}} \cdot \frac{h^{n+1}}{1.2\,(n+1)},$$

and the error made by omitting the terms after the n^{th} is less than

$$\frac{(M-m)\,h^{n+1}}{1.2.3...(n+1)}.$$

Ex. 1. Let $u = x^p$;

$$\therefore \frac{d^{n+1}u}{dx^{n+1}} = p.(p-1).(p-2)...(p-n)x^{p-n-1} = m,$$

and $M = p.(p-1).(p-2)...(p-n).(x+h)^{p-n-1}$;

therefore error committed by omitting the terms of $(x+h)^p$, after the n^{th}

$$\text{is} < \frac{p.(p-1)\,(p-2).....(p-n)}{1.2.3.....(n-1)} \left\{(x+h)^{p-n-1} - x^{p-n-1}\right\}.$$

Ex. 2. Let $u = a^x$;

$$\therefore \frac{d^nu}{dx^n} = A^n a^x, \text{ and } \frac{d^nu_1}{dx^n} = A^n a^{x+h};$$

therefore the true expansion of a^{x+h} lies between the series,

$$a^x \cdot \left\{1 + Ah + \frac{A^2h^2}{1.2} + \&c. + \frac{A^nh^n}{1.2...n} + \frac{A^{n+1}h^{n+1}}{1.2...n+1}\right\},$$

and $a^x \cdot \left\{1 + Ah + \frac{A^2h^2}{1.2} + \&c. + \frac{A^nh^n}{1.2...n} + \frac{A^{n+1}h^{n+1}a^h}{1.2.3...n+1}\right\}$,

and the error committed by omitting the terms after the n^{th},

$$\text{is} < \frac{A^{n+1}h^{n+1}}{1.2...(n+1)}(a^h - 1),$$

$$< \frac{U_nA.h}{n+1}(a^h - 1), \text{ if } U_n \text{ be the } n^{th} \text{ term.}$$

Again, if U_n be the first term that converges,

$$\frac{U_n}{U_{n+1}} > 1, \quad \text{i.e. } \frac{n+1}{Ah} > 1.$$

Let $(n+1) = 2Ah$,

therefore error $< \frac{U_n}{2}.(a^h - 1)$, $< \frac{U_n}{2}.(a^{\frac{n+1}{2A}} - 1)$.

Ex. 3. Let $u = \log x$;

$$\therefore \frac{d^{n+1}u}{dx^{n+1}} = (-1)^n.1.2.3...n.x^{-(n+1)} = m,$$

and the error, by omitting the terms after the n^{th}, is

$$< (-1)^n.\frac{h^{n+1}}{n+1}.\left\{(x+h)^{-(n+1)} - x^{-(n+1)}\right\}.$$

84. Hence it would appear that there is some value which is exactly equal to the sum of the terms after the n^{th}. Let N be this value, therefore $f(x+h)$ becomes

$$u + \frac{du}{dx}h + \frac{d^2u}{dx^2}\frac{h^2}{1.2} + \&c. + \left(\frac{d^nu}{dx^n} + \frac{Nh}{n+1}\right)\frac{h^n}{1.2...n},$$

and to find h, so that $\frac{d^nu}{dx^n}\frac{h^n}{1.2...n}$ shall be greater than the remaining terms of the series, we must have

$$\frac{d^nu}{dx^n} > \frac{Nh}{n+1}, \text{ or } h < \frac{d^nu}{dx^n}.\frac{n+1}{N};$$

it is not necessary that N should be known, we may substitute for it a greater quantity, as M.

85. We may here add some remarks upon a method of notation, by which the Theorems of Taylor and Maclaurin may be put under very simple forms.

We have hitherto considered the letter d prefixed to u, as in du, d^2u, d^3u, &c. to be a symbol of *operation* and not of *quantity*, thus d, d^2 d^3, &c. indicate that u has been differentiated, once, twice, &c. But we may separate the d and its powers from u; and if we treat it as an algebraical quantity, no error can arise, so long as we bear in mind its original signification.

Thus if in Taylor's Theorem we look upon d as a factor of u, we shall have

$$u_1 = u\left\{1 + \frac{d}{dx}.h + \frac{d^2}{dx^2}\frac{h^2}{1.2} + \frac{d^3}{dx^3}\frac{h^3}{2.3} + \&c.\right\};$$

let $\frac{d}{dx} = t$, and $\therefore \frac{d^2}{dx^2} = t^2$, &c.;

$$\therefore u_1 = u\left\{1 + th + \frac{t^2h^2}{1.2} + \frac{t^3h^3}{2.3} + \&c.\right\} = ue^{th};$$

for e^{th} when expanded will produce a series of the required form, and if we take care that the powers of d be referred to operation and not to quantity, no error can arise, and thus Taylor's Theorem may be concisely written

$$u_1 = ue^{\frac{d}{dx}.h}.$$

Again in Maclaurin's Theorem if we may treat the coefficients U_0, U_1, U_2, U_3, &c. as powers U^0, U^1, U^2, U^3, &c., we have

$$u = 1 + Ux + \frac{U^2x^2}{1.2} + \frac{U^3x^3}{2.3} + \&c. = e^{Ux}.$$

Nor can error arise, if we keep in mind the original meaning of the coefficients U_0, U_1, U_2, &c., and if when we expand e^{Ux} we change the indices of U into suffixes, putting U_0 instead of unity. But the utility of this method of notation will be chiefly apparent when the reader enters upon the study of the Calculus of Finite Differences.

CHAPTER VI.

VANISHING FRACTIONS.

86. SOMETIMES the substitution of a particular value for the unknown quantity, makes both the numerator and denominator of a fraction vanish, such a fraction is called a vanishing fraction.

Thus $\frac{x^2-1}{x-1}$ becomes $\frac{0}{0}$ when $x=1$, but since by division, $\frac{x^2-1}{x-1}=x+1$; the true value of the fraction when $x=1$, is $1+1=2$.

Here both the numerator and denominator vanish if $x=1$, because they both contain the factor $x-1$, which vanishes on the same supposition.

87. That the value of the fraction $\frac{x^2-1}{x-1}$ tends to 2 as its limit as x tends to 1, may be shewn by actual substitution; put for x successively the numbers

$$3,\ 2,\ 1\tfrac{1}{2},\ 1\tfrac{1}{10},\ 1\tfrac{1}{100},\ 1\tfrac{1}{1000},\ \&c.$$

then the values of the fraction become

$$4,\ 3,\ 2\tfrac{1}{2},\ 2\tfrac{1}{10},\ 2\tfrac{1}{100},\ 2\tfrac{1}{1000},\ \&c.$$

which sufficiently shew that as x approaches unity, the value of the fraction approaches 2.

88. We proceed to shew that the values of these fractions may be finite, nothing, or infinite; and thus the term *vanishing fraction*, is used with great impropriety.

Let $u=\frac{P}{Q}$ be the fraction, and let $x=a$ be the value of x, which makes $P=0$, and $Q=0$; then P and Q are both divisible by $x-a$, or the powers of $x-a$*;

$$\therefore \text{ let } P=p.\overline{x-a}\Big|^m, \text{ and } Q=q.\overline{x-a}\Big|^n;$$

$$\therefore\ u=\frac{p}{q}.\frac{\overline{x-a}\Big|^m}{\overline{x-a}\Big|^n}.$$

* For divide P by $x-a$, and let m be the quotient and n the remainder, if any; $\therefore P=(x-a)m+n$; make $x=a$; $\therefore P=0$ by hypothesis; $\therefore 0=m(a-a)+n$; $\therefore n=0$; or P is divisible by $x-a$.

(1) Let $m = n$; $\therefore\ u = \frac{p}{q}$; and $= \frac{p'}{q'}$ when $x = a$, which is finite, since neither p nor q contain $x - a$.

(2) Let $m > n$; $\therefore\ u = \frac{p}{q} \cdot \overline{x-a}^{m-n} = 0$; if $x = a$.

(3) Let $m < n$; $\therefore\ u = \frac{p}{q} \cdot \frac{1}{\overline{x-a}^{n-m}} = \frac{1}{0} = \infty$, if $x = a$.

89. Hence it appears the true value of the fraction is found by getting rid of that power of $(x - a)$, which is common both to the numerator and denominator.

When m and n are whole numbers, the value of the fraction may be found by successive differentiation. For

Let $u = \frac{P}{Q}$; $\therefore\ uQ = P$; $\therefore\ u\frac{dQ}{dx} + Q\frac{du}{dx} = \frac{dP}{dx}$.

Let $x = a$; $\therefore\ Q = 0$, by hypothesis;

$$\therefore\ u\frac{dQ}{dx} = \frac{dP}{dx}; \quad \therefore\ u = \frac{\frac{dP}{dx}}{\frac{dQ}{dx}},$$

or is equal to the ratio of the differential coefficients of the numerator and denominator, a being put for x.

But if $x = a$, also makes $\frac{dP}{dx} = 0$ and $\frac{dQ}{dx} = 0$; then by repeating the process,

$$u = \frac{\frac{d}{dx} \cdot \left(\frac{dP}{dx}\right)}{\frac{d}{dx} \cdot \left(\frac{dQ}{dx}\right)} = \frac{\frac{d^2P}{dx^2}}{\frac{d^2Q}{dx^2}},$$

and the differentiation must be continued until one of the differential coefficients becomes finite when $x = a$.

If both be finite at the same time the fraction is finite; it is nothing, if the differential coefficient of the denominator be first finite; and infinite when that is the case with the differential coefficient of the numerator.

90. If m be a fraction, this method is inapplicable, for each differentiation diminishes m by 1, and thus the common factor $(x - a)^m$ will not disappear, and at length $x = a$ will make one of the differential coefficients infinite.

Thus, if $u = \dfrac{(x^2 - a^2)^{\frac{1}{2}}}{\sqrt{x - a}} = \sqrt{x + a} = \sqrt{2a}$;

$$\frac{dP}{dx} = \frac{x}{\sqrt{x^2 - a^2}}, \text{ and } \frac{dQ}{dx} = \frac{1}{2\sqrt{x - a}},$$

both of which become infinite when $x = a$.

In such a case we may have recourse to the following method.

Let $\dfrac{P}{Q}$ be the fraction, where P and Q both vanish when $x = a$. For x put $a + h$, and let the numerator and denominator be expanded by the Binomial Theorem according to the powers of h, the indices increasing, so that the fraction becomes

$$\frac{Ah^{\alpha} + Bh^{\beta} + Ch^{\gamma} + \&c.}{A_1h^{\alpha_1} + B_1h^{\beta_1} + C_1h^{\gamma_1} + \&c.},$$

which is of the proper form; $\because$ if $h = 0$ it becomes $\dfrac{0}{0}$.

There will be three cases, $\alpha = \alpha_1$, $\alpha > \alpha_1$, and $\alpha < \alpha_1$.

(1) If $\alpha = \alpha_1$ divide each term by h^{α}, and we have

$$\therefore \frac{A + Bh^{\beta-\alpha} + Ch^{\gamma-\alpha} + \&c.}{A_1 + B_1h^{\beta_1-\alpha} + C_1h^{\gamma_1-\alpha} + \&c.} = \frac{A}{A_1}, \text{ or finite if } h = 0.$$

(2) $\alpha > \alpha_1$, then the fraction

$$= \frac{Ah^{\alpha-\alpha_1} + Bh^{\beta-\alpha_1} + \&c.}{A_1 + B_1h^{\beta_1-\alpha_1} + \&c.} = 0, \quad \text{when } h = 0.$$

(3) $\alpha < \alpha_1$, then

$$\frac{A + Bh^{\beta-\alpha} + \&c.}{A_1h^{\alpha_1-\alpha} + B_1h^{\beta_1-\alpha} + \&c.} = \frac{A}{0} = \infty, \quad \text{when } h = 0.$$

Cor. 1. If $u = \dfrac{P}{Q} = \dfrac{\infty}{\infty}$, when $x = a$, it $= \dfrac{0}{0}$.

$$\text{For } \frac{P}{Q} = \frac{\frac{1}{Q}}{\frac{1}{P}} = \frac{\frac{1}{\infty}}{\frac{1}{0}} = \frac{0}{0}; \text{ when } x = a.$$

Cor. 2. If $u = \dfrac{1}{P} - \dfrac{1}{Q} = \dfrac{1}{0} - \dfrac{1}{0}$; it also $= \dfrac{0}{0}$.

$$\text{For } \frac{1}{P} - \frac{1}{Q} = \frac{Q - P}{PQ} = \frac{0}{0}, \text{ when } x = a.$$

Cor. 3. Next $P \times Q = 0 \times \infty = \frac{0}{0}$.

For $Q = \frac{1}{Q_1}$, if $Q_1 = 0$, when $x = a$;

$\therefore P \times Q = P \times \frac{1}{Q_1} = \frac{P}{Q_1} = \frac{0}{0}$, if $x = a$.

Ex. 1. Find the value of $u = \frac{x^3 - 1}{x^3 + 2x^2 - x - 2}$, when $x = 1$,

$P = x^3 - 1$; $\therefore \frac{dP}{dx} = 3x^2 = 3$, when $x = 1$,

$Q = x^3 + 2x^2 - x - 2$; $\therefore \frac{dQ}{dx} = 3x^2 + 4x - 1 = 6$, if $x = 1$;

$\therefore u = \frac{0}{0} = \frac{3}{6} = \frac{1}{2}$.

Ex. 2. Find the value of $\frac{a^x - b^x}{x}$, when $x = 0$.

$P = a^x - b^x$, and $Q = x$,

$\frac{dP}{dx} = a^x \log a - b^x \log b = \log a - \log b = \log \frac{a}{b}$, when $x = 0$,

and $\frac{dQ}{dx} = 1$; $\therefore u = \log \left(\frac{a}{b}\right)$.

Ex. 3. $u = \frac{x^x - x}{1 - x + \log x} = \frac{0}{0}$, if $x = 1$.

$P = x^x - x$, and $Q = 1 - x + \log x$,

$\frac{dP}{dx} = x^x(1 + \log x) - 1 = 0$, if $x = 1$,

$\frac{dQ}{dx} = -1 + \frac{1}{x} = 0$, if $x = 1$,

$\frac{d^2P}{dx^2} = x^x(1 + \log x)^2 + \frac{x^x}{x} = 2$, if $x = 1$,

$\frac{d^2Q}{dx^2} = -\frac{1}{x^2} = -1$, if $x = 1$;

$\therefore u = \frac{2}{-1} = -2$.

91. It is sometimes more convenient to dispense with differentiation.

Ex. 4. $u = \dfrac{e^x - 1 - \log(1+x)}{x^2} = \dfrac{0}{0}$, when $x = 0$.

$$e^x - 1 = x + \frac{x^2}{1.2} + \frac{x^3}{2.3} + \text{\&c.}$$

$$\log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \text{\&c.};$$

$$\therefore \frac{e^x - 1 - \log(1+x)}{x^2} = 1 - \frac{x}{6} + \text{\&c.} = 1, \text{ if } x = 0.$$

Ex. 5. If $u = \dfrac{1}{1-x} - \dfrac{2}{1-x^2} = \infty - \infty$, when $x = 1$.

$$\frac{1}{1-x} - \frac{2}{1-x^2} = \frac{1+x-2}{1-x^2} = -\frac{(1-x)}{1-x^2} = -\frac{1}{1+x} = -\tfrac{1}{2}, \text{ if } x = 1.$$

Ex. 6. If $u = \dfrac{\pi}{4x} - \dfrac{\pi}{2x(e^{\pi x}+1)}$; find it, if $x = 0$.

$$\text{Here } u = \frac{\pi}{4x} \cdot \frac{e^{\pi x} - 1}{e^{\pi x} + 1} = \frac{0}{0}, \text{ if } x = 0.$$

Expand $e^{\pi x}$ by the formula $e^z = 1 + z + \dfrac{z^2}{1.2} + \text{\&c.}$

$$u = \frac{\pi}{4x} \cdot \left(\frac{\pi x + \dfrac{\pi^2 x^2}{1.2} + \text{\&c.}}{2 + \pi x + \dfrac{\pi^2 x^2}{1.2} + \text{\&c.}} \right) = \frac{\pi}{4} \cdot \left(\frac{\pi + \dfrac{\pi^2 x}{1.2} + \text{\&c.}}{2 + \pi x + \dfrac{\pi^2 x^2}{1.2} + \text{\&c.}} \right).$$

Let $x = 0$; $\therefore u = \dfrac{0}{0} = \dfrac{\pi}{8}$.

Ex. 7. $u = \dfrac{\log x}{x}$; find it, when $x = \infty$.

Let $\log x = y$; $\therefore x = e^y$,

$$u = \frac{y}{e^y} = \frac{y}{1 + y + \dfrac{y^2}{1.2} + \dfrac{y^3}{2.3} + \text{\&c.}}$$

$$= \frac{1}{\dfrac{1}{y} + 1 + \dfrac{y}{2} + \dfrac{y^2}{2.3} + \text{\&c.}} = \frac{1}{0 + \infty} = \frac{1}{\infty} = 0, \text{ if } y = \infty.$$

Similarly, if $u = \dfrac{\log x}{x^n}$; and $x = \infty$, $u = 0$.

Ex. 8. $u = \dfrac{\sqrt{a^2 - x^2} + a - x}{\sqrt{a - x} + \sqrt{a^3 - x^3}} = \dfrac{0}{0}$, when $x = a$.

Here $\dfrac{dP}{dx}$ and $\dfrac{dQ}{dx}$ are infinite when $x = a$, and we must use the second method, and since if x be $> a$, $\sqrt{a^2 - x^2}$ is impossible; let $x = a - h$, and making the substitutions

$$u = \frac{\sqrt{2ah - h^2} + h}{\sqrt{h} + \sqrt{h(a^2 + ax + x^2)}} = \frac{\sqrt{2a - h} + \sqrt{h}}{1 + \sqrt{a^2 + ax + x^2}}.$$

Let $x = a$, or $h = 0$. Then $u = \dfrac{\sqrt{2a}}{1 + \sqrt{3a^2}}$.

We might divide at once by $\sqrt{a - x}$, and then

$$u = \frac{\sqrt{a + x} + \sqrt{a - x}}{1 + \sqrt{a^2 + ax + x^2}} = \frac{\sqrt{2a}}{1 + \sqrt{3a^2}}, \text{ when } x = a.$$

Ex. 9. $u = \dfrac{x^4 - x^3 + x^2 - 1}{x^4 - 2x^3 + 2x^2 - 1} = \frac{3}{2}$; $x = 1$.

Ex. 10. $u = \dfrac{1 - \sin x + \cos x}{\sin x + \cos x - 1} = 1$; $x = \dfrac{\pi}{2}$.

Ex. 11. $u = \dfrac{ax^2 - 2acx + ac^2}{bx^2 - 2bcx + bc^2} = \dfrac{a}{b}$; $x = c$.

Ex. 12. $u = \dfrac{\cos x - \cos mx}{\cos x - \cos nx} = \dfrac{1 - m^2}{1 - n^2}$; $x = 0$.

Ex. 13. $u = \dfrac{x^3 - x^2 - 8x + 12}{x^4 - 9x^2 + 4x + 12} = \dfrac{1}{3}$; $x = 2$.

Ex. 14. $u = \dfrac{e^x - e^{\sin x}}{x - \sin x} = 1$; $x = 0$.

Ex. 15. $u = \dfrac{\tan x - \sin x}{(\sin x)^3} = \dfrac{1}{2}$; $x = 0$.

Ex. 16. $u = \dfrac{1 - (n+1).x^n + n.x^{n+1}}{(1 - x)^2} = \dfrac{n(n+1)}{2}$; $x = 1$.

Ex. 17. $u = \dfrac{x^4 - 5x^3 + 9x^2 - 7x + 2}{x^4 - 6x^3 + 12x^2 - 10x + 3} = \frac{1}{2}$; $x = 1$.

Ex. 18. $u = \dfrac{e^{nx} - e^{na}}{(x - a)^4} = \dfrac{n^4 e^{na}}{2.3.4}$; $x = a$.

Ex. 19. $u = \dfrac{e^x - e^{-x}}{\log(1+x)} = 2; \quad x = 0.$

Ex. 20. $u = \dfrac{a - x - a\log\left(\dfrac{a}{x}\right)}{a - \sqrt{a^2 - (a-x)^2}} = -1; \quad x = a.$

Ex. 21. $u = (1-x)\tan\dfrac{\pi x}{2} = \dfrac{1-x}{\cot\dfrac{\pi x}{2}} = \dfrac{2}{\pi}; \quad x = 1.$

Ex. 22. $u = (1-x)\log(1-x) = 0; \quad x = 1.$

Ex. 23. $u = \dfrac{\sqrt{2a^3x - x^4} - a\sqrt[3]{a^2x}}{a - \sqrt[4]{ax^3}} = \dfrac{16a}{9}; \quad x = a.$

Ex. 24. $u = \dfrac{1}{2x^2} - \dfrac{\pi}{2x\tan\pi x} = \dfrac{\pi^2}{6}; \quad x = 0.$

Ex. 25. $u = \dfrac{1 + x - x^2 - x^3}{1 + 2x + 2x^2 + 2x^3 + x^4} = 1; \quad x = -1.$

Ex. 26. $u = \dfrac{x^4 + 3x^3 - 7x^2 - 27x - 18}{x^4 - 3x^3 - 7x^2 + 27x - 18}$

$= 10;$ if $x = 3;$ $= \dfrac{1}{10};$ if $x = -3.$

Ex. 27. $u = \dfrac{xe^{2x} + 1 - e^{2x} - x}{e^{2x} - 1} = -1; \quad x = 0.$

Ex. 28. $u = \dfrac{x^3 - 4ax^2 + 7a^2x - 2a^3 - 2a^2\sqrt{2ax - a^2}}{x^2 - 2ax - a^2 + 2a\sqrt{2ax - x^2}}$

$= -5a; \quad x = a.$

Ex. 29. $u = \dfrac{x}{x-1} - \dfrac{1}{\log x} = \frac{1}{2}; \quad x = 1.$

Ex. 30. $u = \dfrac{a^{\log x} - x}{\log x} = \log\left(\dfrac{a}{e}\right); \quad x = 1.$

Ex. 31. $u = \dfrac{\pi}{4x}.\tan\dfrac{\pi x}{2} = \dfrac{\pi^2}{8}; \quad x = 0.$

Ex. 32. If $u^4 - 96a^2u^2 + 100a^2x^2 - x^4 = 0,$

if $x = 0;$ $\dfrac{du}{dx} = \pm\dfrac{5}{4}\sqrt{\dfrac{2}{3}}.$

CHAPTER VII.

MAXIMA AND MINIMA.

92. If $u = f(x)$ express the relation between the function u, and the variable x, then if $x = a$ make $f(a)$ greater than *both* $f(a + h)$ and $f(a - h)$; $u = f(a)$ is said to be a maximum; but if $f(a)$ be less than *both* $f(a + h)$ and $f(a - h)$, it is called a minimum.

Hence the value of a function is said to be a *maximum* or *minimum*, according as the particular value is greater or less than the values which immediately precede and follow it.

From this definition it appears, that if a quantity either continually increase or constantly decrease, it does not possess the property of a maximum or minimum. Also as the words maximum or minimum are used in a relative and not in an absolute sense, functions may possess many maxima or minima.

For we may easily conceive that a quantity after having reached a maximum value may decrease to a minimum value, and afterwards again increase, and thus many maxima and minima may exist in the same function, but which it is obvious must succeed in order.

Thus the alternate elevation and depression of the waves of the sea will with regard to a horizontal line give maxima and minima altitudes.

93. In the circle the sine* which $= 0$, when the arc $= 0$, increases as the arc increases, till the arc $= 90^0$, when the sine $=$ radius, from this value it decreases, till at the end of the second quadrant it becomes $= 0$.

At 90^0, therefore, it is a maximum; for any two sines drawn on opposite sides of the sin 90^0, and equidistant from it, will be both less than the radius.

In the parabola, the line drawn from the focus to the vertex, is less than either of two focal distances which can be drawn to the curve on opposite sides of it; it is therefore a minimum.

* By the sine is here meant the semichord to which the sine of the angle is proportional.

By reference to figures 1 and 2, we perceive that

NP in fig. 1, is a maximum,

NP.....2, is a minimum.

(1)



(2)



94. One of the chief applications of the Differential Calculus, is that which affords rules for the discovery of these values.

But the following proposition must first be established.

If $y = A_1h + A_2h^2 + A_3h^3 + \&c. + A_nh^n + A_{n+1}h^{n+1} + \&c.$, where the ratio of any coefficient to the one immediately preceding is finite, i.e. $\frac{A_{n+1}}{A_n}$ is finite, h may be so assumed that any one term shall be greater than the sum of all the terms that follow it.

Let r be $>$ the greatest ratio between the coefficients;

$$\therefore \frac{A_2}{A_1} < r, \quad \text{or} \quad A_2 < A_1 r,$$

$$\frac{A_3}{A_2} < r; \quad \therefore A_3 < A_1 r^2,$$

$$\frac{A_4}{A_3} < r; \quad \therefore A_4 < A_1 r^3,$$

&c.

$$\therefore A_1h + A_2h^2 + A_3h^3 + \&c. < A_1h + A_1rh^2 + A_1r^2h^3 + \&c.$$

$$< A_1h\{1 + rh + r^2h^2 + \&c.\}$$

$$< A_1h\frac{1}{1-rh}.$$

Let $rh = \frac{1}{2}$, or $h = \frac{1}{2r}$; $\therefore \frac{1}{1-rh} = 2$;

$$\therefore A_1h + A_2h^2 + A_3h^3 + \&c. < 2A_1h;$$

$$\therefore A_2h^2 + A_3h^3 + \&c. < A_1h;$$

similarly may A_2h^2 be shewn to be $> A_3h^3 + A_4h^4 + \&c.$

We have here supposed the series to proceed to infinity: if it extend to n terms, it is evident, *à fortiori* that any one term is greater than the sum of all that follow it.

In two of the following series the theorem is, and in one it is not applicable.

$$x + 2x^2 + 3x^3 + 4x^4 + 5x^5 + \&c. \ldots\ldots\ldots(1),$$

$$x + 1.2x^2 + 2.3x^3 + 2.3.4x^4 + \&c. \ldots\ldots(2),$$

$$x + 2^2.x^2 + 3^2.x^3 + 4^2.x^4 + \&c. \ldots\ldots\ldots\ldots(3).$$

95. PROP. If $u = f(x)$ be a maximum or minimum when $x = a$. Then on the same supposition, $\frac{du}{dx} = 0$.

Let $u_1 = f(x + h)$, and $u_2 = f(x - h)$.

Now at a maximum or minimum, $u = f(x)$ must be greater or less than both $f(x + h)$, and $f(x - h)$, or greater or less than both u_1 and u_2, and hence, $u_1 - u$ and $u_2 - u$ must both have the same algebraical sign.

$$\text{But } u_1 - u = \frac{du}{dx}h + \frac{d^2u}{dx^2}\frac{h^2}{1.2} + \frac{d^3u}{dx^3}\frac{h^3}{2.3} + \&c.$$

$$\text{and } \therefore u_2 - u = -\frac{du}{dx}h + \frac{d^2u}{dx^2}\frac{h^2}{1.2} - \frac{d^3u}{dx^3}\frac{h^3}{2.3} + \&c.$$

by writing $-h$ for h in the value of $u_1 - u$.

Hence, since the first term of the expansion can be made greater than the sum of all the terms that follow it, (if $x = a$, does not make any of the differential coefficients infinite,) it is clear that whilst the term $\frac{du}{dx}h$ exists, $u_1 - u$ and $u_2 - u$ will have a different algebraical sign: i. e. u_1 and u_2 cannot be both greater or both less than u. Therefore, if there be a maximum or minimum, $\frac{du}{dx} = 0$, and

$$u_1 - u = \frac{d^2u}{dx^2}\frac{h^2}{1.2} + \frac{d^3u}{dx^3}\frac{h^3}{2.3} + \&c.$$

$$u_2 - u = \frac{d^2u}{dx^2}\frac{h^2}{1.2} - \frac{d^3u}{dx^3}\frac{h^3}{2.3} + \&c.$$

Now if $x = a$ does not make $\frac{d^2u}{dx^2} = 0$, the sign of $u_1 - u$ and $u_2 - u$, since h^2 is positive, will depend upon that of $\frac{d^2u}{dx^2}$.

If $\therefore \frac{d^2u}{dx^2}$ be +, $u_1 - u$ and $u_2 - u$ are both +.

If $\frac{d^2u}{dx^2}$ be −, $u_1 - u$ and $u_2 - u$ are both −.

If $\frac{d^2u}{dx^2}$ be +, u_1 and u_2 are both $> u$; or u is a minimum; if $\frac{d^2u}{dx^2}$ be −, u_1 and u_2 are both $< u$; or u is a maximum. Hence this rule. To find whether $u = f(x)$ contain any maxima or minima, put $\frac{du}{dx} = 0$, substitute the values of x thus found in $\frac{d^2u}{dx^2}$, if the results be positive, we have minima; if negative, maxima.

96. Should however $\frac{d^2u}{dx^2} = 0$ when $x = a$,

$$u_1 - u = + \frac{d^3u}{dx^3}\frac{h^3}{2.3} + \&c.$$

$$u_2 - u = - \frac{d^3u}{dx^3}\frac{h^3}{2.3} + \&c.$$

and $u_1 - u$ and $u_2 - u$ have again different signs; and therefore there will be no maxima or minima if $\frac{d^3u}{dx^3}$ exist. Hence it is obvious that we can have a maximum or minimum only when the *first differential coefficient that does not vanish is of an even order.*

Cor. 1. If u = maximum or minimum, any constant multiple of u is a maximum or minimum. For if $\frac{du}{dx} = 0$, $a\frac{du}{dx}$ is also $= 0$; and therefore if u = maximum, au is also a maximum.

Cor. 2. If $f(x)$ be a maximum or minimum, $\{f(x)\}^n$, where n is integral, is also a maximum or minimum.

For let $u = f(x)$, and $U = \{f(x)\}^n$; but $\frac{du}{dx} = f'(x) = 0$; $\therefore$ u is a maximum or minimum.;

$\therefore \frac{dU}{dx} = n\{f(x)\}^{n-1} f'(x) = 0$; or U is a maximum or minimum.

Cor. 3. If $u = f(x)$ be a maximum or minimum, $\log u$ is sometimes a maximum or minimum.

Let $U = \log u$; $\therefore \frac{dU}{dx} = \frac{1}{u} \cdot \frac{du}{dx}$. But $\frac{du}{dx} = 0$; $\therefore \frac{dU}{dx} = 0$, or U is a maximum or minimum, unless $x = a$ makes $u = 0$.

Cor. 4. If u = maximum, $\frac{1}{u}$ is a minimum, and conversely.

For let $v = \frac{1}{u}$; $\therefore \frac{dv}{dx} = -\frac{1}{u^2} \frac{du}{dx}$,

$\frac{d^2v}{dx^2} = \frac{2}{u^3} \cdot \frac{du^2}{dx^2} - \frac{1}{u^2} \cdot \frac{d^2u}{dx^2} = -\frac{1}{u^2} \cdot \frac{d^2u}{dx^2}$, when u = maximum.

Therefore, if $\frac{d^2u}{dx^2}$ be negative, $\frac{d^2v}{dx^2}$ is positive, or if u be a maximum, $\frac{1}{u}$ is a minimum.

97. In the succeeding examples the following results, the principal of which the reader may have met with before, will be found useful.

(1^0). Let a = radius of a circle, then

Area $= \pi a^2$; circumference $= 2\pi a$.

Area of sector of a circle $= \frac{1}{2}$ rad $\times$ arc.

Solidity of sphere $= \frac{4}{3}\pi a^3$.

The surface of sphere $= 4\pi a^2$.

Surface of a segment $= 2\pi a x$; where x is altitude of segment.

(2^0). Let $2a$, and $2b$ be the axes of an ellipse;

$\therefore$ area of ellipse $= \pi ab$.

(3^0). Let a = axis, $2b$ greatest double ordinate of parabola;

$\therefore$ area $= \frac{2}{3} a \times 2b = \frac{4}{3} ab$.

Solidity of paraboloid $= \frac{1}{2}\pi b^2 a$.

(4^0). Let a = altitude; b = radius of base of cylinder.

Solidity of cylinder $= \pi b^2 a$.

Convex surface $= 2\pi ab$.

Whole surface $= 2\pi b^2 + 2\pi ab$.

(5^0). Let a = altitude, and b = radius of base of cone.

Solidity of cone $= \frac{1}{3}\pi b^2 a$.

Convex surface* $= \pi b\sqrt{b^2 + a^2}$.

Whole surface $= \pi b^2 + \pi b\sqrt{b^2 + a^2}$.

EXAMPLES.

(1) Let $u = x^3 - 6x^2 + 11x - 6$; find the values of x which make u a maximum or minimum.

$$\frac{du}{dx} = 3x^2 - 12x + 11 = 0;$$

$$\therefore\ x^2 - 4x + 4 = \tfrac{1}{3};\quad \therefore\ x = 2 \pm \frac{1}{\sqrt{3}} = 2 \pm \frac{\sqrt{3}}{3},$$

$$\frac{d^2u}{dx^2} = 6x - 12.$$

Let $x = 2 + \frac{\sqrt{3}}{3}$; $\therefore\ \frac{d^2u}{dx^2} = 2\sqrt{3}$ indicates a minimum,

$x = 2 - \frac{\sqrt{3}}{3}$; $\therefore\ \frac{d^2u}{dx^2} = -2\sqrt{3}$.....a maximum.

(2) Let $y = x\tan\theta - \frac{x^2}{4h\cos^2\theta}$; find x that y may be a maximum or minimum.

$$\frac{dy}{dx} = \tan\theta - \frac{x}{2h\cos^2\theta};\quad \frac{d^2y}{dx^2} = -\frac{1}{2h\cos^2\theta}.$$

From $\frac{dy}{dx} = 0$, $x = 2h\tan\theta\cos^2\theta = 2h\sin\theta\cos\theta$;

also $\frac{d^2y}{dx^2}$ is negative; $\therefore\ y$ is a maximum,

* The surface of a cone when unwrapped becomes the sector of a circle, of which the centre is the vertex of the cone, and radius the slant side, and arc the circumference of the base of the cone.

But sector $= \frac{1}{2}$ rad $\times$ arc; and radius $= \sqrt{b^2+a^2}$; arc $= 2\pi b$;

$\therefore$ Convex surface $= \pi b\sqrt{b^2+a^2}$.

$$\text{and } y = 2h\tan\theta \,.\, \sin\theta\cos\theta - \frac{4h^2\sin^2\theta\cos^2\theta}{4h\cos^2\theta}$$

$$= 2h\sin^2\theta - h\sin^2\theta = h\sin^2\theta.$$

This is the equation to the path of a projectile, and the maximum value of y is the greatest height above the horizontal plane.

(3) $u = (\sin x)^m \,.\, \{\sin(a-x)\}^n$; find x that u may be a maximum or minimum.

$$\frac{du}{dx} = m(\sin x)^{m-1}\sin(a-x)^n \,.\, \cos x$$

$$- n(\sin x)^m \,.\, \sin(a-x)^{n-1}\cos(a-x) = 0;$$

$$\therefore m\sin(a-x) \,.\, \cos x - n\sin x\cos(a-x) = 0;$$

$$\therefore \frac{\sin(a-x) \,.\, \cos x}{\cos(a-x) \,.\, \sin x} = \frac{m}{n};$$

$$\therefore \frac{\sin(a-x)\cos x - \cos(a-x)\sin x}{\sin(a-x)\cos x + \sin x\cos(a-x)} = \frac{n-m}{n+m};$$

$$\therefore \frac{\sin(a-2x)}{\sin a} = \frac{n-m}{n+m};$$

$a - 2x$ and $\therefore x$ may be found from the tables.

(4) $u = \dfrac{\log x}{x}$; find x that u may be a maximum.

$$\frac{du}{dx} = \frac{1-\log x}{x^2} = 0; \quad \therefore \log x = 1 = \log e;$$

$$\therefore x = e, \quad \text{and } u = \frac{1}{e}.$$

(5) Find that fraction which exceeds its second power by the greatest possible number.

Let x be the fraction;

$$\therefore u = x - x^2 \text{ is a maximum;}$$

$$\therefore \frac{du}{dx} = 1 - 2x = 0; \quad \therefore x = \tfrac{1}{2},$$

$$\frac{d^2u}{dx^2} = -2, \quad \text{or } x = \tfrac{1}{2}, \text{ is a maximum.}$$

(6) Find the distance of P from A, when $\angle CPB$ is a maximum.



$$AB = a, \qquad AP = x,$$

$$AC = b, \qquad \angle CPB = \theta;$$

$$\therefore \theta = \angle CPA - \angle BPA$$

$$= \tan^{-1}\frac{b}{x} - \tan^{-1}\frac{a}{x},$$

$$\therefore \frac{d\theta}{dx} = -\frac{\frac{b}{x^2}}{1+\frac{b^2}{x^2}} + \frac{\frac{a}{x^2}}{1+\frac{a^2}{x^2}} = \frac{a}{x^2+a^2} - \frac{b}{x^2+b^2} = 0;$$

$$\therefore (a-b)x^2 = a^2b - ab^2; \quad \therefore x = \sqrt{ab};$$

$\therefore AP$ touches the circle circumscribing the $\Delta\ PBC$.

(7) Of all triangles upon the same base, and having the same perimeter, the isosceles has the greatest area;

$2P$ the perimeter and a the given base,

x and y the remaining sides;

$$\therefore \text{area} = \sqrt{P.(P-a).(P-x).(P-y)};$$

and $\because P$ and $P-a$ are constant, and if $\sqrt{u}$ be a maximum, u is also a maximum.

$$\text{Let } u = (P-x).(P-y).$$

$$\text{But } P-y = P-(2P-a-x) = a+x-P;$$

$$\therefore u = (P-x).(a+x-P),$$

$$\frac{du}{dx} = -(a+x-P) + P - x = 0;$$

$$\therefore -a-2x+2P = 0; \quad \therefore x = P - \frac{a}{2};$$

$$y = 2P - (a+x) = 2P - \left(P + \frac{a}{2}\right) = P - \frac{a}{2};$$

and therefore $x = y$, or the triangle is isosceles.

Since $\frac{d^2u}{dx^2} = -2$, the triangle is a maximum,

$$\text{and area} = \frac{a}{2}\sqrt{P.(P-a)}.$$

(8) Divide a number a into two such parts, that the product of the m^{th} power of the one into the n^{th} power of the other may be a maximum.

x one part; $a - x$ the other; $u = x^m.(a-x)^n$;

$$\therefore \frac{du}{dx} = mx^{m-1}.(a-x)^n - x^m n.(a-x)^{n-1}$$

$$= x^{m-1}.(a-x)^{n-1}\{ma - (m+n).x\} = 0,$$

whence $x = 0$, $x = a$, and $x = \frac{ma}{m+n}$;

$$\frac{d^2u}{dx^2} = \{(m-1).x^{m-2}.(a-x)^{n-1} - (n-1).x^{m-1}.(a-x)^{n-2}\}$$

$$\{ma - (m+n).x\} - (m+n).x^{m-1}.(a-x)^{n-1},$$

which vanishes when $x = 0$ and $x = a$, but if $x = \frac{ma}{m+n}$,

$$\frac{d^2u}{dx^2} = -(m+n).\left(\frac{ma}{m+n}\right)^{m-1}.\left(\frac{na}{m+n}\right)^{n-1};$$

$$\therefore x = \frac{ma}{m+n} \text{ gives } u = \text{maximum.}$$

$x = 0$ and $x = a$ will give no results unless m and n are even.

And then $\frac{d^m u}{dx^m} = m.(m-1)(m-2)\ldots 2.1.(a-x)^n + \phi(x)$,

$$\frac{d^n u}{dx^n} = n.(n-1)(n-2)\ldots 2.1.x^m + \phi(a-x),$$

and $\frac{d^m u}{dx^m} = m.(m-1)(m-2)\ldots 2.1.a^n$, when $x = 0$,

and $\frac{d^n u}{dx^n} = n.(n-1)(n-2)\ldots 2.1.a^m$, when $x = a$;

both of which correspond to minima.

(9) $u^3 - 3aux + x^3 = 0$; find x when u is a maximum.

We must differentiate the implicit function;

$$\therefore \frac{du}{dx}.(u^2 - ax) - au + x^2 = 0.$$

$$\text{But } \frac{du}{dx} = 0; \quad \therefore x^2 - au = 0, \text{ or } u = \frac{x^2}{a}.$$

Substitute in the original equation,

$$\frac{x^6}{a^3} - 3x^3 + x^3 = 0 \ldots(1); \quad \therefore\ x^3 = 2a^3; \quad \therefore\ x = a\,.\,\sqrt[3]{2}.$$

Differentiating a second time,

$$\frac{d^2u}{dx^2}(u^2 - ax) + \frac{du}{dx}\,.\,(2u\frac{du}{dx} - a) - a\frac{du}{dx} + 2x = 0.$$

But $\frac{du}{dx} = 0$, and $u^2 - ax = \frac{x^4}{a^2} - ax = \frac{x}{a^2}(x^3 - a^3) = ax$;

$$\therefore\ \frac{d^2u}{dx^2} = \frac{-2x}{ax} = -\frac{2}{a},$$

whence $x = a\sqrt[3]{2}$, gives $u = a\sqrt[3]{4}$, a maximum.

But also from equation (1) $x = 0$; and $\therefore\ u = 0$.

$$\text{Now } \frac{d^2u}{dx^2} = \frac{-2x}{u^2 - ax} = \frac{0}{0}; \text{ if } x = 0.$$

Treating the fraction as a vanishing one,

$$\frac{d^2u}{dx^2} = \frac{-2}{2u\frac{du}{dx} - a} = \frac{2}{a}, \quad \text{when } x = 0;$$

$$\therefore\ x = 0 \text{ gives } u = 0, \text{ a minimum.}$$

(10) Bisect a triangle by the shortest line.



ABC the triangle, and PQ the shortest line.

$CP = x$, $CQ = y$, $PQ = u$ } a, b, c the three sides of the triangle, C the $\angle BCA$.

Then $\because\ \Delta ABC = 2\,\Delta CPQ$;

$$\therefore\ \frac{ab\sin C}{2} = 2\,.\,\frac{xy\sin C}{2} = xy\sin C; \quad \therefore\ ab = 2xy,$$

$$u^2 = x^2 + y^2 - 2xy\cos C = x^2 + \frac{a^2b^2}{4x^2} - ab\cos C = \text{minimum};$$

$$\therefore\ 2u\frac{du}{dx} = 2x - \frac{a^2b^2}{2x^3} = 0;$$

$$\therefore\ x^4 = \frac{a^2b^2}{4}, \text{ or } x = \sqrt{\frac{ab}{2}}, \text{ and } y = \frac{ab}{2x} = \sqrt{\frac{ab}{2}};$$

$$\therefore u^2 = \frac{ab}{2} + \frac{ab}{2} - ab\cos C = ab.(1-\cos C) = ab.\left\{\frac{2ab + c^2 - (a^2+b^2)}{2ab}\right\}$$

$$= \frac{c^2 - (a-b)^2}{2};$$

$$\therefore u = \sqrt{\frac{(c-a+b)(c+a-b)}{2}}.$$

(11) Describe about a given circle ABC, the least isosceles triangle.



DPQ the triangle, DP touching the circle at A.

$DO = x$; $OA = a$; $\therefore DA = \sqrt{x^2 - a^2}$

Now $PB = \frac{DB}{DA} OA = \frac{(x+a) \times a}{\sqrt{x^2 - a^2}}$;

$\therefore \Delta DPQ = PB \times DB$

$$= \frac{a(x+a)}{\sqrt{x^2-a^2}} \times (x+a) = a \cdot \frac{(x+a)^{\frac{3}{2}}}{\sqrt{x-a}} = \text{minimum.}$$

Whence, if $u = \frac{(x+a)^3}{x-a}$, $x = 2a$, and $\Delta = a^2.3\sqrt{3}$.

(12) Find the greatest area that can be included by four given straight lines.



Let a, b, c, d be the four lines,

θ the $\angle$ included by a, b,

ϕ ... $\angle$ ..... c, d,

D the diagonal subtending the two angles, and dividing the quadrilateral into two Δs;

$$\therefore u = \text{area} = \frac{ab.\sin\theta}{2} + \frac{cd.\sin\phi}{2} = \text{maximum};$$

$$\therefore \frac{du}{d\theta} = \tfrac{1}{2}.\left(ab.\cos\theta + cd.\cos\phi.\frac{d\phi}{d\theta}\right) = 0.$$

But $c^2 + d^2 - 2cd\cos\phi = D^2 = a^2 + b^2 - 2ab.\cos\theta$;

$$\therefore cd.\frac{d\phi}{d\theta} = ab.\frac{\sin\theta}{\sin\phi};$$

$\therefore$ substituting this value, and dividing by $\frac{ab}{2}$,

$$\cos\theta + \cos\phi \cdot \frac{\sin\theta}{\sin\phi} = 0;$$

$$\therefore \sin\phi \cdot \cos\theta + \sin\theta \cdot \cos\phi = \sin(\phi + \theta) = 0 = \sin\pi;$$

$$\therefore \phi + \theta = \pi,$$

or the quadrilateral is one which may be inscribed in a circle,

and $u = \frac{ab + cd}{2} \sin\theta = \sqrt{(P-a)(P-b)(P-c)(P-d)}$,

where $P = \frac{a+b+c+d}{2}$.

(13) Through a given point D to draw the shortest line PQ terminated by two given straight lines AB, AC; the angle A being a right angle.



$AQ = x$; $AE = a$; $u = PQ$,

$AP = y$; $ED = b$;

$\therefore u = \sqrt{x^2 + y^2}$, a minimum...(1),

$\frac{a}{x} + \frac{b}{y} = 1$; the equation to PQ...(2).

From (1) $x\,dx + y\,dy = 0$; from (2) $\frac{a\,dx}{x^2} + \frac{b\,dy}{y^2} = 0$;

$$\therefore dx = -\frac{y}{x}dy; \quad \therefore \frac{a}{x^2}\cdot\frac{y}{x} - \frac{b}{y^2} = 0; \quad y = x\sqrt[3]{\frac{b}{a}};$$

$$\therefore \frac{a}{x} + \frac{\sqrt[3]{ab^2}}{x} = 1; \quad \therefore x = \sqrt[3]{a}\,.\,(a^{\frac{2}{3}} + b^{\frac{2}{3}});$$

$$\therefore y = \sqrt[3]{b}\,(a^{\frac{2}{3}} + b^{\frac{2}{3}}); \quad \therefore u^2 = (a^{\frac{2}{3}} + b^{\frac{2}{3}})(a^{\frac{2}{3}} + b^{\frac{2}{3}})^2;$$

$$\therefore u^{\frac{2}{3}} = a^{\frac{2}{3}} + b^{\frac{2}{3}}.$$

(14) Cut the greatest ellipse from a given cone. ABD the cone. PB the elliptic section.



$AC = a$, $CN = x$,

$BC = \beta$, $NP = y$,

PB the axis-major $= 2a$,

and axis minor $= 2b$.

Now area of ellipse $= \pi ab$.

And $2b = \sqrt{PQ \times BD} = \sqrt{2x \times 2\beta} = 2\sqrt{\beta x}$,

$$2a = \sqrt{BN^2 + NP^2} = \sqrt{(\beta + x)^2 + NP^2}.$$

But $NP = CA \times \frac{DN}{CD} = a \cdot \frac{\beta - x}{\beta}$;

$$\therefore\ 2a = \sqrt{(\beta + x)^2 + \frac{a^2}{\beta^2}(\beta - x)^2};$$

$$\therefore \text{ area } = \pi \frac{\sqrt{\beta x}}{2} \cdot \sqrt{(\beta + x)^2 + \frac{a^2}{\beta^2}(\beta - x)^2};$$

$$\therefore\ u = x\left\{(\beta + x)^2 + \frac{a^2}{\beta^2} \cdot (\beta - x)^2\right\}, \text{ a maximum};$$

$$\therefore\ (\beta + x)^2 + \frac{a^2}{\beta^2}(\beta - x)^2 + 2x\left\{\beta + x - \frac{a^2}{\beta^2}(\beta - x)\right\} = 0;$$

whence $$\frac{3(a^2 + \beta^2)x^2}{\beta^2} - \frac{4(a^2 - \beta^2)x}{\beta} = -(a^2 + \beta^2);$$

$$\therefore\ x^2 - \frac{4}{3}\frac{(a^2 - \beta^2)}{(a^2 + \beta^2)}\beta x = -\frac{\beta^2}{3};$$

$$\therefore\ x = \frac{2}{3} \cdot \beta \cdot \frac{(a^2 - \beta^2)}{a^2 + \beta^2} \pm \sqrt{\frac{4}{9}\beta^2 \cdot \left(\frac{a^2 - \beta^2}{a^2 + \beta^2}\right)^2 - \frac{\beta^2}{3}}$$

$$= \frac{2\beta(a^2 - \beta^2) \pm \beta\sqrt{a^4 - 14\beta^2 a^2 + \beta^4}}{3(a^2 + \beta^2)};$$

and the problem is possible if $a^4 - 14\beta^2 a^2 + \beta^4$ is positive. The limit of possibility is when the surd disappears.

Then $a^4 - 14\beta^2 a^2 + 49\beta^4 = 48\beta^4$;

$$\therefore\ a^2 = 7\beta^2 \pm \sqrt{48\beta^4} = \beta^2\{7 \pm 4\sqrt{3}\};$$

$$\therefore\ a = \beta(2 \pm \sqrt{3}),$$

and $$x = \frac{2\beta}{3} \cdot \frac{6 \pm 4\sqrt{3}}{8 \pm 4\sqrt{3}} = \frac{\beta}{3} \cdot \frac{3 + 2\sqrt{3}}{2 + \sqrt{3}} = \frac{\beta}{\sqrt{3}}.$$

(15) The content of a cone being given, find its form when its surface is a maximum.

x the altitude, and y the radius of the base.

Let $\frac{\pi a^3}{3}$ be the given content $= \therefore \frac{\pi y^2 x}{3}$.

Then u = surface = convex surface + base;

$$\therefore u = \pi y \sqrt{x^2 + y^2} + \pi y^2.$$

But $y^2 = \frac{a^3}{x}$; $\therefore y^2 + x^2 = \frac{a^3 + x^3}{x}$;

$$\therefore u = \pi a^{\frac{1}{2}} \left\{ \frac{\sqrt{a^3 + x^3} + a^{\frac{3}{2}}}{x} \right\};$$

whence because $\frac{du}{dx} = 0$; $x^3 - 2a^3 = 2a^{\frac{3}{2}} \sqrt{x^3 + a^3}$;

$$\therefore (x^3 + a^3) - 2a^{\frac{3}{2}} \sqrt{x^3 + a^3} + a^3 = 4a^3;$$

$$\therefore \sqrt{x^3 + a^3} = 3a^{\frac{3}{2}}; \quad \therefore x^3 = 8a^3, \text{ and } x = 2a,$$

$$\text{and } y^2 = \frac{a^3}{x} = \frac{a^2}{2}, \text{ or } y = \frac{a}{\sqrt{2}},$$

$$\text{and } u = \frac{\pi a}{\sqrt{2}} \cdot \sqrt{4a^2 + \frac{a^2}{2}} + \frac{\pi a^2}{2} = 2\pi a^2.$$

98. The following examples are added by way of exercise, the principal steps to their solution being indicated.

(1) Let $u = x^3 - 7x^2 + 8x + 32$.

$x = 4$; $u = 16$ a minimum;

$x = \frac{2}{3}$; $u = 34\frac{14}{27}$ a maximum.

(2) $u = x^3 - 3x^2 - 9x + 30$.

$x = -1$, gives $u = 35$ a maximum;

$x = 3$, gives $u = 3$ a minimum.

(3) $u = \sin^3 x \,.\, \cos x$; $x = 60$; $u =$ a maximum.

(4) Divide a line into two such parts, that their product multiplied by the difference of their squares shall be a maximum.

Let $2a$ be the line, $a + x$ and $a - x$ the parts;

$$\therefore u = (a^2 - x^2) \,.\, 4ax = \text{maximum, whence } x = \frac{a}{\sqrt{3}}.$$

(5) Divide a number a into two such factors, that the sum of their squares shall be a minimum; $x = \sqrt{a}$.

(6) $u = x^{\frac{1}{x}}$; $x = e$; $u = e^{\frac{1}{e}}$ a maximum.

(7) Into how many equal parts must a number a be divided that their continued product may be the greatest possible? $x = \frac{a}{e}$; $u = e^{\frac{a}{e}}$.

(8) Let $u = (mx + n) \cdot (ny + m)$ be a maximum, and $a^{mx} \cdot b^{ny} = c$; find x.

Here $\frac{du}{dx} = m(ny + m) + n(mx + n)\frac{dy}{dx} = 0$;

and because $mx \log a + ny \log b = \log c$, $\frac{dy}{dx}$ may be found,

and $x = \frac{\log\left(\frac{cb^m}{a^n}\right)}{\log(a^{2m})}$.

(9) $u = \frac{\sin x}{1 + \tan x}$ = a maximum; if $x = 45^0$.

(10) $u = \frac{(\tan x)^3}{\tan 3x}$ = a minimum; if $x = 22^0.30'$.

(11) $u = \sec x \cdot (a - b\tan x)^2$ a maximum or minimum.

$$\tan x = \frac{1}{6b}(a \pm \sqrt{a^2 - 24b^2}),$$

the upper sign gives a maximum; the lower a minimum.

(12) $u = 3x^4 - 28ax^3 + 84a^2x^2 - 96a^3x + 48b^4$.

$x = a$; $u = 48b^4 - 37a^4$ a minimum;

$x = 2a$; $u = 48b^4 - 32a^4$ a maximum;

$x = 4a$; $u = 48b^4 - 64a^4$ a minimum.

(13) $u = x(a - x)^2(2a - x)^3$.

$x = \frac{a}{6}(5 - \sqrt{13})$; u is a maximum;

$x = a$; u is a minimum;

$x = \frac{a}{6}(5 + \sqrt{13})$; u is a maximum.

(14) $au^3 - u^2x^2 + x^4 = 0$; u = a minimum; find x.

$x = 2a\sqrt{2}$; $u = 4a$.

(15) Inscribe the greatest rectangle in a given triangle.



$AD = a$, $BC = b$, $AN = x$; $\therefore Pp = \frac{bx}{a}$;

$u = \frac{bx}{a}(a - x)$; $\quad \therefore x = \frac{a}{2}$.

(16) Inscribe the greatest isosceles triangle in a given circle.

Let a = radius, the triangle is equilateral, side $= a\sqrt{3}$, area $= \frac{a^2 3\sqrt{3}}{4}$.

(17) Inscribe the greatest parallelogram within a given triangle ABC, A being one of the angles of the parallelogram. $AE = \frac{1}{2}AB$, then AE is one of the sides.

(18) Of all equiangular and isoperimetrical parallelograms, the equilateral has the greatest area.

(19) Of all triangles on the same base, and having equal vertical angles, the isosceles has the greatest perimeter.

(20) Given the base and vertical angle of a triangle, shew that when it is isosceles its area is a maximum.

(21) Of all triangles on the same base and having the same area, the isosceles has the least perimeter.

(22) Inscribe the greatest rectangle in a semicircle.



$CN = x$, $\quad CA = a$, $\quad NP = \sqrt{a^2 - x^2}$;

$\therefore u = 2PM \cdot CM = 2x\sqrt{a^2 - x^2}$;

$$\therefore x = \frac{a}{\sqrt{2}}, \quad \text{and } u = a^2.$$

(23) The same construction applies to any curve.

Let $AC = b$, $AM = x$; $\therefore PM = f(x)$, and $u = 2(b - x) \cdot f(x)$.

Then if BAD be a parabola;

$$y = 2\sqrt{mx}, \quad \text{and } u = 4(b - x)\sqrt{mx}.$$

(24) Let BAD be a segment of a circle;

$$AM = x, \quad \text{radius} = a;$$

$\therefore\ PM = \sqrt{2ax - x^2}$, and $u = 2(b - x).\sqrt{2ax - x^2}$.

(25) If A be the vertex, S the focus and P a point in a parabola, find the value of the ratio of $AP : SP$, when it is greatest. Ratio $= \frac{2}{\sqrt{3}}$.

(26) Cut the greatest parabola from a given cone.

(27) Required the least triangle TCt which can be described about a given quadrant.



$CA = a$, $CM = x$, $CN = y$;

$u = \frac{1}{2} CT . Ct$; $CT = \frac{a^2}{x}$; $Ct = \frac{a^2}{y}$;

and if u = maximum, $x = y$ and $\angle ACP = 45^0$.

(28) Let APB be a parabolic arc and C the focus.

$AN = x$, $AC = a$, $u = \frac{(x + a)^2\sqrt{a}}{2\sqrt{x}}$, whence $x = \frac{a}{3}$.

(29) Inscribe the greatest ellipse in a given isosceles triangle.



Let $Da = 2x$, $cb = y$; $\therefore\ u = \pi . yx$.

$AD = a$; $DB = b$.

Now $cN = \frac{ca^2}{cA} = \frac{x^2}{a - x}$;

$\therefore\ aN = \frac{ax - 2x^2}{a - x}$, $DN = \frac{a'x}{a - x}$.

But $\frac{BD^2}{AD^2} . AN^2 = PN^2 = \frac{y^2}{x^2}(Na . ND)$;

$\therefore\ b^2\left(\frac{a - 2x}{a - x}\right)^2 = y^2 . \frac{(a - 2x)\,a}{(a - x)^2}$; $\therefore\ y^2 = \frac{b^2}{a}(a - 2x)$;

$\therefore\ u = \pi yx = \frac{\pi b}{\sqrt{a}} . x\sqrt{a - 2x}$; $\therefore\ x = \frac{a}{3}$.

(30) Inscribe the greatest parabola in a given isosceles triangle. Axis $\frac{3}{4}$th of altitude of triangle.

(31) Within a given parabola inscribe the greatest parabola, the vertex of the latter being at the bisection of the base of the former. Axis $= \frac{2}{3}$ of given axis.

(32) The corner of a leaf is turned back, so as just to reach the other edge of the page: find when the length of the crease is a minimum.



$AP = x$, $AB = a$; $\therefore Aa = \sqrt{2ax}$,

also $Aa \,.\, PQ = 2AQ \,.\, AP$,

since $AQaP$ may be inscribed in a circle;

$$\therefore u^2 = PQ^2 = \frac{2x^3}{2x - a}; \quad \therefore x = \frac{3a}{4}.$$

(33) Find when the part turned down is a minimum. $x = \frac{2}{3}a$.

(34) Inscribe the greatest cylinder within a cone.

a, altitude of cone; x, of cylinder; $x = \frac{a}{3}$; $u = \frac{4\pi b^2 a}{27}$.

(35) Inscribe the greatest cone within a sphere.

$$x = \text{altitude of cone} = \frac{4a}{3}; \quad u = \frac{32}{81} \,.\, \pi a^3.$$

(36) Given the surface of a cylinder, find its form that its volume may be a maximum. Altitude = diameter of base.

(37) Given the volume, find when the surface is least. Altitude = diameter of base.

(38) In the trapezium $ABCD$, the base $AB = a$, $AD = BC = b$, find CD, CD being parallel to AB, that the area may be a maximum; $2CD = \sqrt{8b^2 + a^2} + a$.

(39) PQ is a chord in a semicircle parallel to the diameter AB, join AQ, BP cutting in O: find AP that the triangle POQ may be the greatest possible. $AP = 38^\circ \,.\, 40'$.

(40) Through a given point P between two given straight lines AB, AC (page 94): to draw PQ, so that $AP + AQ$ may be a minimum.

$$u = PQ = (\sqrt{a} + \sqrt{b})^2.$$

(41) Next draw PQ so that the triangle PAQ may have the least area.

$$x = 2a; \quad y = 2b; \quad u = 2ab.$$

(42) ACB is a quadrant, C the centre, CB the horizontal radius is bisected in D, a point P is taken in the arc, and CP, PD are joined, shew that the angle CPD is greatest when PD is vertical.

(43) The centres of two spheres (radii $r_1 r_2$) are at the extremities of a line $2a$, on which a circle is described. Find a point in the circumference, from which the greatest portion of spherical surface is visible.

If x and y be the distances of the point from the centres of the two spheres,

$$u = 2\pi r_1 \left(r_1 - \frac{r_1^2}{x}\right) + 2\pi r_2 \left(r_2 - \frac{r_1^2}{y}\right),$$

$$\text{whence } x = \frac{2ar_1}{\sqrt{r_1^2 + r_2^2}}.$$

(44) Find the position of a line passing through one of three given points, so that the rectangle of the perpendiculars from the other two points may be a maximum or minimum.

There are two lines respectively perpendicular, fulfilling the conditions.

(45) In a spherical triangle, find δ when $u = \theta - \phi$ is a minimum, and

$$\cos\theta = -\frac{\sin a + \sin\delta \,.\, \sin l}{\cos\delta \,.\, \cos l},$$

$$\cos\phi = -\frac{\sin\delta \,.\, \sin l}{\cos\delta \,.\, \cos l},$$

differentiating the three equations and eliminating $d\theta$ and $d\phi$,

$$\frac{\sin\theta}{\sin\phi} = \frac{\sin a}{\sin l}\sin\delta + 1;$$

$$\text{whence } \sin\delta = -\sin l \,.\, \tan\left(\tfrac{1}{2}a\right).$$

This is the problem of the shortest twilight, and if Z be the zenith, P the north pole, S the sun when twilight begins, s when it ends, $\theta = ZPS$, $\phi = ZPs$; $\therefore\ \theta - \phi$ converted into time is the duration, δ is the sun's declination, l the latitude of the place, a is the depression of S below the horizon, generally taken $= 18^0$; $\therefore\ \sin\delta = -\sin l \tan 9^0$; the negative sign shews the sun is on the south side of the equator.

THE CELLS OF BEES.

99. The first examination of the comb of the bee-hive presents a collection of waxen cells, the upper surface being parallel to the lower: but these parallel surfaces being sepa-

rated, each cell is found to be of a prismatic form; its base being a regular hexagon, and the other end of the prism formed of three equal rhombuses, composing the solid angle S. And it is remarkable that the two collections of cells, which by their junction form the comb, are so united that the axis of any one cell of one collection is in the continuation of the line of junction of three other cells of the other collection.

That the bases should be regular hexagons is an instance of the economy of nature; for only three figures, triangles, squares, and hexagons can completely occupy space; and of these, including the same area, the hexagon has the least perimeter.

PROP. Join A and C the extremities of $A'A$, $C'C$ of two equal edges of a prism, and let a plane through CA parallel to the base meet the axis in P; and let a plane inclined to the base, also through CA cut the axis in S and $B'B$ in b. Then



$$SP = Bb, \ \triangle AOP = \triangle AOB;$$

$\therefore$ pyramid $ACSP$ = pyramid $ACBb$. Hence whatever may be the inclination SOP of the plane SCb to the base of the cell, the solid content of the cell remains unaltered, but the surface varies with $\angle SOP$.

To find $\angle SOP$ when the surface of the cell is a minimum.

Let $\angle SOP = \theta$, $A'B' = AB = a$, $AA' = b$.

Then $BO = \frac{a}{2}$; $Ob = OS = \frac{BO}{\cos\theta} = \frac{a}{2\cos\theta}$.

$$Bb = SP = OB \tan\theta = \frac{a}{2}\tan\theta;$$

$$\therefore \text{ trapezium } AA'B'b = BA' - \triangle ABb = ba - \frac{a^2}{4}\tan\theta;$$

$$\therefore \text{ lateral surface of cell} = 3a\left(2b - \frac{a}{2}\tan\theta\right).$$

And rhombus $= SO \times AC = SO \times A'C' = \frac{a}{2\cos\theta} \cdot a\sqrt{3}$;

Note. BP in the figure, ought to pass through O.

$$\therefore \text{ surface of the tetrahedron} = \frac{3a^2\sqrt{3}}{2\cos\theta};$$

$$\therefore u = 3a\left\{2b - \frac{a}{2}\tan\theta + \frac{a\sqrt{3}}{2\cos\theta}\right\};$$

$$\therefore \frac{du}{d\theta} = 3a\left\{-\frac{a}{2}\sec^2\theta + \frac{a}{2}\sqrt{3}\,.\sec\theta\,.\tan\theta\right\} = 0;$$

$$\therefore \sec\theta = \sqrt{3}\tan\theta; \ \therefore \sin\theta = \frac{1}{\sqrt{3}}; \ \therefore \theta = 35^0.15'.51''.$$

$$\text{Also } Ob = \frac{a}{2}\sqrt{\frac{3}{2}}; \quad Cb = \sqrt{(CO)^2 + (Ob)^2} = \frac{3a}{2\sqrt{2}},$$

$$\tan OCb = \frac{Ob}{CO} = \frac{1}{\sqrt{2}}; \ \therefore \sin OCb = \frac{1}{\sqrt{3}};$$

$$\therefore \angle OCb = \angle SOP,$$

$$\text{and } \angle SCb = 2\angle SOP.$$

Hence the acute angle of the rhomb is double the inclination of the rhomb to the base of the cell. These results agree with the most exact measurements made in a multitude of cells*.

100. If $x = a$, make $\frac{du}{dx} = \infty$, the preceding rules are inapplicable, since they are founded on the supposition that $f(a+h)$ is expanded according to the ascending integral powers of h by means of Taylor's Theorem; but which is not the case, when the differential coefficients become infinite.

Let then $f(a+h)$ be expanded by the ordinary methods, and assume

$$f(a+h) = f(a) + Ph^\alpha + Qh^\beta + Rh^\gamma + \&c.$$

where α is the least of all the indices of h;

$$\therefore f(a+h) - f(a) = Ph^\alpha + Qh^\beta + Rh^\gamma + \&c.\ldots.(1),$$

$$\text{and } f(a-h) - f(a) = P(-h)^\alpha + Q(-h)^\beta + \&c.$$

by writing $-h$ for h in series (1).

Now if h be made very small, the algebraical sign of the developements will depend on that of their first term. If therefore we have a maximum or minimum, since

* Puissant's Geometry: Cresswell's Maxima and Minima.

$f(a+h)-f(a)$, and $f(a-h)-f(a)$ must have the same signs, Ph^a and $P(-h)^a$; and $\therefore h^a$ and $(-h)^a$ must have the same sign, or a must either be an even number or a fraction with an even number for its numerator.

(1) If a be an even number, it shews that at a maximum or minimum the first existent term of the developement must involve an even power of h, a conclusion we have already come to in the preceding pages.

(2) If a be a fraction, it must be of the form $\frac{2n}{2m+1}$.

Ex. Let $u = b + c(x-a)^{\frac{3}{5}}$.

Here $\frac{du}{dx} = \frac{2c}{3}\frac{1}{(x-a)^{\frac{1}{3}}}$, which is infinite, if $x = a$.

But $x = a$, gives $u = b$,

$x = a + h$ gives $u = b + ch^{\frac{3}{5}}$,

$x = a - h$ ..... $u = b + ch^{\frac{3}{5}}$,

and $f(a+h)$ and $f(a-h)$ are both $> f(a)$, if c be positive,

$< f(a)$, if c be negative.

If $\therefore$ c be positive $x = a$, makes $u = b$ a minimum,

c be negative $x = a$, ... $u = b$ a maximum.

CHAPTER VIII.

FUNCTIONS OF TWO OR MORE VARIABLES.—IMPLICIT FUNCTIONS.

---

101. As yet we have only treated of functions of a single variable; we next proceed to the case in which $u = f(xy)$, where x and y are independent of each other, and the value of u corresponding to new values $x + h$, and $y + k$, of x and y, is required.

Now when u is a function of x and y, u may vary on three suppositions; 1st, x may vary, and y remain constant; 2nd, y may vary, and x remain constant; and 3rd, x and y may both vary together.

Thus suppose $u = xy^2$, and let x become $x + h$, and y remain constant; therefore if u' be the value of u,

$$u' = (x + h)\, y^2 = xy^2 + y^2 h.$$

Next let y become $y + k$, and x be constant, and let u_1 be the value of u;

$$\therefore\ u_1 = x(y + k)^2 = xy^2 + 2xyk + xk^2.$$

Again, in the equation $u = xy^2$ write $x + h$ for x, and $y + k$ for y, and let u_2 be the value of u, or $u_2 = f(x + h, y + k)$;

$$\therefore\ u_2 = (x + h)(y + k)^2 = xy^2 + y^2 h + 2xyk + 2ykh + xk^2 + k^2 h,$$

the same result as would have been obtained had we put

$$y + k \text{ for } y \text{ in } u', \text{ or } x + h \text{ for } x \text{ in } u_1.$$

102. Next considering the question in a general point of view.

Let $u = f(x, y)$, then if y remain constant while x becomes $x + h$, we have, by Taylor's Theorem,

$$f(x + h,\ y) = u + \frac{du}{dx}h + \frac{d^2u}{dx^2}\frac{h^2}{1.2} + \frac{d^3u}{dx^3}\frac{h^3}{2.3} + \&c.;$$

or, if x remain constant while y becomes $y + k$,

$$f(x,\ y + k) = u + \frac{du}{dy}k + \frac{d^2u}{dy^2}\frac{k^2}{1.2} + \frac{d^3u}{dy^3}\cdot\frac{k^3}{2.3} + \&c.$$

Suppose now that x and y both vary; or x become $x + h$, and y become $y + k$; it is not possible to make both these assumptions at once: but if we use either of the two series,

for $f(x+h, y)$ or $f(x, y+k)$, and in the former put $y+k$ for y, or in the latter $x+h$ for x, we shall in either case have $f(x+h, y+k)$, and its true developement.

Assuming the first expansion,

$$f(x+h.y) = u + \frac{du}{dx}.h + \frac{d^2u}{dx^2}\frac{h^2}{1.2} + \frac{d^3u}{dx^3}\frac{h^3}{2.3} + \&c.$$

But $u = f(xy)$, and therefore $\frac{du}{dx}$, $\frac{d^2u}{dx^2}$, are also functions of x and y, if therefore y become $y+k$; u, $\frac{du}{dx}$, $\frac{d^2u}{dx^2}$, &c. will become functions of $y+k$, and may be expanded by Taylor's Theorem, x being considered constant.

Let therefore y become $y+k$;

$$\therefore u \text{ becomes } u + \frac{du}{dy}.k + \frac{d^2u}{dy^2}\frac{k^2}{1.2} + \frac{d^3u}{dy^3}.\frac{k^3}{2.3} + \&c.....(a),$$

and to obtain the values of $\frac{du}{dx}$, $\frac{d^2u}{dx^2}$, &c. we must write $\frac{du}{dx}$, $\frac{d^2u}{dx^2}$, &c. for u in the series (a);

$$\therefore \frac{du}{dx} \text{ becomes } \frac{du}{dx} + \frac{d.\left(\frac{du}{dx}\right)}{dy}k + \frac{d^2.\left(\frac{du}{dx}\right)}{dy^2}.\frac{k^2}{1.2} + \&c.$$

$$\frac{d^2u}{dx^2} \text{ ..... becomes } \frac{d^2u}{dx^2} + \frac{d.\left(\frac{d^2u}{dx^2}\right)}{dy}.k + \&c.$$

$$\frac{d^3u}{dx^3} \text{ ..... becomes } \frac{d^3u}{dx^3} + \frac{d.\left(\frac{d^3u}{dx^3}\right)}{dy}.k + \&c.$$

But it has been agreed to write $\frac{d^2u}{dy.dx}$ for $\frac{d.\left(\frac{du}{dx}\right)}{dy}$, which expresses that the function has been differentiated twice, 1st considering x, and then y as variable;

and $\frac{d.\left(\frac{d^2u}{dx^2}\right)}{dy}$ is written $\frac{d^3u}{dydx^2}$, and $\frac{d^n.\left(\frac{d^mu}{dx^m}\right)}{dy^n}$ is written

$\frac{d^{m+n} \cdot u}{dy^n \cdot dx^m}$, denoting the differential coefficient when the function has been differentiated m times with regard to x, and n times with regard to y.

Making these substitutions, and multiplying the expansion of $\frac{du}{dx}$ by h, that of $\frac{d^2u}{dx^2}$ by $\frac{h^2}{1.2}$, &c. we shall have

$$f(x+h,\ y+k) = u + \frac{du}{dy}k + \frac{d^2u}{dy^2}\frac{k^2}{1.2} + \frac{d^3u}{dy^3}\frac{k^3}{2.3} + \&c.$$
$$+ \frac{du}{dx}h + \frac{d^2u}{dy.dx}hk + \frac{d^3u}{dy^2.dx}.\frac{hk^2}{1.2} + \&c.$$
$$+ \frac{d^2u}{dx^2}\frac{h^2}{1.2} + \frac{d^3u}{dy.dx^2}.\frac{h^2k}{1.2} + \&c.$$
$$+ \frac{d^3u}{dx^3}\frac{h^3}{2.3} + \&c.$$
$$+ \&c..... + \&c.$$

103. But this developement was obtained, by first supposing x, and then y to vary; but manifestly we should have had an equal result, had y first become $y+k$, and then x become $x+h$. On this supposition we have

$$f(x,\ y+k) = u + \frac{du}{dy}k + \frac{d^2u}{dy^2}\frac{k^2}{1.2} + \frac{d^3u}{dy^3}\frac{k^3}{2.3} + \&c.;$$

put $x+h$ for x,

$$\therefore\ u \text{ becomes } u + \frac{du}{dx}h + \frac{d^2u}{dx^2}\frac{h^2}{1.2} + \frac{d^3u}{dx^3}\frac{h^3}{2.3} + \&c.,$$
$$\frac{du}{dy} \text{ ..... becomes } \frac{du}{dy} + \frac{d^2u}{dxdy}h + \frac{d^3u}{dx^2dy}.\frac{h^2}{1.2} + \&c.,$$
$$\frac{d^2u}{dy^2} \text{ ..... becomes } \frac{d^2u}{dy^2} + \frac{d^3u}{dxdy^2}\frac{h}{1} + \&c.,$$
$$\text{..... becomes ..... } + \&c.;$$

whence by substitution the total developement becomes

$$f(x+h,\ y+k) = u + \frac{du}{dx}h + \frac{d^2u}{dx^2}\frac{h^2}{1.2} + \frac{d^3u}{dx^3}\frac{h^3}{2.3} + \&c.$$
$$+ \frac{du}{dy}k + \frac{d^2u}{dxdy}kh + \frac{d^3u}{dx^2dy}\frac{h^2k}{1.2} + \&c.$$

$$+\frac{d^2u}{dy^2}\frac{k^2}{1.2}+\frac{d^3u}{dxdy^2}\frac{k^2h}{1.2}+\&c.$$

$$+\frac{d^3u}{dy^3}\frac{k^3}{2.3}+\&c.$$

$$+\&c.$$

Cor. 1. Since the series are equal, the coefficients of the same powers of h and k ought to be equal;

$$\therefore \frac{d^2u}{dydx}=\frac{d^2u}{dxdy},$$

$$\frac{d^3u}{dy^2dx}=\frac{d^3u}{dxdy^2},$$

$$\frac{d^3u}{dydx^2}=\frac{d^3u}{dx^2dy},$$

$$\&c.=\&c.$$

$$\text{and } \frac{d^{m+n}.u}{dy^m dx^n}=\frac{d^{m+n}u}{dx^n dy^m}.$$

Whence it appears that the order of differentiation is indifferent; or that the differential coefficient of u differentiated m times with respect to x, and n times with respect to y, is equal to the differential coefficient when u has been first differentiated n times with regard to y, and then m times with regard to x.

Cor. 2. Again, $\because \frac{d^2u}{dydx}=\frac{d^2u}{dxdy}$; $\therefore$ writing $\frac{du}{dx}$ for u,

$$\frac{d^2.\left(\frac{du}{dx}\right)}{dydx}=\frac{d^2.\left(\frac{du}{dx}\right)}{dxdy},$$

$$\text{or } \frac{d^3u}{dydx^2}=\frac{d^3u}{dxdydx},$$

$$\text{and } \frac{d^2.\left(\frac{du}{dy}\right)}{dydx}=\frac{d^2.\left(\frac{du}{dy}\right)}{dxdy};$$

$$\therefore \frac{d^3.u}{dydxdy}=\frac{d^3.u}{dxdy^2}.$$

104. Since $\frac{du}{dx}$, $\frac{d^2u}{dx^2}$, $\frac{d^3u}{dx^3}$, &c. have been obtained by the consideration of x alone being the independent variable,

such differential coefficients have been called *partial* differential coefficients, and for the same reason $\frac{du}{dy}$, $\frac{d^2u}{dy^2}$, &c. are also called partial differential coefficients, and these partial differential coefficients are frequently included within brackets, thus $\left(\frac{du}{dx}\right)$ is the partial differential coefficient with respect to x, and $\left(\frac{du}{dy}\right)$ is the partial differential coefficient with respect to y, and $\left(\frac{du}{dx}\right)dx$, and $\left(\frac{du}{dy}\right)dy$, are the partial differentials of u, with regard to x and y respectively.

105. The term $\frac{du}{dx}h + \frac{du}{dy}k$, which involves only the first powers of h and k is called the *total* differential of u, and putting dx for h, and dy for k, is thus written;

$$du = \left(\frac{du}{dx}\right).dx + \left(\frac{du}{dy}\right)dy,$$

or the total differential of $u = f(xy)$ is the sum of the partial differentials.

106. From the first differential of u, we may form by differentiation the successive differentials d^2u, d^3u; &c.

$$\text{For } du = \left(\frac{du}{dx}\right)dx + \left(\frac{du}{dy}\right)dy \ldots\ldots\ldots(\beta).$$

And differentiating, considering $\left(\frac{du}{dx}\right)$ and $\left(\frac{du}{dy}\right)$ as functions of x and y, and dx and dy constant, we have, by writing successively, $\left(\frac{du}{dx}\right)$ and $\left(\frac{du}{dy}\right)$ for u in (β),

$$d.\left(\frac{du}{dx}\right) = \left(\frac{d^2u}{dx^2}\right).dx + \frac{d^2u}{dydx}.dy,$$

$$d.\left(\frac{du}{dy}\right) = \frac{d^2u}{dxdy}.dx + \left(\frac{d^2u}{dy^2}\right)dy.$$

Then substituting these values, since

$$d^2u = d.\left(\frac{du}{dx}\right).dx + d\left(\frac{du}{dy}\right).dy,$$

$$d^2u = \frac{d^2u}{dx^2}.dx^2 + 2.\frac{d^2u}{dx.dy}.dy.dx + \frac{d^2u}{dy^2}.dy^2.$$

Again, to find d^3u, substituting as before

$$d.\left(\frac{d^2u}{dx^2}\right) = \left(\frac{d^3u}{dx^3}\right).dx + \frac{d^3u}{dydx^2}.dy,$$

$$d.\left(\frac{d^2u}{dxdy}\right) = \frac{d^3u}{dx^2dy}.dx + \frac{d^3u}{dydydx}.dy,$$

$$d.\left(\frac{d^2u}{dy^2}\right) = \frac{d^3u}{dxdy^2}.dx + \left(\frac{d^3u}{dy^3}\right)dy;$$

$$\therefore\ d^3u = \left(\frac{d^3u}{dx^3}\right)dx^3 + 3.\frac{d^3u}{dx^2dy}dx^2dy$$

$$+ 3.\frac{d^3u}{dy^2dx}.dy^2.dx + \left(\frac{d^3u}{dy^3}\right)dy^3.$$

107. The law of continuity is almost obvious; for the numerical coefficients appear to be those of the terms of the expansion of the binomial $(h+k)^n$: but to prove that they do follow this law, assume that

$$d^nu = \frac{d^nu}{dx^n}dx^n + n.\frac{d^nu}{dx^{n-1}dy}.dx^{n-1}dy$$

$$+ n.\frac{n-1}{2}\frac{d^nu}{dx^{n-2}dy^2}dx^{n-2}dy^2 + \&c.$$

Then, differentiating the successive terms by means of

$$du = \frac{du}{dx}dx + \frac{du}{dy}.dy,$$

$$d.\left(\frac{d^nu}{dx^n}\right) = \frac{d^{n+1}u}{dx^{n+1}}.dx + \frac{d^{n+1}u}{dx^ndy}.dy.....(1),$$

$$d.\left(\frac{d^nu}{dx^{n-1}dy}\right) = \frac{d^{n+1}u}{dx^ndy}.dx + \frac{d^{n+1}u}{dx^{n-1}dy^2}.dy.....(2),$$

$$d.\left(\frac{d^nu}{dx^{n-2}dy^2}\right) = \frac{d^{n+1}u}{dx^{n-1}dy^2}.dx + \frac{d^{n+1}u}{dx^{n-2}dy^3}.dy.....(3).$$

$$\&c. = \&c.$$

Then, multiply (1) by dx^n, (2) by $n.dx^{n-1}dy$, (3) by $n.\left(\frac{n-1}{2}\right).dx^{n-2}dy^2$, and adding

$$d^{n+1}u = \frac{d^{n+1}u}{dx^{n+1}}.dx^{n+1} + (n+1).\left(\frac{d^{n+1}u}{dx^ndy}\right).dx^ndy$$

$$+ \frac{(n+1).n}{2}.\frac{d^{n+1}u}{dx^{n-1}dy^2}.dx^{n-1}dy^2 + \&c.$$

or if the formula be true for the index n, it is true for $n+1$, and we have seen that it is true when $n=3$; it is therefore always true.

Cor. If instead of dx and dy we write h and k, we have

$$du = \frac{du}{dx}h + \frac{du}{dy}k,$$

$$d^2u = \frac{d^2u}{dx^2}h^2 + 2\frac{d^2u}{dydx}hk + \frac{d^2u}{dy^2}k^2,$$

$$d^3u = \frac{d^3u}{dx^3}h^3 + 3\frac{d^3u}{dx^2dy}h^2k + 3\frac{d^3u}{dy^2dx}hk^2 + \frac{d^3u}{dy^3}k^3,$$

&c.

and therefore $u_2 = f\{(x+h),\ (y+k)\}$

$$= u + du + \frac{d^2u}{1.2} + \frac{d^3u}{2.3} + \&c.$$

$$+ \frac{d^nu}{1.2.3...n} + \&c.,$$

or the expansion of $f(x+h, y+k)$ may be found from the successive differentiation of $u = f(x, y)$.

108. Again, if $u = f(x, y, z)$, and if $x+h$, $y+k$, $z+m$, be new values of x, y, z, and u_2 the value of u,

$$u_2 = u + \frac{du}{dx}h + \frac{du}{dy}k + \frac{du}{dz}m$$
$$+ Ah^2 + Bk^2 + Cm^2 + \&c.$$

For, supposing z to be constant while x and y become $x+h$, and $y+k$ respectively;

$$\therefore f(x+h, y+k, z) = u + \frac{du}{dx}h + \frac{du}{dy}k + \&c.$$

Now, let z become $z+m$;

$$\therefore u \text{ becomes } u + \frac{du}{dz}m + \&c.$$

$$\frac{du}{dx} \text{ ..... } \frac{du}{dx} + \frac{d^2u}{dzdx}m + \&c.$$

$$\frac{du}{dy} \text{ ..... } \frac{du}{dy} + \frac{d^2u}{dzdy}m + \&c.$$

$$\&c. \quad = \quad \&c.$$

$$u_2 = f(x+h,\ y+k,\ z+m) = u + \frac{du}{dz}m + \frac{du}{dx}h + \frac{du}{dy}k + \&c.$$

and in a similar manner may the expansion of a function of four or more variables be effected.

Cor. Hence, if for m, h, and k, we put dz, dx, and dy,

$$du = d.f(x, y, z) = \left(\frac{du}{dx}\right)dx + \left(\frac{du}{dy}\right)dy + \left(\frac{du}{dz}\right)dz.$$

This result may however be obtained in the following manner:

109. Let $u = f(x, y, z)$; find du.

Let $n = \phi(y, z)$, so that we may put $u = f(x, n)$;

$$\therefore\ du = \left(\frac{du}{dx}\right)dx + \left(\frac{du}{dn}\right)dn.$$

But $\because\ n = \phi(y, z)$; $\quad \therefore\ dn = \dfrac{dn}{dy}dy + \dfrac{dn}{dz}.dz$;

$$\therefore\ du = \frac{du}{dx}dx + \frac{du}{dn}.\frac{dn}{dy}dy + \frac{du}{dn}.\frac{dn}{dz}dz.$$

But $\dfrac{du}{dn}.\dfrac{dn}{dy} = \dfrac{du}{dy}$, and $\dfrac{du}{dn}.\dfrac{dn}{dz} = \dfrac{du}{dz}$;

$$\therefore\ du = \frac{du}{dx}.dx + \frac{du}{dy}.dy + \frac{du}{dz}.dz;$$

and the same method may be extended to any number of variables; whence it appears that the differential of a function of any number of variables equals the sum of the partial differentials.

110. From the preceding reasoning we may find the differentials of a function of two functions of the variable x.

For suppose $u = f(y, z)$, where y and z are functions of x, and therefore u is also a function of x; to find the total differential of u.

Let k and m be the increments of y and z, if x become $x + h$;

$$\therefore\ u_1 = u + \left(\frac{du}{dy}\right)k + \left(\frac{du}{dz}\right)m + \&c.$$

But $k = \dfrac{dy}{dx}h + \&c.$; $\quad m = \dfrac{dz}{dx}h + \&c.$;

$$u_1 = u + \left\{\left(\frac{du}{dy}\right)\frac{dy}{dx} + \left(\frac{du}{dz}\right).\frac{dz}{dx}\right\}h + \&c.;$$

$$\therefore\ du = \left\{\left(\frac{du}{dy}\right)\frac{dy}{dx} + \left(\frac{du}{dz}\right)\cdot\frac{dz}{dx}\right\}h;$$

or since $\dfrac{dy}{dx}h = dy$, and $\dfrac{dz}{dx}h = dz$,

$$du = \left(\frac{du}{dy}\right)\cdot dy + \left(\frac{du}{dz}\right)dz,$$

which is the total differential of a function of two functions.

111. Again, if $u = f(y, z, v)$, and y, z, v are functions of x; to find the same: let n be the increment of v;

$$\therefore\ u_1 = u + \left(\frac{du}{dy}\right)k + \left(\frac{du}{dz}\right)m + \left(\frac{du}{dv}\right)n + \&c.$$

$$= u + \left\{\left(\frac{du}{dy}\right)\frac{dy}{dx} + \left(\frac{du}{dz}\right)\frac{dz}{dx} + \left(\frac{du}{dv}\right)\cdot\frac{dv}{dx}\right\}h + \&c.;$$

$$\therefore\ du = \left\{\left(\frac{du}{dy}\right)\frac{dy}{dx} + \left(\frac{du}{dz}\right)\frac{dz}{dx} + \left(\frac{du}{dv}\right)\cdot\frac{dv}{dx}\right\}h$$

$$= \left(\frac{du}{dy}\right)dy + \left(\frac{du}{dz}\right)dz + \left(\frac{du}{dv}\right)dv.$$

Similarly may the total differential of a function of n functions be found.

EXAMPLES.

(1) Let $u = x^m y^n$; find du.

$$\left(\frac{du}{dx}\right) = my^n x^{m-1};\quad \left(\frac{du}{dy}\right) = nx^m y^{n-1};$$

$$\therefore\ du = \left(\frac{du}{dx}\right)dx + \left(\frac{du}{dy}\right)dy$$

$$= my^n x^{m-1}dx + nx^m y^{n-1}dy$$

$$= x^{m-1}y^{n-1}(mydx + nydy),$$

$$\frac{d^2u}{dydx} = mny^{n-1}x^{m-1} = \frac{d^2u}{dxdy}.$$

(2) $\quad u = \dfrac{x^m}{y^n};\quad du = \dfrac{x^{m-1}(mydx - nxdy)}{y^{n+1}}.$

(3) $u = \sin(x^2 y)$; find $\dfrac{d^2u}{dxdy}$.

$$\frac{d^2u}{dxdy} = 2x(\cos x^2 y - x^2 y \sin x^2 y).$$

(4) $u = \sin^{-1}\dfrac{x}{y}$; $du = \dfrac{ydx - xdy}{y\sqrt{y^2 - x^2}}$.

(5) $u = \log\left(\dfrac{x + \sqrt{x^2 - y^2}}{x - \sqrt{x^2 - y^2}}\right)$; $du = 2 \cdot \dfrac{ydx - xdy}{y\sqrt{x^2 - y^2}}$.

(6) $u = x^y$; $du = x^y\left(\dfrac{y}{x} + \log x\right)$.

$$\frac{d^2u}{dydx} = x^y\left(\frac{1}{x} + \frac{y}{x}\log x\right) = \frac{d^2u}{dxdy}.$$

(7) $u = \dfrac{x^2 y}{a^2 - z^2}$; find du, and shew that,

$$\frac{d^2u}{dxdy} = \frac{2x}{a^2 - z^2} = \frac{d^2u}{dydx},$$

$$\frac{d^2u}{dxdz} = \frac{4xyz}{(a^2 - z^2)^2} = \frac{d^2u}{dzdx},$$

$$\frac{d^2u}{dydz} = \frac{2x^2 z}{(a^2 - z^2)^2} = \frac{d^2u}{dzdx},$$

$$\frac{d^3u}{dxdydz} = \frac{4xz}{(a^2 - z^2)^2} = \frac{d^3u}{dzdydx} = \frac{d^3u}{dydxdz}.$$

(8) Let $u = \dfrac{1}{\sqrt{(a-x)^2 + (b-y^2) + (c-z)^2}}$, shew that

$$\frac{d^2u}{dx^2} + \frac{d^2u}{dy^2} + \frac{d^2u}{dz^2} = 0.$$

Here $\dfrac{1}{u^2} = (a-x)^2 + (b-y)^2 + (c-z)^2$;.....(1);

$$\therefore \frac{du}{dx} = u^3(a-x);$$

$$\therefore \frac{d^2u}{dx^2} = 3u^2\frac{du}{dx}(a-x) - u^3 = 3u^5(a-x)^2 - u^3,$$

and from the symmetric form of (1),

$$\frac{d^2u}{dy^2} = 3u^5(b-y)^2 - u^3,$$

$$\frac{d^2u}{dz^2} = 3u^5(c-z)^2 - u^3;$$

$$\therefore \frac{d^2u}{dx^2} + \frac{d^2u}{dy^2} + \frac{d^2u}{dz^2} = 3u^5\frac{1}{u^2} - 3u^3 = 0,$$

an equation of great importance in physical science.

112. The following theorem is important in the integration of homogenous differential equations.

Let u be a homogenous function of x, y, z, &c. and let n be the sum of the exponents in each term, then

$$nu = \frac{du}{dx}x + \frac{du}{dy}y + \frac{du}{dz}z + \&c.$$

For x, y, z, &c. put $x + mx$, $y + my$, $z + mz$, &c. then u becomes $(1+m)^n u$;

$$\begin{aligned}\therefore (1+m)^n u = u &+ \frac{du}{dx}mx + \frac{d^2u}{dx^2}\cdot\frac{m^2x^2}{1.2} + \&c.\\ &+ \frac{du}{dy}my + \frac{d^2u}{dydx}m^2xy + \&c.\\ &+ \frac{du}{dz}mz + \frac{d^2u}{dzdx}\cdot m^2zx + \&c.\\ &+ \frac{d^2u}{dy^2}\cdot\frac{m^2y^2}{1.2} + \&c.\\ &+ \&c.,\end{aligned}$$

$$\text{also } = u + num + n\frac{n-1}{2}um^2 + \&c.;$$

whence by equating the same powers of m,

$$nu = \frac{du}{dx}x + \frac{du}{dy}y + \frac{du}{dz}z + \&c.,$$

$$\begin{aligned}\text{and } n(n-1)u = &\frac{d^2u}{dx^2}x^2 + \frac{d^2u}{dy^2}y^2 + \frac{d^2u}{dz^2}z^2 + \&c.\\ &+ 2\frac{d^2u}{dydx}xy + 2\frac{d^2u}{dzdx}zx + 2\frac{d^2u}{dzdy}zy + \&c.\end{aligned}$$

(1) Let $u = (x + y + z)^3$; here $n = 3$;

$$\therefore \frac{du}{dx} = \frac{du}{dy} = \frac{du}{dz} = 3(x + y + z)^2;$$

$$\therefore \frac{du}{dx}x + \frac{du}{dy}y + \frac{du}{dz}z = 3(x + y + z)^3 (x + y + z) = 3u.$$

(2) Let $u = \dfrac{xyz}{x + y + z}$; here $n = 2$,

$$\text{and } \frac{du}{dx}x + \frac{du}{dy}y + \frac{du}{dz}z = 2u.$$

(3) Let $u = \dfrac{x^3 + y^3}{x - y}$; here $n = 2$,

$$\text{and } \frac{du}{dx}x + \frac{du}{dy}y = 2u.$$

(4) Let $u = \dfrac{1}{x^2 + y^2}$; here $n = -2$,

$$\text{and } \frac{du}{dx}x + \frac{du}{dy}y = -2u.$$

(5) Let $u = \dfrac{\sqrt{x} + \sqrt{y}}{x + y}$; here $n = -\frac{1}{2}$,

$$\text{and } \frac{du}{dx}x + \frac{du}{dy}y = -\tfrac{1}{2}u.$$

(6) Let $u = \dfrac{xy}{ax + bz}$; here $n = 1$,

$$\text{and } \frac{du}{dx}x + \frac{du}{dy}y + \frac{du}{dz}z = u.$$

(7) Let $u = \sin^{-1}\sqrt{\dfrac{x - y}{x + y}}$; here $n = 0$,

$$\text{and } \frac{du}{dx}x + \frac{du}{dy}y = 0.$$

(8) Let $u = (x^2 + y^2)^2$; here $n = 4$;

$$\therefore \frac{d^2u}{dx^2}x^2 + 2\frac{d^2u}{dy\,dx}xy + \frac{d^2u}{dy^2}y^2 = 12u.$$

IMPLICIT FUNCTIONS.

113. When there is an implicit function of y and x, it is frequently impossible to solve the equation with respect to y, and obtain $y = f(x)$; but by considering $f(x, y,) = 0$ to be a function of two variables, we may from the preceding expansions for such functions obtain rules easy of application.

Let $u = f(x, y) = 0$, and let u_1 represent u when x becomes $x + h$, and therefore y becomes $y + k$;

$$\therefore\ u_1 = u + \left(\frac{du}{dx}\right) h + \left(\frac{du}{dy}\right) k + \&c.$$

But $\because\ u = 0$, whatever x and y are; $\therefore\ u_1 = 0$;

$$\therefore\ 0 = \left(\frac{du}{dx}\right) h + \left(\frac{du}{dy}\right) k + \&c.$$

But $k = \dfrac{dy}{dx} h + \&c.$, $\because\ y = f(x)$;

therefore, substituting for k,

$$0 = \left\{\left(\frac{du}{dx}\right) + \left(\frac{du}{dy}\right)\frac{dy}{dx}\right\} h + Bh^2 + Ch^3 + \&c.;$$

$$\therefore\ \left(\frac{du}{dx}\right) + \left(\frac{du}{dy}\right)\frac{dy}{dx} = 0,$$

a theorem by which $\dfrac{dy}{dx}$ may be found from the partial differentials $\left(\dfrac{du}{dx}\right)$ and $\left(\dfrac{du}{dy}\right)$.

Ex. $y^3 - 3axy + x^3 = 0$; find $\dfrac{dy}{dx}$.

Let $u = y^3 - 3axy + x^3$;

$$\therefore\ \left(\frac{du}{dx}\right) = -3ay + 3x^2;\ \left(\frac{du}{dy}\right) = 3y^2 - 3ax.$$

But $\left(\dfrac{du}{dx}\right) + \left(\dfrac{du}{dy}\right).\dfrac{dy}{dx} = 0$;

$$\therefore\ -3ay + 3x^2 + (3y^2 - 3ax)\frac{dy}{dx} = 0;\ \therefore\ \frac{dy}{dx} = \frac{ay - x^2}{y^2 - ax}.$$

Cor. 1. Since $\left\{\left(\dfrac{du}{dx}\right) + \left(\dfrac{du}{dy}\right).\dfrac{dy}{dx}\right\} h = 0$; $\therefore\ du = 0$;

$$\therefore \left(\frac{du}{dx}\right) dx + \left(\frac{du}{dy}\right) \cdot \frac{dy}{dx} dx = 0; \text{ putting } dx \text{ for } h;$$

$$\text{or } \left(\frac{du}{dx}\right) dx + \left(\frac{du}{dy}\right) dy = 0. \text{ Since } y = f(x).$$

Cor. 2. Hence, since if $u = 0$, $du = 0$; $\therefore$ if $du = 0$, $d^2u = 0$; and thus if $u = 0$, $d^nu = 0$.

114. From the equation $\left(\frac{du}{dx}\right) + \left(\frac{du}{dy}\right) \cdot \frac{dy}{dx} = 0$ (1), to find $\frac{d^2u}{dx^2}$ (where d^2u means the second total differential of u), and thence to deduce $\frac{d^2y}{dx^2}$.

$$\text{Since } \frac{du}{dx} = \left(\frac{du}{dx}\right) + \left(\frac{du}{dy}\right) \cdot \frac{dy}{dx} = 0;$$

$$\therefore \frac{d^2u}{dx^2} = \frac{d}{dx} \cdot \left(\frac{du}{dx}\right) + \frac{d}{dx}\left\{\left(\frac{du}{dy}\right) \cdot \frac{dy}{dx}\right\} \ldots\ldots\ldots\ldots (2).$$

Put in (1) $\frac{du}{dx}$ for u, and then $\frac{du}{dy}$ for u;

$$\therefore \frac{d}{dx} \cdot \left(\frac{du}{dx}\right) = \left(\frac{d^2u}{dx^2}\right) + \frac{d^2u}{dxdy} \cdot \frac{dy}{dx},$$

$$\frac{d}{dx}\left\{\left(\frac{du}{dy}\right) \cdot \frac{dy}{dx}\right\} = \left(\frac{du}{dy}\right) \cdot \frac{d^2y}{dx^2} + \frac{dy}{dx} \cdot \frac{d}{dx} \cdot \left(\frac{du}{dy}\right)$$

$$= \left(\frac{du}{dy}\right) \cdot \frac{d^2y}{dx^2} + \frac{dy}{dx}\left\{\frac{d^2u}{dxdy} + \frac{d^2u}{dy^2} \cdot \frac{dy}{dx}\right\};$$

$\therefore$ substituting in (2), we have

$$\frac{d^2u}{dx^2} = \left(\frac{d^2u}{dx^2}\right) + 2\frac{d^2u}{dxdy} \cdot \frac{dy}{dx} + \left(\frac{du}{dy}\right) \cdot \frac{d^2y}{dx^2} + \left(\frac{d^2u}{dy^2}\right) \cdot \frac{dy^2}{dx^2} = 0 \ldots (3),$$

and because from equation (1) $\frac{dy}{dx}$ may be found in terms of the partial differential coefficients $\left(\frac{du}{dx}\right)$, $\left(\frac{du}{dy}\right)$, and $\left(\frac{d^2u}{dx^2}\right)$ and $\left(\frac{d^2u}{dy^2}\right)$ and $\frac{d^2u}{dxdy}$ being similarly found, $\frac{d^2y}{dx^2}$ may be determined. In the same manner $\frac{d^3u}{dx^3}$, and $\therefore$ $\frac{d^3y}{dx^3}$ and diffe-

rential coefficients of higher orders may be found; but the investigations lead to equations of great complexity, and which in practice are seldom needed.

115. Next, let $u = 0$ be a function of three variables x, y, z, or let z be an implicit function of (x, y); and let $z + m$ be the value of z when the independent variables, x and y, become respectively $x + h$ and $y + k$;

$$\therefore \text{ since } u_1 = 0 = f(x + h,\ y + k,\ z + m),$$

$$0 = \left(\frac{du}{dx}\right) h + \left(\frac{du}{dy}\right) k + \left(\frac{du}{dz}\right) m + Ah^2 + Bk^2 + Cm^2 + \&c.$$

But $z + m = \phi(x + h,\ y + k)$;

$$\therefore\ m = \left(\frac{dz}{dx}\right) h + \left(\frac{dz}{dy}\right) k + \&c.$$

therefore, substituting for m, we have

$$0 = \left\{\left(\frac{du}{dx}\right) + \left(\frac{du}{dz}\right) \cdot \frac{dz}{dx}\right\} h + \left\{\left(\frac{du}{dy}\right) + \left(\frac{du}{dz}\right) \cdot \frac{dz}{dy}\right\} k + \&c.$$

whence, since h and k are independent, we have

$$\left(\frac{du}{dx}\right) + \left(\frac{du}{dz}\right) \cdot \frac{dz}{dx} = 0 \ldots\ldots\ldots\ldots (1),$$

$$\left(\frac{du}{dy}\right) + \left(\frac{du}{dz}\right) \cdot \frac{dz}{dy} = 0 \ldots\ldots\ldots\ldots (2);$$

whence $dz = \frac{dz}{dx} dx + \frac{dz}{dy} dy$ may be found.

116. The differential coefficients of the superior orders, can be found by differentiating the equations

$$\left(\frac{du}{dx}\right) + \left(\frac{du}{dz}\right) \cdot \frac{dz}{dx} = 0 \ldots\ldots\ldots\ldots (1),$$

$$\left(\frac{du}{dy}\right) + \left(\frac{du}{dz}\right) \cdot \frac{dz}{dy} = 0 \ldots\ldots\ldots\ldots (2).$$

117. Thus to obtain $\frac{d^2z}{dx^2}$, $\frac{d^2z}{dy\,dx}$ and $\frac{d^2z}{dy^2}$.

We must consider equations (1) and (2) as functions of x, y, z, and that $\left(\frac{du}{dx}\right)$, $\left(\frac{du}{dy}\right)$ and $\left(\frac{du}{dz}\right)$ are also functions of the same variables.

(1°) Let equation (1) be differentiated with respect to x; it must be considered as a function of x and z, and therefore from (Art. 114), putting z for y,

$$\left(\frac{d^2u}{dx^2}\right) + 2 \cdot \frac{d^2u}{dz \cdot dx} \cdot \frac{dz}{dx} + \left(\frac{d^2u}{dz^2}\right) \cdot \frac{dz^2}{dx^2} + \left(\frac{du}{dz}\right) \cdot \frac{d^2z}{dx^2} = 0 \ldots (3).$$

(2°) Differentiate (2), considered a function of y and z.

Write y for x in equation (3);

$$\left(\frac{d^2u}{dy^2}\right) + 2\frac{d^2u}{dz\,dy} \cdot \frac{dz}{dy} + \left(\frac{d^2u}{dz^2}\right) \cdot \frac{dz^2}{dy^2} + \left(\frac{du}{dz}\right) \cdot \frac{d^2z}{dy^2} = 0 \ldots (4).$$

Now either differentiate (1) with respect to y, or (2) with respect to x; and since in the former case $\frac{dz}{dx}$ becomes $\frac{d^2z}{dy\,dx}$, and that in the latter $\frac{dz}{dy}$ beeomes $\frac{d^2z}{dx\,dy}$, and that $\frac{d^2z}{dy\,dx} = \frac{d^2z}{dx\,dy}$, the results will be identical.

Let equation (1) be differentiated with respect to y; and to do this put $\left(\frac{du}{dx}\right)$ and $\left(\frac{du}{dz}\right)$ for u in equation (2), whence we have

$$\frac{d^2u}{dx \cdot dy} + \frac{d^2u}{dx \cdot dz} \cdot \frac{dz}{dy} + \frac{d^2u}{dz \cdot dy} \cdot \frac{dz}{dx} + \left(\frac{d^2u}{dz^2}\right) \cdot \frac{dz}{dx} \cdot \frac{dz}{dy} + \left(\frac{dz}{dx}\right) \cdot \frac{d^2z}{dy \cdot dx} = 0 \ldots (5).$$

From the equations (3), (4), (5), $\frac{d^2z}{dx^2}$, $\frac{d^2z}{dy^2}$ and $\frac{d^2z}{dy\,dx}$ may be found, while from equations (1) and (2), $\frac{dz}{dx}$ and $\frac{dz}{dy}$ may be found.

118. From this it is obvious, that if $u = f(x, y, z)$, be an explicit function of x, y and z, where z is a function of x and y, that the differential coefficients of u may be found.

For $$\frac{du}{dx} = \left(\frac{du}{dx}\right) + \left(\frac{du}{dz}\right) \cdot \frac{dz}{dx} \ldots\ldots\ldots\ldots (1),$$

$$\frac{du}{dy} = \left(\frac{du}{dy}\right) + \left(\frac{du}{dz}\right) \cdot \frac{dz}{dy} \ldots\ldots\ldots\ldots (2),$$

$$\frac{d^2u}{dx^2} = \left(\frac{d^2u}{dx^2}\right) + 2\frac{d^2u}{dzdx}\cdot\frac{dz}{dx} + \left(\frac{d^2u}{dz^2}\right)\cdot\frac{dz^2}{dx^2} + \left(\frac{du}{dz}\right)\cdot\frac{d^2z}{dx^2} \quad ..(3),$$

$$\frac{d^2u}{dy^2} = \frac{d^2u}{dy^2} + 2\frac{d^2u}{dzdy}\cdot\frac{dz}{dy} + \left(\frac{d^2u}{dz^2}\right)\cdot\frac{dz^2}{dy^2} + \left(\frac{du}{dz}\right)\cdot\frac{d^2z}{dy^2} \quad .....(4),$$

$$\frac{d^2u}{dxdy} = \frac{d^2u}{dxdy} + \frac{d^2u}{dxdz}\cdot\frac{dz}{dy} + \frac{d^2u}{dydz}\cdot\frac{dz}{dx} + \left(\frac{d^2u}{dz^2}\right)\cdot\frac{dz}{dx}\cdot\frac{dz}{dy} + \frac{dx}{dz}\cdot\frac{d^2z}{dydx} \quad .....(5).$$

From which $\frac{dz}{dx}$, $\frac{dz}{dy}$, $\frac{d^2z}{dx^2}$, $\frac{d^2z}{dy^2}$, $\frac{d^2z}{dydx}$ may be found.

By a similar process the differentials of the third and higher orders may be obtained.

ELIMINATION BY MEANS OF DIFFERENTIATION.

119. We have seen that if a constant quantity be connected with the function by the signs $\pm$, it disappears from the differential coefficients. Should it however be multiplied into the function or any term of the function, it will still appear in the value of the differential coefficient.

Thus if $u = 0$ be a function of x and y, involving a constant a, both $u = 0$ and $du = 0$ will contain a, but between these two equations it may be eliminated, and an equation will arise independent of a, which is called a differential equation.

Thus, let $y = ax^2$; $\therefore \frac{dy}{dx} = 2ax$, $= \frac{2y}{x}$;

an equation from which a has disappeared.

Irrational and transcendental quantities may also be eliminated by differentiation.

Thus, let $y = (a^2 + x^2)^{\frac{m}{n}}$;

$$\therefore \frac{dy}{dx} = 2x\cdot\frac{m}{n}\cdot(a^2 + x^2)^{\frac{m}{n}-1} = \frac{2mx(a^2 + a^2)^{\frac{m}{n}}}{n(a^2 + x^2)} = \frac{2mxy}{n(a^2 + x^2)}.$$

If there be two constants as a and b involved in the equation $y = f(x)$, then to eliminate them, the equations $u = 0$, $du = 0$, and $d^2u = 0$ must be combined.

Ex. 1. Let $u = y - ax^2 - bx = 0$, or $y = ax^2 + bx$;

$$\therefore \frac{dy}{dx} = 2ax + b, \quad \frac{d^2y}{dx^2} = 2a; \quad b = \frac{dy}{dx} - x\frac{d^2y}{dx^2};$$

$$\therefore y - \frac{x^2}{2}\cdot\frac{d^2y}{dx^2} - x\frac{dy}{dx} + x^2\frac{d^2y}{dx^2} = 0;$$

$$\therefore \frac{d^2y}{dx^2} - \frac{2}{x}\cdot\frac{dy}{dx} + \frac{2y}{x^2} = 0.$$

Ex. 2. Let $y = a \cdot \cos mx + b \cdot \sin mx$; eliminate a and b.

$$\frac{dy}{dx} = -ma \sin mx + mb \cos mx,$$

$$\frac{d^2y}{dx^2} = -m^2 a \cos mx - m^2 b \sin mx$$

$$= -m^2\{a \cos mx + b \sin mx\} = -m^2 y;$$

$$\therefore \frac{d^2y}{dx^2} + m^2 y = 0.$$

Ex. 3. Let $y = ae^{2x} \sin(3x + b)$; eliminate a and b.

$$\frac{dy}{dx} = 2ae^{2x} \sin(3x + b) + 3ae^{2x} \cos(3x + b)$$

$$= 2y + 3y \cot(3x + b),$$

$$\frac{d^2y}{dx^2} = 4ae^{2x} \sin(3x + b) + 6ae^{2x} \cos(3x + b)$$

$$+ 6ae^{2x} \cos(3x + b) - 9ae^{2x} \sin(3x + b)$$

$$= -5y + 12y \cot(3x + b)$$

$$= -5y + 4\frac{dy}{dx} - 8y;$$

$$\therefore \frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 13y = 0.$$

120. Again, if $u = f(xyz) = 0$, or $z = f(xy)$.

We may by means of the partial differential coefficients $\frac{dz}{dy}$ and $\frac{dz}{dx}$ eliminate two constants from $z = f(xy)$, and by proceeding to the second differential, we have three other equations for $\frac{d^2z}{dx^2}$, $\frac{d^2z}{dy\,dx}$ and $\frac{d^2z}{dy^2}$, and therefore five constants

may be eliminated; and not only constants, but indeterminate functions.

Ex. 4. Let $z = f(ax + by)$; eliminate the arbitrary function.

$$\text{Let } (ax + by) = v; \quad \therefore\ z = f(v),$$

$$\text{and } \frac{dz}{dx} = \frac{dz}{dv}\cdot\frac{dv}{dx}. \quad \text{But } \frac{dz}{dv} = f'(v), \text{ and } \frac{dv}{dx} = a;$$

$$\therefore\ \frac{dz}{dx} = af'(v); \quad \frac{dz}{dy} = \frac{dz}{dv}\cdot\frac{dv}{dy} = f'(v).b,$$

$$\text{or } b.\frac{dz}{dx} = abf'(v), \text{ and } a.\frac{dz}{dy} = ab.f'(v);$$

$$\therefore\ b\frac{dz}{dx} - a\frac{dz}{dy} = 0, \text{ or } bp - aq = 0.$$

As an example. Let $z = \sin(ax + by)$;

$$\therefore\ p = a\cos(ax + by), \quad q = b\cos(ax + by);$$

$$\therefore\ bp - aq = 0,$$

and similarly if $z = (ax + by)^n$, or $z = \log(ax + by)$, the differential equation will be verified.

Ex. 5. Let $z = (x + y)^m \phi(x^2 - y^2)$; eliminate the function.

$$p = m.(x + y)^{m-1}\phi(x^2 - y^2) + 2(x + y)^m \phi'(x^2 - y^2).x \ldots (1),$$

$$q = m.(x + y)^{m-1}\phi(x^2 - y^2) - 2(x + y)^m \phi'(x^2 - y^2).y \ldots (2).$$

Multiply (1) by (y), and (2) by x, and add;

$$\therefore\ yp + xq = m(x + y)^m \phi(x^2 - y^2) = mz,$$

$$\text{or } py + qx = mz.$$

Ex. 6. Let $zx = f\left(\frac{x}{y}\right)$; or $z = \frac{1}{x}.f\left(\frac{x}{y}\right)$;

$$\therefore\ p = -\frac{1}{x^2}f\left(\frac{x}{y}\right) + \frac{1}{xy}.f'\left(\frac{x}{y}\right),$$

$$q = \ldots\ldots\ldots\ldots - \frac{1}{y^2}.f'\left(\frac{x}{y}\right);$$

$$\therefore\ px + z = \frac{1}{y}.f'\left(\frac{x}{y}\right); \quad qy = -\frac{1}{y}.f'\left(\frac{x}{y}\right);$$

$$\therefore\ px + qy + z = 0.$$

Ex. 7. Let $z = f(y + ax) + \phi(y - ax)$,

$$p = a.f'(y + ax) - a\phi'(y - ax),$$

$$q = f'(y + ax) + \phi'(y - ax);$$

$$\therefore \frac{d^2z}{dx^2} = a^2 f''(y + ax) + a^2\phi''(y - ax),$$

$$\frac{d^2z}{dy^2} = f''(y + ax) + \phi''(y - ax);$$

$$\therefore \frac{d^2z}{dx^2} - a^2.\frac{d^2z}{dy^2} = 0.$$

The equation to vibrating chords, which also occurs in other investigations in Natural Philosophy.

Ex. 8. If $z = xf(a) + y\phi(a) + \psi(a)$.....(1),

$$0 = xf'(a) + y\phi'(a) + \psi'(a).....(2),$$

where $f'(a) = \frac{d.f(a)}{da}$, to eliminate the arbitrary functions:

Differentiate (1) successively with respect to x and y, considering a as a function of x and y;

$$\therefore \frac{dz}{dx} = f(a) + \{xf'(a) + y\phi'(a) + \psi'(a)\}\frac{da}{dx},$$

$$\frac{dz}{dy} = \phi(a) + \{xf'(a) + y\phi'(a) + \psi'(a)\}\frac{da}{dy};$$

therefore from (2), $\frac{dz}{dx} = f(a)$; $\frac{dz}{dy} = \phi(a)$;

$$\therefore \frac{dz}{dx} = F\left(\frac{dz}{dy}\right);$$

$$\therefore \frac{d^2z}{dx^2} = F'\left(\frac{dz}{dy}\right).\frac{d^2z}{dxdy}.....(3),$$

$$\frac{d^2z}{dydx} = F'\left(\frac{dz}{dy}\right).\frac{d^2z}{dy^2}.....(4),$$

whence multiplying crossways,

$$\left(\frac{d^2z}{dx^2}\right)\left(\frac{d^2z}{dy^2}\right) = \left(\frac{d^2z}{dxdy}\right)^2,$$

which is the equation to developable surfaces.

Ex. 9. $y = xe^{cx}$; eliminate c.

$$x\,dy - y\,dx = y(\log y - \log x).$$

Ex. 10. Eliminate a and b from $y^2 = ax + bx^2$.

Ex. 11. If $y = a\sin x + b\sin 2x$, shew that

$$\frac{d^4y}{dx^4} + 5\frac{d^2y}{dx^2} + 4y = 0.$$

Ex. 12. If $z = \frac{x^2}{2} + f(y + \log x)$; $\quad px - q = x^2$.

Ex. 13. If $z = f\left(\frac{y^2 - x^2}{x}\right)$; $\quad 2xyp + (x^2 + y^2)q = 0$.

Ex. 14. $x^2 + y^2 + z^2 = f(ax + by + c)$.

$$(y - bz)p - (x - az)q = bx - ay.$$

Ex. 15. $z = ax + by + c$; eliminate a, b, c.

$$\frac{d^2z}{dx^2}\cdot\frac{d^2y}{dx^2} - \frac{d^2z}{dx^2}\cdot\frac{d^2y}{dx^2} = 0.$$

Ex. 16. If $z = xf\left(\frac{y}{x}\right) + \phi(xy)$, shew that

$$x^2\cdot\frac{d^2z}{dx^2} = y^2\cdot\frac{d^2z}{dy^2}.$$

CHAPTER IX.

MAXIMA AND MINIMA OF FUNCTIONS OF TWO VARIABLES.

121. If $u = f(x, y)$ be an equation between the function u, and the two independent variables, x and y, there may be some particular value of x, and also of y, which will make the function greater or less than the values which immediately precede or follow it. It is then a maximum or minimum. We proceed to find the relation between the differential coefficients, when this circumstance takes place.

122. Let u_1 be the value of u, when $x + h$ and $y + k$ are written for x and y respectively; and u_2 the value of u, when $x - h$ and $y - k$ are substituted for the same quantities. Also put A for $\frac{d^2u}{dx^2}$, B for $\frac{d^2u}{dy\,dx}$, and C for $\frac{d^2u}{dy^2}$. Then

$$u_1 = u + \frac{du}{dx}h + \frac{du}{dy}k + \tfrac{1}{2}\{Ah^2 + 2Bhk + Ck^2\} + \&c.$$

and $$u_2 = u - \left(\frac{du}{dx}h + \frac{du}{dy}k\right) + \tfrac{1}{2}\{Ah^2 + 2Bhk + Ck^2\} - \&c.$$

Now since the values of h and k may be assumed so small that, (as long as the differential coefficients $\frac{du}{dx}$ and $\frac{du}{dy}$ remain finite) the algebraical sign of $u_1 - u$ and $u_2 - u$ will depend upon that of the term $\left(\frac{du}{dx}h + \frac{du}{dy}k\right)$, it is manifest, that if this term exist, $u_1 - u$ and $u_2 - u$ cannot be both positive or both negative, or there cannot be a minimum or maximum of u. Therefore at a maximum or minimum $\frac{du}{dx}h + \frac{du}{dy}k$ must = 0. A condition which can only be fulfilled, since h and k are independent quantities, by making $\frac{du}{dx} = 0$, and $\frac{du}{dy} = 0$. Hence at a maximum or minimum,

$$u_1 - u = \tfrac{1}{2}(Ah^2 + 2Bhk + Ck^2) + \&c.$$

$$= \frac{h^2}{1.2}\{A + 2Bn + Cn^2\} + \&c., \text{ if } k = nh.$$

Therefore the sign of $u_1 - u$, and also of $u_2 - u$, will depend upon that of the coefficient of $\frac{h^2}{2}$, that is, upon $A + 2Bn + Cn^2$. Hence, this term must not change its sign whatever be the value of n; which it will not do, if it can be put under the form of the sum of two squares, as $(x + \alpha)^2 + \beta^2$.

$$\text{Now } A + 2Bn + Cn^2 = \frac{1}{C}\{CA + 2BCn + C^2n^2\}$$

$$= \frac{1}{C}\{CA - B^2 + (B + Cn)^2\}$$

$$= \frac{1}{C}\left\{CA - B^2 + C^2\left(\frac{B}{C} + n\right)^2\right\},$$

which is of the requisite form, if CA be not less than B^2: or to have a maximum or minimum of a function of two variables, we must first have $\frac{du}{dx} = 0$ and $\frac{du}{dy} = 0$; and secondly, $\frac{d^2u}{dx^2} \times \frac{d^2u}{dy^2}$ not less than $\left(\frac{d^2u}{dydx}\right)^2$.

123. It is obvious that $\frac{d^2u}{dx^2}$ and $\frac{d^2u}{dy^2}$ must have the same algebraical sign; and also if they be both *negative*, u is a maximum, if both *positive*, u is a minimum.

If the second differential coefficient of u become = 0, when the first does, there will not be a maximum or minimum, unless the third differential coefficient vanishes, and the fourth neither vanishes nor changes its sign, whatever be the value of n.

Ex. 1. Let $u = x^3 + y^3 - 3axy$,

$$\frac{du}{dx} = 3x^2 - 3ay = 0; \quad \therefore\ y = \frac{x^2}{a},$$

$$\frac{du}{dy} = 3y^2 - 3ax = 0; \quad \therefore\ y^2 - ax = \frac{x^4}{a^2} - ax = 0;$$

$\therefore\ x = 0$, and $x^3 - a^3 = 0$; whence $x = a$; the other two roots are impossible; and $y = \frac{x^2}{a} = a$; or $= 0$.

$$\text{Also } \frac{d^2u}{dx^2} = 6x, \quad \frac{d^2u}{dy^2} = 6y, \quad \text{and} \quad \frac{d^2u}{dydx} = -3a.$$

If $x = 0$, $A = 0$, $C = 0$, and $B = -3a$.

If $x = a$, $A = 6a$, $C = 6a$, $B = -3a$,

$$AC = 36a^2, \text{ and } B^2 = 9a^2;$$

$\therefore x = a$, $\because A$ is +, gives a minimum, and $u = -a^3$; $x = 0$ gives neither a maximum nor minimum.

Ex. 2. $u = x^3y^2(a - x - y)$.

$$\frac{du}{dx} = 3x^2y^2(a - x - y) - x^3y^2 = 0,$$

$$\frac{du}{dy} = 2x^3y(a - x - y) - x^3y^2 = 0;$$

$$\therefore 3(a - x - y) = x; \quad 2(a - x - y) = y; \quad \therefore 2x = 3y;$$

$$\therefore 3a - 3x - 2x = x, \quad \text{or } x = \frac{a}{2},$$

$$2a - 3y - 2y = y, \quad \text{or } y = \frac{a}{3};$$

$$\therefore a - x - y = a - \frac{a}{2} - \frac{a}{3} = \frac{a}{6},$$

$$\frac{d^2u}{dx^2} = 6xy^2(a - x - y) - 6x^2y^2 = 6\left\{\frac{a}{2}\cdot\frac{a^2}{9}\cdot\frac{a}{6} - \frac{a^2}{4}\cdot\frac{a^2}{9}\right\} = -\frac{a^4}{9},$$

$$\frac{d^2u}{dy^2} = 2x^3(a - x - y) - 4x^3y = 2\left\{\frac{a^3}{8}\cdot\frac{a}{6} - 2\cdot\frac{a^3}{8}\cdot\frac{a}{3}\right\} = -\frac{a^4}{8},$$

$$\frac{d^2u}{dxdy} = 6x^2y(a - x - y) - 3x^2y^2 - 2x^3y = \frac{a^4}{12} - \frac{a^4}{12} - \frac{a^4}{12} = -\frac{a^4}{12};$$

$$\therefore AC = \frac{a^8}{72}, \text{ and } B^2 = \frac{a^8}{144}; \quad \therefore AC \text{ is} > B^2;$$

and $u = \frac{a^3}{8} \times \frac{a^2}{9} \times \frac{a}{6} = \frac{a^6}{432}$ is a maximum, $\because A$ is $-$.

Ex. 3. $u = (x + 1)\cdot(y + 1)\cdot(z + 1)$, where $a^xb^yc^z = A$;

$$\therefore x\log a + y\log b + z\log c = \log A.$$

Now $$\frac{du}{dx} = (y + 1)\cdot\left\{z + 1 + (x + 1)\cdot\frac{dz}{dx}\right\} = 0,$$

$$\frac{du}{dy} = (x + 1)\cdot\left\{z + 1 + (y + 1)\cdot\frac{dz}{dy}\right\} = 0,$$

$$\text{and } \frac{dz}{dx} = -\frac{\log a}{\log c}, \text{ and } \frac{dz}{dy} = -\frac{\log b}{\log c};$$

$$\therefore\ z + 1 - (x+1).\frac{\log a}{\log c} = 0,$$

or $(z+1)\log c = (x+1)\log a$, and $\therefore\ c^{z+1} = a^{x+1}$.

$$\text{Also } z + 1 - (y+1).\frac{\log b}{\log c} = 0;$$

and $\therefore\ c^{z+1} = b^{y+1} = a^{x+1}$.

$$\text{Also } \because\ z + 1 = (x+1).\frac{\log a}{\log c} = (y+1).\frac{\log b}{\log c};$$

$$\therefore\ z = \frac{(x+1)\log a - \log c}{\log c},$$

$$y = \frac{(x+1)\log a - \log b}{\log b};$$

$$\therefore\ x\log a + (x+1)\log a - \log b + (x+1)\log a - \log c = \log A;$$

$$\therefore\ 3x\log a + 2\log a - \log bc = \log A;$$

$$\therefore\ x = \frac{\log Abc - 2\log a}{3\log a};$$

$$\therefore\ x + 1 = \frac{\log Abc + \log a}{3\log a} = \frac{\log (Abca)}{3\log a},$$

$$y + 1 = \frac{\log (Abca)}{3\log b}, \text{ and } z + 1 = \frac{\log (Abca)}{3\log c};$$

$$\therefore\ u = \frac{(\log Aabc)^3}{27\log a \,.\, \log b \,.\, \log c}.$$

Ex. 4. Inscribe the greatest triangle within a given circle.



R the radius.

a, b, c the sides.

$\theta = \angle B$, $\phi = \angle C$.

$$\text{But } u = \frac{bc}{2}.\sin A = \frac{bc}{2}.\sin(\theta+\phi);$$

and $b = 2R\,.\,\sin\theta$; $c = 2R\,.\,\sin\phi$;

$$\therefore\ u = 2R^2 \sin\theta \sin\phi \sin(\phi+\theta) = \text{maximum};$$

$$\therefore \frac{du}{d\theta} = 2R^2\{\cos\theta \cdot \sin(\phi+\theta) + \sin\theta \cdot \cos(\phi+\theta)\}\sin\phi = 0,$$

$$\text{and } \frac{du}{d\phi} = 2R^2\{\cos\phi \cdot \sin(\phi+\theta) + \sin\phi \cdot \cos(\phi+\theta)\}\sin\theta = 0;$$

$$\therefore \sin(\phi+2\theta) = 0 = \sin\pi, \text{ and } \sin(\theta+2\phi) = 0 = \sin\pi;$$

$$\therefore \phi + 2\theta = \pi, \text{ and } \theta + 2\phi = \pi;$$

$$\therefore \theta - \phi = 0, \text{ or } \theta = \phi; \quad \therefore 3\theta = \pi, \text{ and } \theta = 60^0 = \phi,$$

and $\therefore A = 60^0$; and the triangle is equiangular.

Ex. 5. Inscribe the greatest parallelopipedon within a given ellipsoid.

Let $2x$, $2y$, $2z$ be the edges,

$2a$, $2b$, $2c$ the principal diameters of the ellipsoid;

$$\therefore u = 8xyz, \text{ and } \frac{z^2}{c^2} + \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1;$$

$$\therefore z^2 = c^2\left(1 - \frac{x^2}{a^2} - \frac{y^2}{b^2}\right);$$

$$\therefore \frac{du}{dx} = 8yz + 8yx\frac{dz}{dx} = 0,$$

$$\frac{du}{dy} = 8xz + 8yx\frac{dz}{dy} = 0.$$

$$\text{But } \frac{dz}{dx} = -\frac{x}{z}\cdot\frac{c^2}{a^2}, \text{ and } \frac{dz}{dy} = -\frac{y}{z}\cdot\frac{c^2}{b^2};$$

$$\therefore z - \frac{x^2}{z}\cdot\frac{c^2}{a^2} = 0, \text{ and } z - \frac{y^2}{z}\cdot\frac{c^2}{b^2} = 0;$$

$$\therefore \frac{z^2}{c^2} = \frac{x^2}{a^2}, \text{ and } \frac{z^2}{c^2} = \frac{y^2}{b^2} = \quad \therefore \frac{x^2}{a^2};$$

$$\therefore \frac{3x^2}{a^2} = 1, \; x = \frac{a}{\sqrt{3}}; \quad \therefore \frac{y^2}{b^2} = \frac{z^2}{c^2} = \frac{1}{3}; \quad \therefore y = \frac{b}{\sqrt{3}}; \quad \therefore z = \frac{c}{\sqrt{3}};$$

$$\text{and } u = \frac{8}{3}\frac{abc}{\sqrt{3}}.$$

But if V = volume of ellipsoid; $V = \frac{4}{3}\pi abc$;

$$\therefore u = \frac{2V}{\pi\sqrt{3}} \text{ or } u : V :: 2 : \pi\sqrt{3}.$$

124. In the following examples, some only of the steps are indicated.

Ex. 6. If $u = x^4 + y^4 - 4axy^2$; find x and y, when u = maximum or minimum.

$$x = \pm a\sqrt{2}; \quad y = \pm a\sqrt[4]{8},$$

and $x = 0$; $y = 0$; give minima.

Ex. 7. $u = a\{\sin x + \sin y + \sin(x + y)\}$.

$x = 60$; $y = 60$; give $u = 3a\dfrac{\sqrt{3}}{2}$ a maximum.

Ex. 8. Given the perimeter of a triangle, shew that its area is greatest when it is equilateral.

Ex. 9. Divide a quantity a into three such parts x, y, z, that $u = x^m y^n z^p$ may be a maximum; and shew that it is a maximum and not a minimum.

$$x = \frac{ma}{m+n+p}; \quad y = \frac{na}{m+n+p}; \quad z = \frac{pa}{m+n+p}.$$

Ex. 10. Given the surface of a rectangular parallelopipedon; find when its content is a maximum.

If x, y, z be the edges of the solid:

$$\text{Surface} = 2xy + 2xz + 2yz = 2a^2,$$

and $u = xyz$ a maximum,

whence $x = y = z = a$; and solid is a cube.

Ex. 11. If the content of the rectangular parallelopipedon be given, find its form when the surface is a minimum.

It is a cube, as in the preceding question.

Ex. 12. Let $u = ax + by + cz$ a maximum,

and $x^2 + y^2 + z^2 = 1$; find x, y, z;

$$\therefore u = ax + by + c\sqrt{1 - x^2 - y^2};$$

$$\therefore \frac{du}{dx} = a - \frac{cx}{z} = 0; \quad \therefore az = cx,$$

$$\frac{du}{dy} = b - \frac{cy}{z} = 0; \quad \therefore bz = cy;$$

$$\therefore (a^2 + b^2 + c^2)z^2 = c^2(x^2 + y^2 + z^2) = c^2;$$

$$\therefore z = \frac{c}{\sqrt{a^2 + b^2 + c^2}}; \quad x = \frac{a}{\sqrt{a^2 + b^2 + c^2}}; \quad y = \frac{b}{\sqrt{a^2 + b^2 + c^2}}.$$

Ex. 13. Given the radius of a circle described about a triangle; find its form when the perimeter is a minimum.

If θ and ϕ be two of the angles, and if r be the radius of the circle,

$$u = 2r\{\sin\theta + \sin\phi + \sin(\theta + \phi)\};$$

whence $\theta = \phi = 60$; and the triangle is equilateral.

Ex. 14. Given the sum of the three axes of an ellipsoid; find them when the volume of the ellipsoid is greatest.

If $2x$, $2y$, $2z$ be the three axes,

$$u = \frac{4}{3}\pi xyz, \text{ a maximum,}$$

and $2x + 2y + 2z = 2a$, the given length; whence

$x = y = z = a$, and ellipsoid becomes a sphere.

Ex. 15. Find that point within a triangle, from which if lines be drawn to the angular points, the sum of their squares shall be a minimum.



Let ABC be a triangle, and P a point within it, a, b, c, the sides of the triangle.

Draw PN, AD perpendicular to the base; join AP, BP, CP.

Let $CN = x$; $NP = y$; then $AD = b\sin C$; $CD = b\cos C$.

Then $CP^2 = x^2 + y^2$; $BP^2 = y^2 + (a-x)^2 = y^2 + x^2 + a^2 - 2ax$,

$$AP^2 = (b\cos C - x)^2 + (b\sin C - y)^2$$

$$= b^2 + x^2 + y^2 - 2b(x\cos C + y\sin C);$$

$$\therefore u = 3x^2 + 3y^2 + a^2 + b^2 - 2ax - 2b(x\cos C + y\sin C);$$

$$\therefore x = \tfrac{1}{3}(a + b\cos C); \quad y = \tfrac{1}{3}b\sin C;$$

$$\therefore CP = \sqrt{x^2 + y^2} = \tfrac{1}{3}\sqrt{2a^2 + 2b^2 - c^2}.$$

The point P is the centre of gravity of the triangle.

Ex. 16. Find a point within a triangle, from which if perpendiculars be let fall upon the sides, the sum of their squares shall be a minimum.

ABC the triangle as before, P the point within it, draw PN, PM, PQ respectively perpendicular to CB, CA, AB.

Let $CN = x$; $PM = p$;

$NP = y$; $PQ = q$;

Then $u = y^2 + p^2 + q^2$.

Now if δ be the perpendicular from a point (α, β) on a line $y = mx + b$,

$$\delta = \frac{\beta - b - m\alpha}{\sqrt{m^2+1}}.$$

(1^0) If $\delta = p$; $\beta = y$; $\alpha = x$; $b = 0$; $m = \tan C$;

$$\therefore\ p = \frac{y - x\tan C}{\sec C} = y\cos C - x\sin C.$$

(2^0) If $\delta = q$; $m = -\tan B$; $b = a\tan B$;

$$\therefore\ q = \frac{y - (a-x)\tan B}{\sec B} = y\cos B - (a-x)\sin B;$$

$$\therefore\ u = y^2 + (y\cos C - x\sin C)^2 + \{y\cos B - (a-x)\sin B\}^2,$$

whence $$y = \frac{a\sin A\,.\,\sin B\,.\,\sin C}{2\,(1 - \cos^2 B\,.\,\cos^2 C + \sin B\,.\,\sin C\,.\,\cos B\,.\,\cos C)}$$

$$= \frac{abc\sin A}{a^2+b^2+c^2};$$

and $$\therefore\ p = \frac{abc\sin B}{a^2+b^2+c^2};\quad q = \frac{abc\sin C}{a^2+b^2+c^2}.$$

Ex. 17. Find a point P within a quadrilateral figure $ABCD$, from which if lines be drawn to the angular points, the sum of their squares shall be the least possible.



$AB = a$; $BC = b$;

$AD = c$; $AN = x$; $NP = y$;

$$\therefore\ u = x^2 + y^2 + y^2 + (a-x)^2 + (b\sin B - y)^2 + (a - x - b\cos B)^2 + (c\sin A - y)^2 + (x - c\cos A)^2$$

$$= 4y^2 + 2x^2 + 2(a-x)^2 - 2b\{y\sin B + (a-x)\cos B\} + b^2 + c^2 - 2c(y\sin A + x\cos A).$$

whence $x = \frac{1}{4}(2a + c\cos A - b\cos B)$,

$$y = \tfrac{1}{4}(c\sin A + b\sin B).$$

Also $\because \dfrac{d^2u}{dx^2} = 8 = \dfrac{d^2u}{dy^2}$; $\dfrac{d^2u}{dxdy} = 0$; u is a minimum.

Ex. 18. Of all triangular pyramids of a given base and altitude, to find that which has the least surface.

Let a, b, c be the sides of the base, h the altitude of the pyramid, θ, ϕ, ψ the inclination of the faces to the base.

Then because if p be perpendicular from vertex on the side a, $p\sin\theta = h$, and area of face $= \frac{1}{2}ap = \frac{1}{2}ah\operatorname{cosec}\theta$;

$$\therefore\ u = \tfrac{1}{2}h(a\operatorname{cosec}\theta + b\operatorname{cosec}\phi + c\operatorname{cosec}\psi).....(1).$$

Also since the base of the pyramid may be divided into three triangles, whose altitudes are respectively

$$h\cot\theta,\ h\cot\phi,\ h\cot\psi;\ \text{if } m^2 \text{ be its area,}$$

$$m^2 = \tfrac{1}{2}h(a\cot\theta + b\cot\phi + c\cot\psi).....(2),$$

from which combined with (1) a minimum, we have $\theta = \phi = \psi$, or the faces are equally inclined to the base.

Ex. 19. Two points P and Q are given above a plane; find a point R in a plane, such that $PR + RQ$ may be a minimum.

Let the given plane be that of xy; from P and Q draw lines perpendicular to it, let the axis of z pass through P, and the axis of x pass through the foot of the perpendicular from Q.

Then if c = the co-ordinate of P, a and b that of Q, x and y of R;

$$\therefore\ u = PR + QR = \sqrt{x^2 + y^2 + c^2} + \sqrt{y^2 + (a-x)^2 + b^2};$$

$$\therefore\ \frac{du}{dx} = \frac{x}{\sqrt{x^2 + y^2 + c^2}} - \frac{a-x}{\sqrt{y^2 + (a-x)^2 + b^2}} = 0...(1);$$

$$\therefore\ \frac{du}{dy} = \frac{y}{\sqrt{x^2 + y^2 + c^2}} - \frac{y}{\sqrt{y^2 + (a-x)^2 + b^2}} = 0...(2).$$

From (2), $y = 0$, and therefore the point R is in the axis of x.

From (1), $\dfrac{x}{PR} = \dfrac{a-x}{QR}$; or the cosines of the angles which

PR and QR make with the axis of x are equal,

$$\text{also } \because \frac{x}{\sqrt{x^2+c^2}} = \frac{a-x}{\sqrt{(a-x)^2+b^2}}; \quad x = \frac{ac}{b+c}.$$

125. When $u = f(xyz)$ is a maximum or minimum, we must put $\frac{du}{dx} = 0$; $\frac{du}{dy} = 0$; $\frac{du}{dz} = 0$; and the equation of condition is $(AC - B^2)(AD - E^2) > (AF - BE)^2$ *,

$$\text{where } A = \frac{d^2u}{dx^2}; \quad B = \frac{d^2u}{dy\,dx}; \quad C = \frac{d^2u}{dy^2},$$

$$\text{and } D = \frac{d^2u}{dz^2}; \quad E = \frac{d^2u}{dz\,dx}; \quad F = \frac{d^2u}{dz\,dy}.$$

Ex. 20. $u = ax^2y^3z^4 - x^3y^3z^4 - x^3y^4z^4 - x^3y^3z^5 =$ a maximum.

$$x = \frac{1}{5}a; \quad y = \frac{3a}{10}; \quad z = \frac{2a}{5}.$$

Ex. 21. $u = \dfrac{xyz}{(x+a)(x+y)(y+z)(z+e)} =$ maximum.

$$x = \sqrt[4]{a^3e}; \quad y = \sqrt[4]{a^2e^2}; \quad z = \sqrt[4]{ae^3}.$$

* Lacroix, *Calcul. Diff.* Vol. 1. Art. 166.

CHAPTER X.

EQUATIONS TO CURVES.

126. We proceed to treat briefly of the equations to a straight line, to the circle, the conic sections, and some other curves, which will be frequently referred to in the succeeding pages; but for complete investigations of the properties of the conic sections and curves in general, we must refer to works expressly written on these subjects: since the object of this Chapter is merely to furnish the student with such a knowledge of the nature of certain curves, as may make the applications of the Differential Calculus to them obvious and interesting.

It is necessary to make some prefatory remarks before the nature of these equations is explained.

From a point A, assumed at pleasure, draw two lines, Ay, Ax, perpendicular to each other; then the position of a point P, situated within the angle yAx, will be known if the perpendicular distances PN and PM from P upon Ax, and Ay be known.



Ay and Ax are called axes, and PM and PN the *ordinates* of the point P: but since $AN = PM$, the line PM is seldom drawn, but the position of P determined by taking AN equal to it, and then from N drawing NP perpendicular to Ax.

NP is then called the ordinate, and AN the abscissa; and AN and NP the co-ordinates of P.

Ax is termed the axis of abscissas, and Ay the axis of ordinates.

127. Now a curve is a series of points, and if their respective distances from two such axes as Ay and Ax be known, the curve itself may be drawn. Also every possible equation $y = f(x)$ may be represented by a series of points; for if it be assumed that the values of x may be taken along the line Ax, and those of y be drawn perpendicular to the axis, we shall have, when $x = AN$, $y = f(AN)$, which may be represented by some line as NP. Hence Ax is called the axis of x, and Ay the axis of y, and the point A the origin of the co-ordinates; since the values of x and y commence at A.

128. If $AN = a$, and $NP = b$,

then $x = a$, and $y = b$,

are the equations to a point P.



If also we take $AN_1 = AN$, and $MP_1 = MP$, and complete the rectangle PP_2, then unless some assumption be made with respect to the algebraical sign of a and b, we shall be unable to tell in which of the angles yAx, yAx_1, x_1Ay_1, or y_1Ax the point is, since P, P_1, P_2, P_3, in the annexed figure, are each at the same distances from the lines yAy_1 and xAx_1.

If however the values of x, when measured from A along Ax, or to the right hand of A, be reckoned positive, and those in the direction of Ax_1, or to the left of A, be negative; and if the ordinates which are above xAx_1 be called positive, and negative when measured below the same line, no difficulty can arise, and then

$$\left.\begin{aligned} x &= +a \\ y &= +b \end{aligned}\right\} \text{ determines the point } P,$$

$$\left.\begin{aligned} x &= -a \\ y &= +b \end{aligned}\right\} \ldots\ldots\ldots\ldots\ldots\ldots\ P_1,$$

$$\left.\begin{aligned} x &= -a \\ y &= -b \end{aligned}\right\} \ldots\ldots\ldots\ldots\ldots\ldots\ P_2,$$

$$\left.\begin{aligned} x &= +a \\ y &= -b \end{aligned}\right\} \ldots\ldots\ldots\ldots\ldots\ldots\ P_3.$$

Cor. 1. The distance of a point P from the origin, or

$$AP = \sqrt{AN^2 + PN^2} = \sqrt{x^2 + y^2}.$$



Cor. 2. The distance between two points P and P_1 is thus found.

A the origin. $AN = x$, $AN_1 = x_1$,

$$NP = y, \quad NP_1 = y_1;$$

$$\therefore PP_1 = \text{distance} = \sqrt{Pm^2 + P_1m^2}$$

$$= \sqrt{(AN_1 - AN)^2 + (P_1N_1 - PN)^2}$$

$$= \sqrt{(x_1 - x)^2 + (y_1 - y)^2}.$$

129. If in the curve CPQ the relation between PN and AN be known, and PN be a function of AN, the equation to the curve is said to be known; and from it, the curve itself may be drawn. Sometimes the equation $y = f(x)$ is to be found from some given property of the curve; as in the circle, of which the property is, that all lines drawn from the centre to the circumference are equal. To questions of this description our attention in this Chapter will be solely directed.



When AN and NP are drawn at right angles to each other, the equation $y = f(x)$ is called an equation to rectangular co-ordinates.

130. Many curves however cannot be expressed by an equation between rectangular co-ordinates. Such are the Spirals, which may be conceived to be described by the extremity of a straight line of variable length, which revolves round a fixed point, called the pole of the spiral. The revolving line (called the radius vector) may be considered as a function of the angle described.

Thus if S be the pole, SP the radius vector, SA the original position of SP, and $\angle ASP = \theta$; $r = f(\theta)$ is the equation to the spiral, and r and θ are called polar co-ordinates.



THE STRAIGHT LINE.

131. Find the equation to the straight line.

Ax, Ay the two axes of x and y.

$$\left.\begin{aligned} AN &= x \\ NP &= y \\ \angle PCA &= \theta \end{aligned}\right\}, \qquad \begin{aligned} &Bn \perp \text{ to } PN. \\ &AB = b. \end{aligned}$$



Then $$\frac{Pn}{Bn} = \frac{BA}{CA} = \tan\theta,$$

or $$\frac{y-b}{x} = \tan\theta = m, \text{ by writing } m \text{ for } \tan\theta;$$

$$\therefore\ y = mx + b.$$

Cor. 1. If the line be drawn through a given point; let α and β be the co-ordinates of the point; $\therefore$ if $x = \alpha$, $y = \beta$;

$$\therefore \beta = m\alpha + b, \quad \text{and} \quad y = mx + b;$$

$$\therefore y - \beta = m(x - \alpha).$$

Cor. 2. If the line be drawn through the origin, $AB = 0$; and $\therefore b = 0$; and $y = mx$ is the equation to a line drawn through A.

132. If two lines intersect, find the point of intersection.

Let $y = mx + b$, and $y = m_1x + b_1$ be the equations of the two lines; then, at the point of intersection, the values of the co-ordinates are the same for both lines;

$$\therefore mx + b = m_1x + b_1; \quad x = \frac{b_1 - b}{m - m_1},$$

$$\text{and } y = \frac{mb_1 - mb}{m - m_1} + b = \frac{mb_1 - m_1b}{m - m_1}.$$

133. Find the equation to a line passing through two given points.

Let $y = mx + b$ be the equation to the line where m and b are to be determined.

α and β, α_1 and β_1 the co-ordinates of the two points;

$$\therefore \beta = m\alpha + b; \quad \beta_1 = m\alpha_1 + b;$$

$$\therefore \beta - \beta_1 = m.(\alpha - \alpha_1); \quad \therefore m = \frac{\beta - \beta_1}{\alpha - \alpha_1}.$$

But $\because y = mx + b$, and $\beta = m\alpha + b$;

$$\therefore y - \beta = m.(x - \alpha) = \frac{\beta - \beta_1}{\alpha - \alpha_1}.(x - \alpha).$$

134. To find the angle which two straight lines make with each other at the point of intersection.

$$y = mx + b, \quad \text{and} \quad y = m_1x + b_1,$$

the equations to the two lines.

PQR and P_1QR_1 the lines.

From A draw An parallel to PR, and Am parallel to $P'R'$;

$$\therefore \angle nAm = \angle PQP';$$



$$\therefore\ PQP' = nAx - mAx = \tan^{-1} m - \tan^{-1} m_1,$$

$$\text{and } \tan PQP' = \frac{m - m_1}{1 + mm_1}.$$

Cor. 1. If the lines be parallel, $PQP' = 0$; $m - m_1 = 0$; and $m_1 = m$;

$$\left.\begin{array}{l} \therefore\ y = mx + b \\ \text{and } y = mx + b_1 \end{array}\right\}, \text{ are the equations to two parallel lines.}$$

Cor. 2. If the lines be perpendicular,

$$\tan PQP' = \frac{1}{0} = \frac{m - m_1}{1 + mm_1};$$

$$\therefore\ 1 + mm_1 = 0, \text{ and } m_1 = -\frac{1}{m};$$

therefore, if $y = mx + b$ be the equation to a line, $y = -\frac{1}{m}x + b_1$ is the equation to a line perpendicular to it.

135. Find the equation to a line drawn through a given point perpendicular to a given line.

$y = mx + b$, the equation to the given line, α and β the co-ordinates of the given point;

$$\therefore\ y = -\frac{1}{m}x + b_1 \text{ is the equation to the perpendicular,}$$

$$\text{also } \beta = -\frac{1}{m}\alpha + b_1, \text{ since it passes through } (\alpha, \beta);$$

$$\therefore\ (y - \beta) = -\frac{1}{m}(x - \alpha) \text{ is the equation required.}$$

136. Find the perpendicular distance of a given point from a given line.

$y = mx + b$ the equation to the given line, and (β, α) the given point;

$$\therefore\ (y - \beta) = -\frac{1}{m}(x - \alpha)$$

is the equation to a perpendicular from a given point upon the given line.

Then if δ be the distance required, and y_1 and x_1 the co-ordinates of the point of intersection of the given line with the perpendicular,

$$\delta = \sqrt{(x_1 - \alpha)^2 + (y_1 - \beta)^2} = (x_1 - \alpha) \cdot \sqrt{\frac{m^2 + 1}{m}}.$$

But $mx_1 + b = \beta - \frac{x_1 - a}{m}$;

$\therefore\ x_1(m^2+1) = m\beta + a - mb$; $\therefore\ x_1 = \frac{m\beta + a - mb}{m^2+1}$,

$$x_1 - a = \frac{m\beta - mb - m^2 a}{m^2+1} = \frac{m}{m^2+1}(\beta - b - ma);$$

$$\therefore\ \delta = \pm\frac{\beta - b - ma}{\sqrt{m^2+1}};$$

the upper or lower sign being taken according as the numerator is positive or negative.

Cor. 1. If the point be the origin, then $a = 0$; $\beta = 0$;

$$\therefore\ \delta = \frac{b}{\sqrt{m^2+1}} = \frac{y - mx}{\sqrt{m^2+1}}$$

$$= y\cos\theta - x\sin\theta,$$

since $m = \tan\theta$, and $\sqrt{m^2+1} = \sec\theta$.

Cor. 2. If the line pass through the origin, $b = 0$;

$$\therefore\ \delta = \frac{\beta - ma}{\sqrt{m^2+1}} = \beta\cos\theta - a\sin\theta.$$

Cor. 3. If neither the point be in the origin, nor the line pass through it.

$$\delta = \frac{\beta - ma - y + mx}{\sqrt{m^2+1}}$$

$$= \frac{(\beta - y) - m(a - x)}{\sqrt{m^2+1}}$$

$$= (\beta - y)\cos\theta - (a - x)\sin\theta.$$

137. Find the equation to a straight line, which cuts the axis of y at a distance b from the origin, and the axis of x at a distance a from the origin, in terms of b and a.

$y = mx + b$ the equation to the line,

when $x = 0$, $y = b$,

and $y = 0$, $x = a$; $\therefore\ ma + b = 0$; $\therefore\ m = -\frac{b}{a}$;

$\therefore\ y = -\frac{b}{a}x + b$; or $\frac{y}{b} + \frac{x}{a} = 1$, is the equation.

THE CIRCLE.

138. The circle is a curve of which the property is, that every point in its circumference is equi-distant from the centre.

Let a and β be the co-ordinates of the centre,

x and y of a point in the curve; a = radius.

Then the distance between two points a, β, and x, y

$$= \sqrt{(x-a)^2 + (y-\beta)^2} = a;$$

$$\therefore\ y^2 + x^2 - 2\beta y - 2ax + a^2 + \beta^2 - a^2 = 0,$$

is the equation to the circle.

Cor. 1. If the origin be in the circumference, and the axis of x pass through the centre.

$\beta = 0$, $a = a$; $\therefore\ y^2 + x^2 - 2ax = 0$, or $y^2 = 2ax - x^2$.

Cor. 2. If the origin be in the centre.

$a = 0$, $\beta = 0$; $\therefore\ y^2 + x^2 - a^2 = 0$, and $y^2 = a^2 - x^2$.



139. If S be a point in the circumference of a circle, to find the equation between SP and a perpendicular on the tangent, SY.

Join OP and draw $SN \perp PO$ produced,

$SP = r$; $SY = p$; $OP = a$; then $SY = PN$.

Now $SP^2 = SO^2 + OP^2 + 2OP \cdot ON$;

$$\therefore\ r^2 = 2a^2 + 2a(p-a) = 2ap;$$

$$\therefore\ p = \frac{r^2}{2a}$$ the equation required.

TRANSFORMATION OF CO-ORDINATES.

140. In some problems it is necessary to change the place of the origin, and the position of the axes, these cases will be separately treated.

(1) Let the origin be changed.



A the origin at first.

B the new origin.

P a point in the curve.

$\left.\begin{matrix} AC = \alpha \\ BC = \beta \end{matrix}\right\}$ the co-ordinates of B.

$$AN = x, \quad BM = x_1,$$
$$NP = y, \quad MP = y_1.$$

Then $x = x_1 + \alpha$, and $y = y_1 + \beta$.

Substitute these values for x and y, and the equation is transformed, and the co-ordinates are measured from B.

(2) Let the axes be changed but still rectangular.



Ax, Ay, the old axes, Ax_1, Ay_1, the new ones.

$$AN = x, \quad AM = x_1, \quad NP = y, \quad MP = y_1, \quad \angle xAx_1 = \theta,$$

Draw $Mm \perp$ to PN, and $Mn \perp$ to Ax;

$$\therefore x = An - Nn = x_1 \cos\theta - y_1 \sin\theta,$$

$y = Nm + Pm = x_1 \sin\theta + y_1 \cos\theta$. For $\angle mPM = \theta$.

(3) If the origin and position of the axes be changed. Let α and β be the co-ordinates of the origin, therefore we have

$$x = \alpha + x_1 \cos\theta - y_1 \sin\theta,$$
$$y = \beta + x_1 \sin\theta + y_1 \cos\theta.$$

141. To transform rectangular co-ordinates into polar, the origin being the pole.



$$AP = r, \quad \angle PAN = \theta;$$
$$\therefore x = r\cos\theta; \quad y = r\sin\theta,$$

which put for x and y in the equation.

But if the point S be the pole, draw $SB \perp Ax$, and $Sm \perp PN$.



$$AB = a, \qquad SP = r,$$

$$BS = \beta, \qquad \angle PSm = \theta;$$

$$\therefore\ x = AB + BN = a + r\cos\theta,$$

$$y = BS + Pm = \beta + r\sin\theta.$$

Ex. 1. Find the polar equation to the circle round a point S, co-ordinates α and β.

$$x^2 + y^2 = a^2;$$

$$\therefore\ (\alpha + r\cos\theta)^2 + (\beta + r\sin\theta)^2 = a^2;$$

$$\therefore\ r^2 + 2r.(\alpha\cos\theta + \beta\sin\theta) + \alpha^2 + \beta^2 - a^2 = 0.$$

Ex. 2. Transform $(x^2 + y^2)^2 = a^2(x^2 - y^2)$ into polar co-ordinates, the origin being the pole.

$$x^2 + y^2 = r^2, \text{ and } x = r\cos\theta,\ y = r\sin\theta;$$

$$\therefore\ r^4 = a^2r^2.(\cos^2\theta - \sin^2\theta);$$

$$\therefore\ r^2 = a^2\cos 2\theta.$$

Ex. 3. Transform the equation $x^2 - y^2 = a^2$ into another, the co-ordinates of which are rectangular, but the axis of y is inclined at an $\angle 45^0$ to the axis of x.

$$x = x_1\cos\theta - y_1\sin\theta,$$

$$y = x_1\sin\theta + x_1\cos\theta,$$

$$\theta = 2\pi - 45;\quad \therefore\ \cos\theta = \frac{1}{\sqrt{2}};\quad \sin\theta = -\frac{1}{\sqrt{2}};$$

$$\therefore\ x = \frac{x_1 + y_1}{\sqrt{2}};\quad y = \frac{x_1 - y_1}{\sqrt{2}};$$

$$\therefore\ x^2 - y^2 = \frac{(x_1 + y_1)^2 - (x_1 - y_1)^2}{2} = \frac{4x_1y_1}{2} = 2x_1y_1 = a^2;$$

$$\therefore\ x_1y_1 = \frac{a^2}{2}.$$

THE PARABOLA.

142. If from a fixed line QDq perpendicular lines, as QP, be drawn, intersecting lines equal in length, but drawn from a fixed point S, the locus of P is the *parabola*.



Draw $SD \perp Qq$, bisect SD in A, then the curve passes through A.

Let $SA = AD = a$, $AN = x$, $NP = y$.

Now QP or $DN = SP$; $\therefore DA + AN = \sqrt{NP^2 + SN^2}$;

$$\therefore a + x = \sqrt{y^2 + (x - a)^2};$$

$$\therefore (a + x)^2 \text{ or } (x - a)^2 + 4ax = y^2 + (x - a)^2;$$

$$\therefore y^2 = 4ax.$$

Cor. 1. If $x = a$; $y^2 = 4a^2$; $y = 2a$, let this value of y be $\frac{1}{2}l$; $\therefore l = 4a$, and $y^2 = lx$; l, which is the double ordinate through the focus, is called the Latus Rectum.

Cor. 2. The polar Equation. Let $SP = r$, $\angle ASP = \theta$.

Then $r = DN = 2a + SN = 2a + r\cos PSN$

$$= 2a - r\cos\theta;$$

$$\therefore r = \frac{2a}{1 + \cos\theta} = \frac{a}{\cos^2\frac{\theta}{2}}.$$

THE ELLIPSE.

143. If from two fixed points S and H two lines SP



and PH be drawn and intersect, and $SP + PH =$ a constant line, the locus of P is the *ellipse*.

Let $SP+PH=2a$. Bisect SH in C, and take $CA=CM=a$, the curve passes through A and M. Through C draw BCb $\perp$ to SH. With centre S and radius $=a$ cut this line in the points B and b, the curve will pass through B and b, since HB and Hb each $=a$; make $CB=b$, and let CS : CA :: e : 1; $\therefore$ $CS=ae$, which is called the eccentricity.

Let $CN=x$; $NP=y$; $SP=D$; $HP=D_1$;

$$\therefore D^2=SN^2+NP^2=(ae+x)^2+y^2,$$

$$D_1^2=HN^2+NP^2=(ae-x)^2+y^2;$$

$$\therefore D^2+D_1^2=2(a^2e^2+x^2+y^2), \text{ and } D^2-D_1^2=4aex.$$

But $D+D_1=2a$; $\therefore$ $D-D_1=2ex$;

$$\therefore D=a+ex, \text{ and } D_1=a-ex;$$

$$\therefore D^2+D_1^2=2a^2+2e^2x^2=2(a^2e^2+x^2+y^2);$$

$$\therefore y^2=a^2.(1-e^2)-x^2.(1-e^2)=(1-e^2)(a^2-x^2).$$

But $1-e^2=1-\frac{CS^2}{a^2}=\frac{a^2-CS^2}{a^2}=\frac{SB^2-CS^2}{a^2}=\frac{b^2}{a^2}$;

$$\therefore y^2=\frac{b^2}{a^2}(a^2-x^2); \text{ and } \frac{y^2}{b^2}+\frac{x^2}{a^2}=1.$$

Cor. 1. If A be the origin. Make $AN=x_1$;

$$\therefore x_1=a+x, \text{ or } x=x_1-a;$$

$$\therefore a^2-x^2=2ax_1-x_1^2;$$

$$\therefore y^2=\frac{b^2}{a^2}(2ax_1-x_1^2).$$

Cor. 2. If S be the pole, and $ASP=\theta$, and $SP=r$;

$$\therefore (2a-r)^2=HP^2=HN^2+NP^2=(2ae-SN)^2+r^2\sin^2\theta,$$

and $SN=r\cos PSH=-r\cos\theta$;

$$\therefore 4a^2-4ar+r^2=(2ae+r\cos\theta)^2+r^2\sin^2\theta$$

$$=4a^2e^2+4aer\cos\theta+r^2;$$

$$\therefore r=\frac{a(1-e^2)}{1+e\cos\theta}.$$

Cor. 3. If C be the pole, $CP=r$, and $PCM=\theta$.

Then $x=r\cos\theta$, and $y=r\sin\theta$;

$$\therefore \frac{x^2}{a^2} + \frac{y^2}{b^2} = r^2 . \left(\frac{\cos^2\theta}{a^2} + \frac{\sin^2\theta}{b^2}\right) = 1 ;$$

$$\therefore r = \frac{ab}{\sqrt{b^2\cos^2\theta + a^2\sin^2\theta}} = \frac{ab}{\sqrt{a^2(1-e^2)\cos^2\theta + a^2\sin^2\theta}}$$

$$= \frac{b}{\sqrt{1 - e^2\cos^2\theta}}.$$

THE HYPERBOLA.

144. If the difference between SP and PH be constant, the locus of P is the *hyperbola*.

Let the difference be $2a$; bisect SH in C. Take $CA = a = CM$, and the curve passes through A.



$\left.\begin{matrix} CN = x \\ NP = y \end{matrix}\right\}$. Let $CS = e \,.\, CA = ea$, where $e > 1$.

Then $HP^2 = HN^2 + NP^2 = (ea + x)^2 + y^2 = D_1^2$,

$$SP^2 = SN^2 + NP^2 = (ea - x)^2 + y^2 = D^2;$$

$$\therefore D_1^2 + D^2 = 2 \,.\, (a^2e^2 + x^2 + y^2); \quad D_1^2 - D^2 = 4aex.$$

Also $D_1 - D = 2a$; $\therefore D_1 + D = 2ex$;

$$\therefore D_1 = a + ex, \text{ and } D = ex - a;$$

$$\therefore 2a^2 + 2e^2x^2 = 2(a^2e^2 + x^2 + y^2);$$

$$\therefore y^2 = (e^2 - 1) \,.\, x^2 - (e^2 - 1)a^2 = (e^2 - 1)(x^2 - a^2)$$

$$= \frac{b^2}{a^2} \,.\, (x^2 - a^2); \quad \text{making } b^2 = a^2(e^2 - 1);$$

$$\therefore \frac{y^2}{b^2} - \frac{x^2}{a^2} = -1.$$

Cor. 1. If A be the origin, and $AN = x_1$; $\therefore x = x_1 + a$.

$$\therefore x + a = x_1 + 2a; \quad x^2 - a^2 = x_1^2 + 2ax_1;$$

$$\text{and } y^2 = \frac{b^2}{a^2}(2ax_1 + x_1^2).$$

Cor. 2. To find the polar equation, S being the pole.

$$SP = r, \quad \angle ASP = \theta.$$

Then $(2a + r)^2 = HP^2 = PN^2 + HN^2$

$$= PN^2 + (2CS - SN)^2$$

$$= r^2 \sin^2\theta + (2ae - r\cos\theta)^2;$$

$$\therefore\ 4a^2 + 4ar + r^2 = r^2 + 4a^2e^2 - 4aer\cos\theta;$$

$$\therefore\ r = \frac{a(e^2 - 1)}{1 + e\cos\theta}.$$

Cor. 3. Let C be the pole; $CP = r$; $\angle ACP = \theta$.

$$\therefore\ x = r\cos\theta, \text{ and } y = r\sin\theta;$$

$$\therefore\ \frac{x^2}{a^2} - \frac{y^2}{b^2} = r^2\left(\frac{\cos^2\theta}{a^2} - \frac{\sin^2\theta}{b^2}\right) = 1;$$

$$\therefore\ r = \frac{ab}{\sqrt{b^2\cos^2\theta - a^2\sin^2\theta}} = \frac{b}{\sqrt{e^2.\cos^2\theta - 1}}.$$

145. The asymptotes being the axes, and the centre the origin, find the equation to the hyperbola.



The asymptotes are lines, as CO and Co, drawn through the centre, making an angle $= \tan^{-1}\frac{b}{a}$ with the axis of the hyperbola.

$CN = x$, $CM = x_1$, and $OCA = oCA = \theta$,

$NP = y$, $MP = y_1$.

Draw $Mn \perp$ to CAN, and $Pm \perp$ to Mn.

Since MP is parallel to Co, and Pm to CN, $\therefore\ \angle MPm = \theta$.

Now $x = Cn + nN = x_1\cos\theta + y_1\cos\theta = (x_1 + y_1)\cos\theta$,

$$y = Mn - Mm = x_1\sin\theta - y_1\sin\theta = (x_1 - y_1)\sin\theta;$$

$$\therefore\ \frac{x^2}{a^2} - \frac{y^2}{b^2} = \frac{(x_1 + y_1)^2}{a^2}\cos^2\theta - \frac{(x_1 - y_1)^2}{b^2}\sin^2\theta = 1.$$

But $\tan\theta = \frac{b}{a}$; $\therefore\ 1 + \tan^2\theta = \frac{1}{\cos^2\theta} = \frac{b^2 + a^2}{a^2}$;

$$\therefore\ \frac{\cos^2\theta}{a^2} = \frac{1}{b^2 + a^2}, \text{ and } \frac{\sin^2\theta}{b^2} = \frac{\cos^2\theta}{a^2} = \frac{1}{b^2 + a^2};$$

$$\therefore\ \frac{(x_1 + y_1)^2 - (x_1 - y_1)^2}{b^2 + a^2} = 1;$$

$$\text{i.e. } 4x_1y_1 = a^2 + b^2, \quad x_1y_1 = \frac{a^2 + b^2}{4}.$$

Cor. If the hyperbola be rectangular, $b = a$, and $x_1y_1 = \frac{a^2}{2}$.

146. The curves whose equations have been just investigated are termed Conic Sections, since they may be supposed to arise from the intersection of a cone by a plane; and they are also called curves of the second degree, since the sum of the indices of x and y in any one term does not exceed two.

The general equation of the second degree is of the form

$$Ay^2 + Bxy + Cx^2 + Dy + Ex + F = 0.$$

Now if the centre be the origin, the equation to the curve ought to remain the same when $(-x)$ and $(-y)$ are put for x and y: consequently the origin of the co-ordinates of the general equation is not in the centre; since Dy and Ex will both change their signs, when $(-y)$ and $(-x)$ are put for y and x.

To get rid of these terms, transform the equation to the centre by putting $x + \alpha$ and $y + \beta$ for x and y, and make the coefficients of x and y respectively $= 0$; there will be two equations to determine α and β;

$$\text{and } \alpha = \frac{2AE - BD}{B^2 - 4AC}, \quad \beta = \frac{2CD - BE}{B^2 - 4AC}.$$

The equation now becomes $Ay^2 + Bxy + Cx^2 + F_1 = 0$.

Next, to get rid of the term Bxy; let the axes be changed to others, making an angle θ with the axis of x, by putting

$$x = x\cos\theta - y\sin\theta,$$

$$\text{and } y = x\sin\theta + y\cos\theta.$$

Therefore the coefficient of xy becomes

$$2A\sin\theta\cos\theta + B(\cos^2\theta - \sin^2\theta) - 2C\sin\theta\cos\theta = 0,$$

$$\text{or } (A - C)\sin 2\theta = -B\cos 2\theta; \quad \therefore \tan 2\theta = \frac{-B}{A - C},$$

an equation which is always possible, since the tangent passes through all degrees of magnitude from zero to infinity. The reduced equation finally becomes $My^2 + Nx^2 + F_1 = 0$, which may be made to coincide with the equations to the circle, the ellipse, or the hyperbola, by giving proper values to M, N, and F_1.

147. To find the relation between M, N, A, B and C, or to prove that $4MN = 4AC - B^2$.

By putting in the general equation $x\cos\theta - y\sin\theta$ for x, and $x\sin\theta + y\cos\theta$ for y; we have

$$M = A\cos^2\theta - B.\sin\theta\cos\theta + C.\sin^2\theta,$$

$$N = A\sin^2\theta + B.\sin\theta\cos\theta + C.\cos^2\theta;$$

$$\therefore\ M + N = A + C,$$

$$M - N = (A - C).\cos 2\theta - B.\sin 2\theta.$$

But since $\tan 2\theta = \dfrac{-B}{A-C}$,

$$\cos 2\theta = \frac{A-C}{\sqrt{(A-C)^2 + B^2}},\ \text{and}\ \sin 2\theta = \frac{-B}{\sqrt{(A-C)^2+B^2}};$$

$$\therefore\ M - N = \frac{(A-C)^2 + B^2}{\sqrt{(A-C)^2+B^2}} = \sqrt{(A-C)^2 + B^2};$$

$$\therefore\ 2M = A + C + \sqrt{(A-C)^2 + B^2},$$

$$2N = A + C - \sqrt{(A-C)^2 + B^2};$$

$$\therefore\ 4MN = (A+C)^2 - (A-C)^2 - B^2 = 4AC - B^2.$$

Whence, if $4AC > B^2$, M and N have the same sign;

if $4AC < B^2$, ... different signs;

if $4AC = B^2$, either M or N must $= 0$.

Cor. 1. If $B^2 = 4AC$, α and β are infinite, and the curve has not a centre: the first transformation is not therefore applicable, but the second is. Change the inclination of the axes, and the equation without the term Bxy becomes

$$My^2 + Nx^2 + Px + Ry + F = 0.$$

Now $4M.N = 4AC - B^2 = 0$; $\therefore$ M or $N = 0$.

Let $N = 0$; $\therefore$ $My^2 + Px + Ry + F = 0$.

Again, to get rid of the terms Ry and F, make

$$x = x + a,\quad \text{and}\quad y = y + b;$$

$$\therefore\ My^2 + (2Mb + R)\,y + Px + Pa + Rb + Mb^2 + F = 0;$$

and we find a and b, $\because$ $2Mb + R = 0$, or $b = \dfrac{-R}{2M}$,

and $Pa + Rb + Mb^2 + F = 0$; $\therefore\ a = \dfrac{4FM - R^2}{4MP}$;

and $\therefore\ My^2 + Px = 0$, the equation to the parabola.

If $M = 0$, we shall have $Nx^2 + Ry = 0$.

Cor. 2. If $My^2 + Nx^2 = F$ be an ellipse, find the axes.

Let a, b, be the semi-axes; $\therefore\ \dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1$;

but from the given equation $\dfrac{M}{F}y^2 + \dfrac{N}{F}x^2 = 1$;

$\therefore\ a^2 = \dfrac{F}{N}$; $b^2 = \dfrac{F}{M}$; $\therefore\ a = \sqrt{\dfrac{F}{N}}$; $b = \sqrt{\dfrac{F}{M}}$.

Ex. 1. Prove that the curve defined by $\sqrt{y} + \sqrt{x} = \sqrt{a}$, is a parabola.

Ex. 2. The equation $y^2 - 2yx + x^2 - 8x + 16 = 0$ belongs to a parabola, its axes are inclined at an angle of 45^0 to the axis of the parabola, and its equation when transformed to that axis is $(y + \sqrt{2})^2 = 2(x\sqrt{2} - 3)$.

Ex. 3. Find the centre and the axes of the curve

$$y^2 - 2xy + 3x^2 + 2y - 4x = 3.$$

Ex. 4. Find the centre of $y = \dfrac{x}{x + 1}$.

THE CISSOID.

148. AQB is a semicircle. Take AN and BM equal. Draw the ordinates NQ, MR. Join AR cutting NQ in P. The locus of P is the cissoid.



$$AN = x,\ NP = y,\ AB = 2a.$$

Now $\dfrac{AN^2}{NP^2} = \dfrac{AM^2}{MR^2} = \dfrac{AM^2}{AM \cdot MB} = \dfrac{AM}{MB}$,

or $\dfrac{x^2}{y^2} = \dfrac{2a - x}{x}$; $\therefore\ y^2 = \dfrac{x^3}{2a - x}$.

Cor. To find the Polar Equation.

$$AP = r, \quad \angle PAN = \theta, \quad x = r\cos\theta, \quad y = r\sin\theta,$$

$$\frac{y^2}{x^2} = \frac{\sin^2\theta}{\cos^2\theta} = \frac{x}{2a - x} = \frac{r\cos\theta}{2a - r\cos\theta};$$

$$\therefore\ 2a\sin^2\theta = r\cos\theta\,(\sin^2\theta + \cos^2\theta);$$

$$\therefore\ r = 2a\frac{\sin\theta}{\cos\theta}.\sin\theta = 2a\tan\theta.\sin\theta.$$

THE CONCHOID OF NICOMEDES.

149. The line CP revolves round a fixed point C, cutting the line ARN: RP is always of the same length; then the point P will trace out the conchoid.



Let $CA = a$, $AM = x$,

$RP = AB = b$, $MP = y$.

$$\frac{MP^2}{CM^2} = \frac{AR^2}{CA^2} = \frac{RN^2}{NP^2} = \frac{RP^2 - NP^2}{NP^2};$$

$$\text{or}\ \frac{y^2}{(a+x)^2} = \frac{b^2 - x^2}{x^2};$$

$$\therefore\ x^2y^2 = (a+x)^2(b^2 - x^2).$$

Cor. Let $CP = r$, $\angle PCM = \theta$,

$$r = CP = PR + CR = a + \frac{b}{\cos\theta}.$$

THE WITCH.

150. AQB is a semi-circle, and NP is taken a fourth proportional to AN, AB, and NQ,

Then the locus of P is the Witch.



$AN = x$, $AB = 2a$,

$NP = y$; $\therefore\ NQ = \sqrt{2ax - x^2}$,

$x : 2a :: \sqrt{2ax - x^2} : y$;

$$\therefore\ y = \frac{2a\sqrt{2ax - x^2}}{x} = 2a\sqrt{\frac{2a - x}{x}}.$$

151. The Logarithmic Curve.

In this curve, the abscissa is the logarithm of the ordinate, or if a be the base of the system, the equation to the curve is $y = a^x$;



$$\therefore\ AB = a^0 = 1,$$

or the ordinate through the origin is always unity.

It is obvious that as the abscissa increases in arithmetic, the ordinate increases in geometric progression.

152. The Quadratrix of Dinostratus.

While the ordinate RN of the quadrant AQB moves uniformly from A to BC, the radius revolves from CA to CB, cutting RN in P; the locus of P is the curve required.



$$AN = x, \qquad CB = a,$$
$$NP = y, \qquad \angle QCA = \theta.$$

$$\text{Then } \theta : \frac{\pi}{2} :: x : a; \quad \therefore\ \theta = \frac{\pi x}{2a};$$

$$\frac{PN}{CN} = \frac{y}{a - x} = \tan\theta = \tan\frac{\pi x}{2a};$$

$$y = (a - x) \cdot \tan\frac{\pi x}{2a}.$$

Cor. When $x = a$; $y = Cb = \dfrac{2a}{\pi}$.

153. If RN move as before, and a line as QPM parallel to AC move uniformly from AC, the intersection P of RN and QM will trace the Quadratrix of Tschirnhausen.



Here $AQ : AQB :: x : a$;

$$\therefore\ AQ = \frac{x}{a} \cdot \frac{\pi a}{2} = \frac{\pi x}{2};$$

$$\therefore\ y = a \cdot \sin\left(\frac{AQ}{a}\right) = a \cdot \sin\frac{\pi x}{2a} \text{ is the equation.}$$

THE LEMNISCATA.

154. If SH be a straight line bisected in C, and if SP and HP revolve round S and H, and intersect in P, so that $SP \times HP = CS^2$, the locus of P is the Lemniscata.



$$CN = x; \quad NP = y; \quad CS = a,$$

$$SP = \sqrt{PN^2 + SN^2} = \sqrt{y^2 + (a+x)^2},$$

$$HP = \sqrt{PN^2 + HN^2} = \sqrt{y^2 + (a-x)^2};$$

$$\therefore \sqrt{y^2 + (a+x)^2} \times \sqrt{y^2 + (a-x)^2} = a^2;$$

$$\therefore \{(y^2 + a^2 + x^2) + 2ax\}\{(y^2 + a^2 + x^2) - 2ax\} = a^4;$$

$$\therefore (y^2 + a^2 + x^2)^2 - 4a^2x^2 = a^4;$$

$$\therefore y^4 + 2a^2y^2 + x^4 = 2a^2x^2 - 2a^2y^2;$$

$$\therefore (y^2 + x^2)^2 = 2a^2(x^2 - y^2).$$

Cor. If $CP = r$; and $\angle PCH = \theta$.

Then $x = r\cos\theta$; $y = r\sin\theta$; $x^2 + y^2 = r^2$;

$$\therefore r^4 = 2a^2r^2(\cos^2\theta - \sin^2\theta); \quad \therefore r^2 = 2a^2\cos 2\theta,$$

which is the Polar Equation.

THE CYCLOID.

155. *The Cycloid* is described by a point in the circumference of a circle, which rolls along a horizontal line.



Let BQD be the circle, O the centre; and when its diameter is perpendicular to the horizontal line at A, let the point P, which generates the curve, also be at A.

Then Ab must = arc Pb, since each point of Pb has been in contact with each successive point of Ab.

(1°) Let $AN = x$, $BD = 2a$,

$NP = y$, $\angle QOB = \theta$;

$\therefore\ x = Ab - Nb = a\theta - a \sin\theta = a(\theta - \sin\theta)$;

$\therefore\ y = bm = a \text{ ver. } \sin\theta = a(1 - \cos\theta)$;

an algebraic equation cannot be found between x and y, but since $x = a \text{ ver. } \sin^{-1}\frac{y}{a} - \sqrt{2ay - y^2}$;

$$\therefore\ \frac{dx}{dy} = \frac{a}{\sqrt{2ay - y^2}} - \frac{a - y}{\sqrt{2ay - y^2}} = \frac{y}{\sqrt{2ay - y^2}},$$

a differential equation.

(2°) To find the equation from the vertex D.

Let $DM = x$; $MP = y$; $\angle DOQ = \theta$.

Join Pb and QB, then these being equal and parallel,

$$PQ = Bb = AB - Ab = AB - Pb = DQ.$$

Then $y = PM = MQ + PQ = a \sin\theta + a\theta = a(\theta + \sin\theta)$,

$x = DM = a \text{ ver. } \sin\theta = a(1 - \cos\theta)$.

Cor. Since $x = a \text{ ver. } \sin\theta$; $\therefore\ \theta = \text{ver. } \sin^{-1}\frac{x}{a}$, and

$$a \sin\theta = \sqrt{2ax - x^2};$$

$$\therefore\ y = \sqrt{2ax - x^2} + a \text{ ver. } \sin^{-1}\frac{x}{a};$$

$$\therefore\ \frac{dy}{dx} = \frac{a - x}{\sqrt{2ax - x^2}} + \frac{a}{\sqrt{2ax - x^2}}$$

$$= \frac{2a - x}{\sqrt{2ax - x^2}} = \frac{\sqrt{2ax - x^2}}{x},$$

which is the equation most commonly used.

156. From (2°) may be derived a mechanical method of describing the cycloid; for the point P is found by drawing MP perpendicular to DB, and equal to the sum of the ordinate QM and the arc DQ of the circle.

157. If we take MP equal to DQ only, then the locus of

P is a curve called the Companion to the Cycloid: its equations therefore are

$$y = a\theta; \quad x = a(1 - \cos\theta),$$

whence $\frac{dy}{dx} = \frac{a}{\sqrt{2ax - x^2}}$.

THE TROCHOID.

158. The trochoid is the curve traced in space by a point B in the circumference of the inner circle BRb, whilst the outer circle AQ rolls upon a horizontal line.

P a point in the trochoid. Through P draw a horizontal



line $MRPm$. Take O and o the centres of the circles.

Draw ORQ and oP.

Then $Pm = RM$, and $\angle AOQ = \angle A_1oP$.

Let $\left.\begin{matrix} OA = a, & AN = x \\ OB = b, & NP = y \end{matrix}\right\}$, $\angle AOR = \theta$.

Then it is obvious that arc $AQ = AA_1$;

$$\therefore x = AA_1 - NA_1 = a\theta - b\sin\theta,$$

$$y = NP = oA_1 + om = a - b\cos\theta.$$

Let $e = \frac{b}{a}$; $\quad \therefore x = a(\theta - e\sin\theta)$; $\quad y = a(1 - e\cos\theta)$.

If $e = 1$, or $b = a$, the trochoid becomes the common cycloid and their equations coincide.

THE EPICYCLOID AND HYPOCYCLOID.

159. If one circle revolve upon another circle and in the same plane with it, the curve described by any point in the circumference of the revolving circle is called the Epicycloid; but if the revolving circle move within the other circle, the curve described by the point is called the Hypocycloid.





P the describing point of epicycloid.

$$CA = a; \quad CB = b; \quad \angle ACQ = \theta;$$

$$CN = x; \quad NP = y; \quad \angle PBQ = \phi;$$

$$\therefore\ x = Cn + Nn = (a + b)\cos\theta + b \sin PBm.$$

But $PBm = PBQ - (90 - \theta) = \phi + \theta - 90.$

Now $AQ = PQ; \quad \therefore\ a\theta = b\phi; \quad \therefore\ \phi = \frac{a\theta}{b};$

$$\therefore\ x = (a + b)\cos\theta - b\cos\left(\frac{a+b}{b}.\theta\right);$$

$$y = Bn - Bm = (a + b)\sin\theta - b\sin\left(\frac{a+b}{b}.\theta\right).$$

160. If b be negative, the curve is the hypocycloid.

$$\therefore\ x = (a - b)\cos\theta + b\cos\left(\frac{a-b}{b}.\theta\right),$$

$$\text{and } y = (a - b)\sin\theta - b\sin\left(\frac{a-b}{b}.\theta\right),$$

or we may derive it from the figure.

$$\text{For } PBm = PBC - nBC = 180 - \frac{a\theta}{b} - (90 - \theta)$$

$$= 90 - \frac{a-b}{b}.\theta,$$

$$\text{and } x = (a - b)\cos\theta + b\sin PBm$$

$$= (a - b)\cos\theta + b\cos\left(\frac{a-b}{b}.\theta\right),$$

$$y = (a - b)\sin\theta - b\cos PBm$$

$$= (a - b)\sin\theta - b\sin\left(\frac{a-b}{b}.\theta\right).$$

161. If the describing point P be not in the circumference, but within the revolving circle and at a distance b_1 from the centre. The curves described are then respectively called the Epitrochoid and Hypotrochoid: and their equations are

$$\left.\begin{aligned} x &= (a + b)\cos\theta - b_1\cos\left(\frac{a+b}{b}.\theta\right) \\ y &= (a + b)\sin\theta - b_1\sin\left(\frac{a+b}{b}.\theta\right) \end{aligned}\right\} \ldots\ldots (1).$$

$$\left.\begin{aligned} x &= (a-b)\cos\theta + b_1 \cos\left(\frac{a-b}{b}.\theta\right) \\ y &= (a-b)\sin\theta - b_1 \sin\left(\frac{a-b}{b}.\theta\right) \end{aligned}\right\} \ldots\ldots(2).$$

162. If in the epicycloid $a = b$,

$$x = a(2\cos\theta - \cos 2\theta),$$

$$y = a(2\sin\theta - \sin 2\theta);$$

$\therefore$ squaring and adding,

$$x^2 + y^2 = a^2(4 - 4\cos\theta + 1);$$

$$\therefore\ x^2 + y^2 - a^2 = 4a^2(1 - \cos\theta).$$

But $x = a(2\cos\theta - 2\cos^2\theta + 1);$

$$\therefore\ x - a = 2a\cos\theta(1 - \cos\theta);$$

$$y = 2a\sin\theta(1 - \cos\theta);$$

$$\therefore\ (x-a)^2 + y^2 = 4a^2(1 - \cos\theta)^2.$$

But $16a^2(1 - \cos\theta)^2 = (x^2 + y^2 - a^2)^2;$

$$\therefore\ 4a^2\{y^2 + (x-a)^2\} = (x^2 + y^2 - a^2)^2;$$

the equation to rectangular co-ordinates.

Cor. If $x - a = r\cos\phi$, and $y = r\sin\phi$,

$$4a^2r^2 = (r^2 + 2ar\cos\phi)^2;$$

$$\therefore\ r = 2a(1 - \cos\phi);$$

the polar equation to the curve, called from its form the Cardioid.

163. To find the equation to the epicycloid in terms of the radius vector CP and the perpendicular on the tangent CY.

Produce CB to q; join PQ, Pq, then since for an instant the revolving circle turns on Q, the motion of the point P must be perpendicular to QP, and therefore in the direction Pq: hence qP produced is a tangent to the curve.

Produce qP, and draw $CY \perp$ to it, and make

$CP = r$; $CY = p$; then $\because$ CY is parallel to PQ,

$$\frac{PY^2}{CQ^2} = \frac{qY^2}{Cq^2};$$ also, $Cq = a + 2b$;

$$\therefore \frac{r^2 - p^2}{a^2} = \frac{(a+2b)^2 - p^2}{(a+2b)^2} = \frac{c^2 - p^2}{c^2}; \text{ if } c = a + 2b;$$

$$\therefore r^2c^2 - p^2c^2 = a^2c^2 - a^2p^2;$$

$$\therefore p^2 = c^2 \cdot \left(\frac{r^2 - a^2}{c^2 - a^2}\right).$$

Cor. For the hypocycloid, $c = a - 2b$ is $< a$;

$$\therefore p^2 = a^2 \cdot \left(\frac{a^2 - r^2}{a^2 - c^2}\right).$$

---

164. SPIRALS.

(1) The spiral of Archimedes. In this spiral the radius vector varies directly as the angle described,

or $r \propto \theta$; $\therefore r = a\theta$ is the equation.

Its equation may be thus found. Let a line revolve uniformly round S, while a point P moves uniformly from S along it, then P will trace the spiral of Archimedes.



Or let $\angle ASP = \theta$, $SP = r$; and let c be the value of r when $\theta = 2\pi$;

$$\therefore r : c :: \theta : 2\pi;$$

$$\therefore r = \frac{c}{2\pi}\theta = a\theta \text{ by putting } a = \frac{c}{2\pi}.$$

(2) The logarithmic spiral. Here the angle described is the logarithm of the radius vector; its equation is $r = a^\theta$.

This curve is also called the equiangular spiral, since, as it will be shewn, the angle at which it cuts the radius is constant.

(3) The hyperbolic spiral. In this spiral as the angle increases the radius vector decreases, and its equation is $r = \frac{a}{\theta}$.

(4) The *lituus*, so called from its form; where $\theta = \frac{a^2}{r^2}$.

(5) The spiral of Archimedes, the hyperbolic, and the lituus, are included under the general equation $r = a\theta^n$, as we shall see by putting $n = 1$, -1, or $\frac{-1}{2}$.

THE INVOLUTE OF THE CIRCLE.

165. This curve is described by the extremity of a string which is unwound from the circumference of a circle.



A the point from which the string began to be unwrapped, QP the string once coincident with, and therefore $= AQ$; PY a tangent to AP, SY perpendicular to PY, join SP.

$$SP = r, \quad SY = p, \quad SQ = a;$$

$$\therefore\ SQ = PY = \sqrt{SP^2 - SY^2};$$

$$\therefore\ a^2 = r^2 - p^2; \quad \therefore\ p^2 = r^2 - a^2, \text{ is the equation.}$$

Cor. If $\phi = \sec^{-1}\frac{r}{a} = PSQ$, and $\theta = \angle ASP$,

$$\theta + \phi = \frac{\sqrt{r^2 - a^2}}{a}; \quad \therefore\ \theta = \frac{\sqrt{r^2 - a^2}}{a} - \sec^{-1}\frac{r}{a};$$

$$\therefore\ a\frac{d\theta}{dr} = \frac{\sqrt{r^2 - a^2}}{r} = \frac{SY}{SP}.$$

CHAPTER XI.

TANGENTS TO CURVES.

166. DEF. A TANGENT to a curve is a straight line which has a point in common with the curve, but which, if produced, does not cut the curve in the neighbourhood of the point.

Let PP_1 be the curve, and suppose a line to pass through the points P and P_1, and then to revolve round P, so as to make P_1 continually approach to P; the position of the revolving line, when P_1 coincides with P, will be that of the tangent.



Let $AN = x$, Ax, Ay the axes of x and y.

$NP = y$, QPT the last position of the line.

$y = f(x)$ the equation to the curve.

$y_1 = ax_1 + b$ that of the line through P and P'.

$NN_1 = h$; $N_1P_1 = y'$; $AN_1 = x' = x + h$.

Then because the line passes through P and P_1,

$$y_1 - y = \frac{y' - y}{x' - x} \cdot (x_1 - x) \text{ is its equation.}$$

But $y' = f(x + h) = y + \frac{dy}{dx}h + Ph^2$;

$$\therefore \frac{y' - y}{x' - x} = \frac{\frac{dy}{dx}h + Ph^2}{h} = \frac{dy}{dx} + Ph.$$

Now since by the revolution of the line (supposed to pass through P_1 and P), P_1 approaches P; therefore h decreases,

and the right-hand member of the equation continually approaches $\frac{dy}{dx}$, and when P_1 actually coincides with P, $h = 0$, and $\frac{y'-y}{x'-x} = \frac{dy}{dx}$; but then the line becomes a tangent,

and $y_1 - y = \frac{dy}{dx}(x_1 - x)$ is the equation required.

Cor. 1. The equation to the tangent may be written

$$y_1 = \frac{dy}{dx}x_1 + y - x\frac{dy}{dx},$$

which compared with the general equation to the straight line, $y = mx + b$, gives

$$m = \frac{dy}{dx}; \quad b = y - x\frac{dy}{dx},$$

which shew that the tangent is a line inclined to the axis of x at an angle whose tangent is $\frac{dy}{dx}$, and which cuts off from the axis of y a line $= y - x\frac{dy}{dx}$; from the former circumstance it is that the Differential Calculus has been called the Method of Tangents.

Cor. 2. Hence $AD = y - x\frac{dy}{dx}$, and

$$\because \frac{AD}{AT} = \tan DTA = \frac{dy}{dx}; \quad \therefore AT = y\frac{dx}{dy} - x.$$

Cor. 3. These values may be derived from the general equation; for

$$\text{let } x_1 = 0; \qquad \therefore y_1 = AD = y - x\frac{dy}{dx},$$

$$y_1 = 0; \quad \therefore -x_1 = AT = y\frac{dx}{dy} - x;$$

or if these values be called y_0 and x_0, we have

$$y_0 = y - x\frac{dy}{dx}; \quad x_0 = x - y\frac{dx}{dy},$$

which are the parts cut off from the axes of y and x by the tangent.

Hence, having found $\frac{dy}{dx}$ from the given equation to the curve, AD and AT may be found; then join TD, it produced is the tangent.

Cor. 4. If the axes be oblique, we shall obtain similar results, but $\frac{dy}{dx}$ will not be the tangent of PTN, but the ratio of the sines of the angles the tangent makes with the axes of x and y.

Cor. 5. The line NT is called the subtangent, and is useful in drawing the tangent,

$$\text{and } NT = AN + AT = x + y\frac{dx}{dy} - x = y\frac{dx}{dy}.$$

Hence to draw a tangent, find the value of NT. Join P, T, and we have the tangent required.

Cor. 6. The length PT of the tangent

$$= \sqrt{PN^2 + NT^2} = \sqrt{y^2 + y^2\frac{dx^2}{dy^2}} = y\sqrt{1 + \frac{dx^2}{dy^2}}.$$

167. Def. A line PG drawn from the point of contact P, perpendicularly to the tangent, and meeting the axis in G, is called the normal.

Since if $y = mx + b$ is the equation to a line;

$$\therefore y = -\frac{1}{m}x + b_1 \text{ is that of a perpendicular to it;}$$

$$\therefore \text{since } y_1 = \frac{dy}{dx}x_1 + y - x\frac{dy}{dx} \text{ is equation to tangent;}$$

$$\therefore y_1 = -\frac{dx}{dy}x_1 + b_1 \text{ is that of the normal;}$$

$$\therefore y = -\frac{dx}{dy}x + b_1 \text{ since it passes through } P;$$

$$\therefore y_1 - y = -\frac{dx}{dy}(x_1 - x) \text{ is the equation required;}$$

and $\frac{dx}{dy}$ being found from the given equation to the curve, the normal may be drawn.

Cor. 1. Hence $\tan PGx = -\frac{dx}{dy}$; and $\therefore \tan PGN = \frac{dx}{dy}$;

also if $y_1 = 0$; $x_1 = AG = y\frac{dy}{dx} + x$;

$\therefore$ AG, called the subnormal $= AG - AN = y\frac{dy}{dx}$.

But AG may also be found from similar triangles NTP, PGN.

For $NG = \frac{NP^2}{NT} = \frac{y^2}{y \cdot \frac{dx}{dy}} = y\frac{dy}{dx}$.

Hence to draw a normal, find NG; join PG, it is the normal.

Cor. 2. The length PG of the normal

$$= \sqrt{PN^2 + NG^2} = \sqrt{y^2 + y^2\frac{dy^2}{dx^2}} = y\sqrt{1 + \frac{dy^2}{dx^2}}.$$

168. The normal is the shortest or longest line that can be drawn from a given point to a curve.

For if x_1 and y_1 be the co-ordinates of the given point,

x and y of a point in the curve,

u = distance between the points;

$\therefore u^2 = (x_1 - x)^2 + (y_1 - y)^2$ a maximum or minimum;

$$\therefore u\frac{du}{dx} = 0 = (x_1 - x) + (y_1 - y)\frac{dy}{dx} = 0;$$

$$\therefore y_1 - y = -\frac{dx}{dy}(x_1 - x),$$

which is, as we have just seen, the equation to the normal.

169. Find the length of the perpendicular from the origin upon the tangent, and the angle which the line from the origin to the point of contact makes with the tangent.

Since if δ be the perpendicular from the origin on a line

$$y = mx + b; \quad \delta = \frac{b}{\sqrt{m^2 + 1}} = \frac{y - mx}{\sqrt{m^2 + 1}};$$

$$\therefore \text{ if } m = \frac{dy}{dx}; \quad \delta = \frac{y - x\dfrac{dy}{dx}}{\sqrt{1 + \dfrac{dx^2}{dy^2}}} = \frac{y - px}{\sqrt{1 + p^2}}.$$

Again, if θ be the angle between the tangent and a line from the origin,

$$\theta = \tan^{-1}\frac{y}{x} - \tan^{-1}\frac{dy}{dx};$$

$$\therefore \tan\theta = \frac{\dfrac{y}{x} - \dfrac{dy}{dx}}{1 + \dfrac{y\,dy}{x\,dx}} = \frac{y - px}{x + py}.$$

Cor. If we use differentials, multiply by dx.

$$\delta = \frac{y\,dx - x\,dy}{\sqrt{dx^2 + dy^2}},$$

$$\text{and } \tan\theta = \frac{y\,dx - x\,dy}{x\,dx + y\,dy}.$$

170. It may be useful to collect these values in a table.

(1) $y_1 - y = \dfrac{dy}{dx}(x_1 - x)$, equation to tangent.

(2) $y_1 - y = -\dfrac{dx}{dy}(x_1 - x)$, equation to normal.

(3) Subtangent $NT = y\dfrac{dx}{dy}$.

(4) Subnormal $NG = y\dfrac{dy}{dx}$.

(5) Tangent $PT = y\sqrt{1 + \dfrac{dx^2}{dy^2}}$.

(6) Normal $PG = y\sqrt{1 + \dfrac{dy^2}{dx^2}}$.

(7) Perpendicular on tangent $= \dfrac{y - px}{\sqrt{1 + p^2}}$; where $p = \dfrac{dy}{dx}$.

(8) Tangent $APT = \dfrac{y - px}{x + py}$.

(9) $AD = y_0 = y - x\dfrac{dy}{dx}$.

(10) $AT = -x_0 = y\dfrac{dx}{dy} - x$;

of these formulas, the first four are the most important to be remembered.

171. Since the tangent of the angle which the tangent makes with the axis of x is $\dfrac{dy}{dx}$; the angle at which the curve cuts the axis may be found.

For the angle which the tangent makes with the axis at the point of section will be the same that the curve makes. Find therefore the co-ordinates of the point of section, and substitute them in the expression for $\dfrac{dy}{dx}$, and the resulting value will be the tangent of the angle required.

Ex. 1. Let $y = \dfrac{x}{1+x}$ be the equation to the curve.

Here if $x = 0$, $y = 0$; and $\therefore$ the origin is the point of section,

$$\text{and } \frac{dy}{dx} = \frac{1}{(1+x)^2} = \frac{1}{1}, \text{ when } x = 0;$$

$$\therefore \tan\theta = 1 = \tan 45^0; \quad \therefore \theta = 45^0.$$

Ex. 2. Let the curve be the cycloid.

$$\text{Here } \frac{dy}{dx} = \frac{\sqrt{2a-y}}{\sqrt{y}} = \sqrt{\frac{2a}{y} - 1},$$

which is infinite if $y = 0$; or at the origin the curve cuts the axis of x at an angle of 90^0.

Ex. 3. Let the curve be the circle, and the origin in the centre.

$$y^2 = a^2 - x^2, \quad \frac{dy}{dx} = p = -\frac{x}{y}; \quad 1 + p^2 = \frac{y^2 + x^2}{y^2} = \frac{a^2}{y^2},$$

$$y - px = y + \frac{x^2}{y} = \frac{a^2}{y}, \quad py + x = -x + x = 0;$$

$$\therefore AY = \frac{\frac{a^2}{y}}{\frac{a}{y}} = a; \quad \tan APT = \frac{\frac{a^2}{y}}{0} = \infty; \quad \therefore \angle APT = 90^0.$$

172. To draw a tangent through a given point.

Let α and β be the co-ordinates of the given point;

x and y be the co-ordinates of the curve;

$(y_1 - y) = \frac{dy}{dx}(x_1 - x)$ is the equation to the tangent.

But because it passes through a point where $y_1 = \beta$ and $x_1 = \alpha$;

$$\therefore \beta - y = \frac{dy}{dx}(\alpha - x);$$

from which, and the given equation to the curve, the point to which the tangent is to be drawn may be found.

173. To draw a tangent parallel to a given line.

Let A = tangent of the angle which the given line makes with the axis x;

$$\therefore \frac{dy}{dx} = A, \text{ since tangent and line are parallel;}$$

and $y_1 - y = A.(x_1 - x)$ is the equation required.

If it pass through a given point, the co-ordinates of the point may be put for x_1 and y_1, and then from the given equation to the curve, and from that of the tangent, the point to which the tangent is to be drawn may be found.

174. To find the locus of the intersections of perpendiculars drawn from the origin upon the tangent, with the tangent.

Let $y = f(x)$ be the equation to the curve;

$$\therefore y_1 - y = \frac{dy}{dx}(x_1 - x) \text{ is the equation to the tangent.}$$

$$y_1 = -\frac{dx}{dy}x_1 \text{ is that of perpendicular from origin.}$$

From these three equations eliminate y, x; and also $\frac{dy}{dx} = f'(x)$; the resulting equation, containing y_1, x_1, and constant quantities, will be that of the curve required.

PROB. If δ be the length of the perpendicular from the origin upon the tangent, shew that $\delta = \sqrt{x_1 x + y y_1}$.

For $\because y_1 - y = \frac{dy}{dx}(x_1 - x)$.....(1),

$$\text{and } y_1 = -\frac{dx}{dy}x_1 \ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots (2),$$

are simultaneous equations for the point of intersection;

$$\therefore (1) \times (2),\quad y_1^2 - yy_1 = -(x_1^2 - xx_1);$$

$$\therefore y_1^2 + x_1^2 = \delta^2 = yy_1 + xx_1;$$

$$\therefore \delta = \sqrt{yy_1 + xx_1}.$$

175. Two curves whose equations are $y = f(x)$; and $y' = \phi(x')$ intersect, find the angle of intersection.

Let θ be the angle between the tangents at the point of intersection, it will therefore be the angle required;

$$\text{also } \theta = \tan^{-1}\frac{dy}{dx} - \tan^{-1}\frac{dy'}{dx'}.$$

$$\text{But } \frac{dy}{dx} = f'(x);\quad \frac{dy'}{dx'} = \phi'(x') = \phi'(x).$$

Since at point of intersection $x' = x$;

$$\therefore \tan\theta = \frac{f'(x) - \phi'(x)}{1 + f'(x).\phi'(x)}.$$

PROB. The vertex of a parabola is in the centre of a circle, and its focus bisects the radius, find the angle of intersection of circle and parabola.

Here $y^2 = 2ax$ (1), and $y^2 = a^2 - x^2$ (2), are the equations to the parabola and the circle.

$$\text{From (1) } \frac{dy}{dx} = f'(x) = \frac{a}{y};\quad (2)\ \frac{dy}{dx} = \phi'(x) = -\frac{x}{y};$$

$$\therefore \tan\theta = \frac{\dfrac{a+x}{y}}{1 - \dfrac{ax}{y^2}} = \frac{y(a+x)}{y^2 - ax} = \frac{y(a+x)}{ax}.$$

But $2ax = a^2 - x^2$ at point of intersection;

$$\therefore x + a = a\sqrt{2};\quad x = a(\sqrt{2} - 1);\quad y = a\sqrt{2}\sqrt{\sqrt{2} - 1};$$

$$\therefore \tan\theta = \frac{2\sqrt{\sqrt{2} - 1}}{\sqrt{2} - 1} = \frac{2}{\sqrt{\sqrt{2} - 1}}.$$

ASYMPTOTES.

176. Asymptotes are tangents to the curve at a point infinitely distant from the origin, and may be drawn, if the values of AD or AT, or of both remain finite, when either x or y, or x and y, are infinite.

Asymptotes may be thus constructed:

(1) If AD and AT be finite, join T, D, and the line produced is the asymptote.

(2) If AD be infinite, and AT finite, the asymptote is perpendicular to the axis of x, passing through T.

(3) If AD be infinite, and $AT = 0$, the asymptote coincides with the axis of y.

(4) If AD be finite, and AT infinite, the asymptote is parallel to the axis of x; and if $AD = 0$ is coincident with it.

EXAMPLE. Draw an asymptote to the hyperbola.

Here $y = \frac{b}{a}\sqrt{2ax + x^2}$, and $\frac{dy}{dx} = \frac{b}{a}\frac{a+x}{\sqrt{2ax+x^2}}$,

$$AD = y - x\frac{dy}{dx} = \frac{b}{a}\cdot\left\{\sqrt{2ax+x^2} - \frac{(a+x)x}{\sqrt{2ax+x^2}}\right\} = \frac{bx}{\sqrt{2ax+x^2}}$$

$$= \frac{b}{\sqrt{1+\frac{2a}{x}}} = b \text{ if } x = \infty;$$

$$AT = y\frac{dx}{dy} - x = \frac{2ax+x^2}{a+x} - x = \frac{ax}{a+x} = \frac{a}{1+\frac{a}{x}} = a; \text{ if } x = \infty;$$

$\therefore$ $AT = \frac{1}{2}$ major-axis, or T and C coincide.

Join CD; it produced, is the asymptote.

177. This method is frequently difficult of application, and the following is more generally useful.

If possible, let the equation to the curve be put under the form $y = Ax + B + \frac{C}{x} + \frac{D}{x^2} + \frac{E}{x^3} + \&c.$ then it is obvious, that as x increases, the terms after B decrease; and when x becomes infinitely great, they vanish, and the equation to the infinite branch of the curve is $y = Ax + B$.

But this is the equation to a straight line cutting the axis of y at a point $y = B$, and making an angle $= \tan^{-1} A$,

with the axis of x. Hence it appears that the infinite branch of the curve is coincident with the line determined by the equation $y = Ax + B$;

$\therefore$ if $y = Ax + B + \frac{C}{x} + \frac{D}{x^2}$ + &c. be the equation to a curve,

$y = Ax + B$ is the equation to the asymptote.

We may observe that this method will not apply to find the asymptotes that are parallel to the axis of y: since then A would be infinite: but asymptotes of this kind are discoverable by simple inspection of the equation. (See Example 10.)

Cor. 1. If the form of the expanded $f(x)$ be

$$y = Ax^2 + Bx + C + \frac{D}{x} + \frac{E}{x^2} + \&c.$$

the asymptote is a parabolic curve, of which the equation is

$$y = Ax^2 + Bx + C.$$

Cor. 2. Since for every finite value of x, the ordinate of the curve $y = Ax + B + \frac{C}{x}$: differs less from that of the original curve, than the ordinate of the rectilinear asymptote, it is obvious that we may have a hyperbolic curve lying between the asymptote and the curve, and ultimately coincident with either; this may be called a hyperbolic asymptote.

EXAMPLES.

THE PARABOLA.

(1) Here $y^2 = 4ax$;

$\therefore \frac{dy}{dx} = \frac{2a}{y}$;

$\therefore y\frac{dx}{dy} = NT = \frac{y^2}{2a} = 2x$;

$\therefore AT = AN$.



Take $\therefore AT = AN$, join PT, PT is the tangent,

$$y\frac{dy}{dx} = 2a; \text{ or } NG = 2AS.$$

Take $\therefore NG = 2AS$; join PG, it is the normal;

equation to tangent is $y_1 - y = \frac{2a}{y}(x_1 - x)$;

$$\therefore\ yy_1 - y^2 = 2ax_1 - 2ax;\quad \therefore\ yy_1 = 2a(x_1 + x).$$

Cor. Since $yy_1 = 2a(x_1 + x)$ is equation to tangent;

$$\therefore\ y_1 = \frac{2a}{y}x_1 + \frac{2ax}{y};\ \text{make}\ \frac{2a}{y} = m;$$

$$\therefore\ \frac{2ax}{y}\ \text{or}\ \frac{y}{2} = \frac{a}{m};$$

$$\therefore\ y_1 = mx_1 + \frac{a}{m};$$

an equation to the tangent which is often very convenient in the solution of problems.

The equation to the normal is

$$y_1 - y = -\frac{y}{2a}(x_1 - x);$$

$$\therefore\ y_1 = -\frac{y}{2a}x_1 + y + \frac{yx}{2a}.$$

Make $\frac{-y}{2a} = m_1;\quad \therefore\ \frac{y^2}{4a^2} = \frac{x}{a} = m_1^2;$

$$\therefore\ y + \frac{yx}{2a} = -2am_1 - am_1^3;$$

$$\therefore\ y_1 = m_1(x_1 - 2a - am_1^2),$$

the equation to the normal in terms of its inclination to the axis of x.

Ex. Two normals to a parabola intersect at right angles; find the locus of their intersection.

$$\therefore\ y + 2am = m(x - am^2) \ldots\ldots\ldots\ldots\ldots\ldots (1),$$

$$\text{and}\ y - \frac{2a}{m} = -\frac{1}{m}\left(x - \frac{a}{m^2}\right) \ldots\ldots\ldots\ldots\ldots\ldots (2);$$

$$\therefore\ 2a\left(m + \frac{1}{m}\right) = x\left(m + \frac{1}{m}\right) - a\left(m^3 + \frac{1}{m^3}\right);$$

$$\therefore\ x - 3a = a\left(m - \frac{1}{m}\right)^2 \ldots\ldots\ldots\ldots (3).$$

But from (2) $my - 2a = -x + \frac{a}{m^2}$,

(1) $\frac{1}{m}y + 2a = x - am^2$;

$$\therefore\ y\left(m+\frac{1}{m}\right)=a\left(\frac{1}{m^2}-m^2\right);$$

$$\therefore\ y=a\left(\frac{1}{m}-m\right)\ldots\ldots\ldots\ldots\ldots\ldots(4);$$

$$\therefore\ y^2=a^2\left(\frac{1}{m}-m\right)^2=a(x-3a),$$

the equation to a parabola.

(2) Find the equation to the tangent in the ellipse;



The centre being the origin,

$$\frac{y^2}{b^2}+\frac{x^2}{a^2}=1;\quad\therefore\ \frac{dy}{dx}=-\frac{b^2}{a^2}\cdot\frac{x}{y};$$

$$\therefore\ y_1-y=-\frac{b^2}{a^2}\frac{x}{y}(x_1-x);$$

$$\therefore\ yy_1-y^2=\frac{-b^2}{a^2}xx_1+\frac{b^2}{a^2}x^2=-\frac{b^2}{a^2}xx_1+b^2-y^2;$$

$$\therefore\ yy_1=\frac{b^2}{a^2}\cdot(a^2-xx_1),\quad\text{or}\quad y_1=\frac{b}{a}\cdot\frac{a^2-xx_1}{\sqrt{a^2-x^2}}.$$

$$\text{Let } y_1=0;\quad\therefore\ x_1=CT=\frac{a^2}{x}=\frac{CA^2}{CN};$$

$$\therefore\ CT\times CN=CA^2,\quad\text{(See } \textit{Conic Sections.)}$$

$$\text{and } NT=CT-CN=\frac{a^2-x^2}{x}=\text{sub-tangent,}$$

$$\text{or } NT\times CN=(a+x)(a-x)=A_1N\times AN.$$

Cor. 1. Make $x_1 = a$; $\therefore y_1 = AD = \dfrac{b.(a-x)}{\sqrt{a^2-x^2}}$,

$x_1 = -a$; $\therefore y_1 = A_1D_1 = \dfrac{b.(a+x)}{\sqrt{a^2-x^2}}$;

$\therefore AD.A_1D_1 = b^2 = CB^2$,

also $NG = -y\dfrac{dy}{dx} = \dfrac{b^2}{a^2}x = \dfrac{b^2}{a^2}CN$.

Cor. 2. The equation to the tangent may be written

$$a^2yy_1 + b^2xx_1 = a^2b^2;$$

$$\therefore y_1 = -\frac{b^2x}{a^2y}x_1 + \frac{b^2}{y}.$$

Let $m = -\dfrac{b^2x}{a^2y}$; $\therefore m^2a^2 = \dfrac{b^4x^2}{a^2y^2}$;

$$\therefore m^2a^2 + b^2 = b^2.\left(\frac{b^2x^2 + a^2y^2}{a^2y^2}\right) = \frac{a^2b^4}{a^2y^2} = \frac{b^4}{y^2};$$

$$\therefore y_1 = mx_1 + \sqrt{m^2a^2 + b^2},$$

an equation to the tangent in terms of the angle it makes with the axis of x.

Ex. Find the locus of the intersection of pairs of tangents to an ellipse at right angles to each other.

$y = mx + \sqrt{m^2a^2 + b^2}$, one tangent;

$$\therefore y = -\frac{x}{m} + \sqrt{\frac{a^2}{m^2} + b^2}, \text{ the other;}$$

$$\therefore y - mx = \sqrt{m^2a^2 + b^2},$$

$$my + x = \sqrt{m^2b^2 + a^2};$$

squaring and adding,

$$(1+m^2)(y^2+x^2) = (1+m^2)(a^2+b^2);$$

$$\therefore y^2 + x^2 = a^2 + b^2,$$

the equation to a circle, radius $= \sqrt{a^2+b^2}$.

Cor. If $b = a$, the ellipse becomes a circle and radius $= a\sqrt{2}$, the chord of a quadrant.

(3) In the hyperbola of which the equation is

$$a^2y^2 - b^2x^2 = -a^2b^2,$$

the equation to the tangent is

$$a^2yy_1 - b^2xx_1 = -a^2b^2;$$

$$\therefore\ y_1 = \frac{b^2x}{a^2y}x_1 - \frac{b^2}{y}$$

$$= mx_1 + \sqrt{m^2a^2 - b^2};\ \text{if}\ m = \frac{b^2x}{a^2y}.$$

Ex. Find the locus of the intersections of tangents to the hyperbola, and perpendiculars from the centre.

$y_1 = mx_1 + \sqrt{m^2a^2 - b^2}$, (1) is the tangent,

$y_1 = -\frac{1}{m}x_1$, (2) is the perpendicular;

$\therefore\ m = -\frac{x_1}{y_1}$; substituting in (1), and omitting the suffix,

$$y = -\frac{x^2}{y} + \sqrt{\frac{a^2x^2}{y^2} - b^2};$$

$$\therefore\ (y^2 + x^2)^2 = a^2x^2 - b^2y^2.$$

Cor. Let $b = a$, or the hyperbola be the rectangular;

$$\therefore\ (y^2 + x^2)^2 = a^2(x^2 - y^2),$$

the equation to the Lemniscata. In fact, the Lemniscata is commonly defined to be the locus of the intersection of the tangent to a rectangular hyperbola, with the perpendicular from the centre on the tangent.

(4) In the Cissoid, find the subtangent and subnormal, and equation to tangent.

Here $y^2 = \frac{x^3}{2a - x}$;

$$\therefore\ \text{subnormal} = y\frac{dy}{dx} = \frac{x^2(3a - x)}{(2a - x)^2};$$

$$\therefore\ \text{subtangent} = y\frac{dx}{dy} = \frac{x(2a - x)}{3a - x}.$$



The equation to the tangent is

Note.—*PTD* in the figure should be a straight line.

$$\therefore\ y_1 = \frac{\sqrt{x}}{(2a-x)^{\frac{1}{2}}} \cdot \{(3a-x)\,x_1 - ax\}.$$

Making y_1 and x_1 successively $= 0$,

$$AT = \frac{a \cdot x}{3a-x}, \text{ and } AD = -y_0 = a \cdot \left(\frac{x}{2a-x}\right)^{\frac{1}{2}}.$$

If $x = 2a$, $\frac{dy}{dx}$ and y_1 are infinite; there is therefore an asymptote through B perpendicular to the axis of x.

(5) Rectangular hyperbola referred to the asymptotes.

Here $yx = \frac{a^2}{2}$; $\therefore\ y = \frac{a^2}{2x}$,

$$\frac{dy}{dx} = -\frac{a^2}{2} \cdot \frac{1}{x^2} = -\frac{y}{x};$$

$$\therefore\ y_1 - y = -\frac{y}{x}(x_1 - x),$$



$$xy_1 - xy = -yx_1 + yx; \quad xy_1 + yx_1 = 2yx = a^2;$$

$$x_1 = 0, \quad y_0 = AD = \frac{a^2}{x}; \quad y_1 = 0, \quad x_0 = AT = \frac{a^2}{y}.$$

The $\Delta\, DAT = \frac{AT \cdot AD}{2} = \frac{a^4}{2xy} = a^2$, which is constant.

(6) Let $\sqrt{y} = \sqrt{a} - \sqrt{x}$; find equation to tangent, and shew that $AD + AT = a$.

$$\frac{dy}{dx} = -\frac{\sqrt{y}}{\sqrt{x}}; \quad \therefore\ y_1 - y = -\frac{\sqrt{y}}{\sqrt{x}}(x_1 - x).$$

Let $x_1 = 0$; $\therefore\ AD = y + \frac{x\sqrt{y}}{\sqrt{x}} = y + \sqrt{xy}$;

$$y_1 = 0; \quad \therefore\ AT = x + \frac{y\sqrt{x}}{\sqrt{y}} = x + \sqrt{xy};$$

$$\therefore\ AD + AT = x + 2\sqrt{xy} + y = (\sqrt{x} + \sqrt{y})^2 = a.$$

(7) Draw a tangent to the cycloid.

$$AN = x, \quad NP = y, \quad AB = 2a.$$

Then $\frac{dy}{dx} = \frac{\sqrt{2ax - x^2}}{x}$;

$\therefore\ y \cdot \frac{dx}{dy} = \frac{y \cdot x}{\sqrt{2ax - x^2}}$;

or $NT = \frac{NP \cdot AN}{NQ}$; or

$NT : NP :: AN : NQ$;



and $\angle N$ is common to Δs ANQ, TPN; $\therefore$ they are similar, and $\angle PTN = \angle QAN$; $\therefore$ the tangent TP is parallel to the chord AQ.

Also since $\angle AQB$ is always $= 90$, PG is parallel to BQ.

(8) Draw an asymptote to the hyperbola.

$$y = \pm \frac{b}{a}\sqrt{2ax + x^2} = \pm \frac{b}{a} x \left(1 + \frac{2a}{x}\right)^{\frac{1}{2}}$$

$$= \pm \frac{b}{a} x \left\{1 + \tfrac{1}{2}\frac{2a}{x} + \frac{\frac{1}{2} \cdot (\frac{1}{2} - 1)}{1 \cdot 2} \cdot \frac{4a^2}{x^2} + \frac{B}{x^3} + \&c.\right\}$$

$$= \pm \frac{b}{a} \cdot \left\{x + a - \frac{1}{2} \cdot \frac{a^2}{x} + \frac{B}{x^2} + \&c.\right\},$$

and therefore $y = \pm \frac{b}{a}(x + a)$ is the equation to two asymptotes;

and since if $x = 0$, $y = \pm b$; and if $y = 0$, $x = -a$, both will pass through the centre, and be equally inclined to the axis of x.

(9) Draw the asymptote to the curve.

$$y^3 = x^3 + ax^2 = x^3 \left(1 + \frac{a}{x}\right);$$

$$\therefore\ y = x \left(1 + \frac{a}{x}\right)^{\frac{1}{3}} = x \left\{1 + \tfrac{1}{3} \cdot \frac{a}{x} + \frac{A}{x^2} + \frac{B}{x^3} + \&c.\right\}$$

$$= x + \frac{a}{3} + \frac{A}{x} + \frac{B}{x^2} + \&c.;$$

$\therefore\ y = x + \frac{a}{3}$ is the equation to the asymptote which cuts the axis of x at an $\angle = 45^0$, and at a point $x = -\frac{a}{3}$.

(10) Let $y.(ax + b^2) = x^3$, draw the asymptotes.

$$y = \frac{x^3}{ax + b^2} = \frac{1}{a}\frac{x^3}{x + c} \text{ by putting } \frac{b^2}{a} = c$$

$$= \frac{1}{a}.\frac{x^2}{1 + \frac{c}{x}} = \frac{1}{a}x^2\left\{1 - \frac{c}{x} + \frac{c^2}{x^2} - \frac{c^3}{x^3} + \&c.\right\}$$

$$= \frac{x^2}{a} - \frac{cx}{a} + \frac{c^2}{a} - \frac{c^3}{ax} + \&c.;$$

$\therefore ay = x^2 - cx + c^2$ is the equation to the asymptotic curve.

This being put under the form

$$ay - \tfrac{3}{4}c^2 = x^2 - cx + \frac{c^2}{4} = \left(x - \frac{c}{2}\right)^2,$$

$$\text{or } \left(x - \frac{c}{2}\right)^2 = a\left(y - \tfrac{3}{4}\frac{c^2}{a}\right),$$

shews that the curve is a parabola, the axis of which is perpendicular to the axis of x, and the vertex determined by making $x_1 = \frac{c}{2}$ and $y_1 = \frac{3}{4}\frac{c^2}{a}$; the latus rectum $= a$.

The curve has also a rectilinear asymptote parallel to the axis of y; for making $x = -c = -\frac{b^2}{a}$, y becomes infinite, and $\frac{dy}{dx}$ is also infinite; therefore a line drawn perpendicularly to the axis of x at a distance $-\frac{b^2}{a}$ from the origin, will when produced to infinity be a tangent to the curve.

178. When the equation between y and x cannot be solved with regard to y, it is frequently very convenient to substitute zx for y; and then from the given equation to find x and y in terms of z; and z being so assumed as to make x and y infinite, the inclination of the asymptote, if any, to the axis of x is known (for $z = \frac{y}{x}$ = tan inclination); and AD and AT being determined, the position of the asymptote may also be determined.

(11) Let $ay^4 - bx^4 + c^3xy = 0$; make $y = xz$;

$$\therefore ax^4z^4 - bx^4 + c^3x^2z = 0;$$

$$\therefore x^2 = \frac{c^3 x}{b - ax^4} = \infty \text{ ; if } ax^4 = b \text{; or } x = \sqrt[4]{\frac{b}{a}};$$

$$\therefore y = x\sqrt[4]{\frac{b}{a}} \text{ is the equation to the asymptote, since}$$

$$AD = y - px = \frac{-2c^3xy}{4ay^3 + c^3x} = 0 \text{; if } y = \infty \text{; and } x = \infty,$$

$$\text{and } AT = x - \frac{y}{p} = \frac{2c^3xy}{4ay^3 + c^2x} = 0 \text{; if } y = \infty \text{; and } x = \infty.$$

(12) Find when the curve, which is the locus of the general equation of the second order, has an asymptote.

Here $Ay^2 + Bxy + Cx^2 + Dy + Ex + F = 0$,

or $y^2 + 2(ax + b)y + cx^2 + ex + f = 0$,

dividing by A, and making the proper substitutions;

$$\therefore y^2 + 2(ax + b)y + (ax + b)^2 = (a^2 - c)x^2 + (2ab - e)x + b^2 - f,$$

$$\text{and } y = -(ax + b) \pm \sqrt{(a^2 - c)x^2 + (2ab - e)x + (b^2 - f)};$$

$$\therefore = -x\left\{\left(a + \frac{b}{x} \pm \sqrt{a^2 - c + \frac{2ab - e}{x} + \frac{b^2 - f}{x^2}}\right\}\right.$$

$$= -x\left\{a + \frac{b}{x} \pm \sqrt{a^2 - c}\left(1 + \frac{1}{2x}\frac{2ab - e}{(a^2 - c)} + \frac{A}{x^2} + \frac{B}{x^3}\right)\right\}$$

$$= -\left\{ax + b \pm \sqrt{a^2 - c}\left(x + \tfrac{1}{2}\frac{2ab - e}{a^2 - c} + \frac{A}{x} + \text{\&c.}\right)\right\};$$

and therefore the equation to the asymptote is

$$y = -(a \pm \sqrt{a^2 - c})x - \left(b \pm \frac{2ab - e}{2\sqrt{a^2 - c}}\right),$$

which is possible when $a^2 > c$, or $\frac{B^2}{4A^2} > \frac{C}{A}$, or $B^2 - 4AC > 0$, or when the curve is the hyperbola.

Cor. If $a^2 = c$, the curve is the parabola, and the equation being of the form $y = -(ax + b) \pm \sqrt{mx} \cdot \sqrt{1 + \frac{n}{x}}$ cannot be reduced to the form $y = Ax + B$.

Prob. Find that tangent line to a given curve which cuts off from the co-ordinate axes the greatest area.

$$\text{Area} = \tfrac{1}{2} x_0 y_0 = \tfrac{1}{2}(y - px)\left(x - \frac{y}{p}\right)$$

$$= -\tfrac{1}{2}\frac{(y-px)^2}{p}; \quad \text{where } p = \frac{dy}{dx};$$

$$\therefore\ u = \frac{(y-px)^2}{p} \text{ a maximum};$$

$$\therefore\ \frac{du}{dx} = \frac{2}{p}(y-px)\left(\frac{dy}{dx} - p - x\frac{dp}{dx}\right) - \frac{(y-px)}{p^2}\frac{dp}{dx} = 0;$$

$$\therefore\ -2px - (y - px) = 0;$$

$$\therefore\ x = \tfrac{1}{2}\left(\frac{y-px}{p}\right) = \tfrac{1}{2}x_0.$$

Therefore also $y = \tfrac{1}{2}y_0$; and the tangent is bisected at the point of contact.

Cor. From this it appears that the least polygon of a given number of sides which will circumscribe a given oval, must be such that every side of the polygon must be bisected at the point of its contact with the oval, since the interior area must be least, when the triangular spaces as in the preceding problem are greatest.

In the problem the axes are supposed rectangular, but if they be oblique, the same results are obtained, but area $= \tfrac{1}{2} x_0 y_0 \times$ sine of inclination of axes.

EXAMPLES.

(1) Let $y^n = a^{n-1}x$; $NT = nx$; $NG = \dfrac{y^2}{nx}$.

If $n = 2$, the curve is the parabola; $NT = 2x$, $NG = \dfrac{a}{2}$.

(2) Let the curve be the Witch:

$$NT = -\frac{2ax - x^2}{a}; \quad NG = -\frac{4a^3}{x^2}.$$

(3) The focus of a parabola is in the centre of a given circle, its vertex bisects the radius; find the point and angle of intersection of circle and parabola.

(4) Shew that the curve defined by $y^2 = 4ax$, intersects the curve defined by $y^2 = \frac{4}{27a}(x - 2a)^3$, at a point where $x = 8a$, and find the angle of intersection.

(5) If $y^2 = 4a(x + a)$ be the equation to a parabola, the origin in the focus, shew that the points of intersection of the tangents, and perpendiculars from the focus, are determined by the equations

$$x_1 = -a, \text{ and } y_1 = \frac{y}{2}.$$

(6) The locus of the intersections of tangents to the parabola with perpendiculars from the vertex, is the cissoid.

(7) Find the length of the perpendicular drawn from the focus of an ellipse upon the tangent, and shew that the locus of their intersection is a circle, radius = a.

(8) Given two points A and B, find the locus of P when the angle PBA is double of the angle PAB, and draw an asymptote to the curve traced by P.

(9) Draw an asymptote to the curve defined by $y^3 + x^3 = 3axy$, and determine the points where the tangents are parallel to the two axes.

(10) Find the point and angle at which the curve $3y^2 = x(x + 2)^2$ cuts the axis. At the origin, angle = 90.

(11) Find the same when $y - 2 = (x - 1)\sqrt{x - 2}$, and the values of x and y when the tangent is perpendicular to the axis of x. (1) $y = 0$; $x = 3$. (2) $y = 2$; $x = 2$.

(12) If $y^2 = 3x^2 - x^3$ draw an asymptote to the curve; find its greatest ordinate; and the angles the curve makes with the axis of x at the points, $x = 0$, $x = 2$ and $x = 3$. Asymptote $y = -x + 1$; ordinate = 2; angles are 90, 0, 90.

(13) Draw the asymptotes, (1) when $y = \frac{x^3 + ax^2 + a^3}{x^2 - a^2}$, (2) when $y^5 = ax^4 + mx^5$, and (3) when $y = a^x$.

(14) If in the ellipse, $\theta = \angle CPG$, and l = the angle PG makes with the axis major; $\tan\theta = \frac{\tan l\,(a^2 - b^2)}{a^2 + b^2\tan^2 l}$.

(15) Shew that the curve whose equation is $\frac{y}{r} = \sin\frac{a}{r}$;

where $y^2 - 2rx + x^2 = 0$; intersects the axis of x at points determined by $x = \frac{2a}{n\pi}$, n being any whole number.

(16) The normal to the curve defined by $y^2 = 4ax$, is a tangent to the curve in which $y^2 = \frac{4}{27a} \cdot (x - 2a)^3$.

(17) Draw a tangent to a circle, cutting the axis of x at 30^0.

(18) In the conchoid, where $x^2y^2 = (a + x)^2 \cdot (b^2 - x^2)$,

$$NT = - \frac{x \cdot (a + x) \cdot (b^2 - x^2)}{x^3 + ab^2}.$$

Before differentiating take the logarithm of the equation.

(19) Draw a tangent to $x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}$; shew that the part of the tangent intercepted between the axes $= a$, and that perpendicular on tangent $= \sqrt[3]{axy}$.

(20). The centre of an ellipse is the vertex of a parabola, the major axis of ellipse is perpendicular to the axis of parabola, and the curves intersect at right angles, prove that major axis : minor axis :: $\sqrt{2}$: 1.

(21) If PY and QY be respectively tangents to corresponding arcs of a cycloid and the generating circle, shew that the locus of Y is the involute of the circle.

(22) Find the angle of intersection of a rectangular hyperbola and a circle having the same centre, radius of circle being $2a$;

$$\text{angle} = \sin^{-1} \tfrac{1}{4} \sqrt{15}.$$

(23) If TP and TQ be tangents to a parabola, and S be the focus, shew that

$$SP \cdot SQ = ST^2.$$

(24) If $2y = c(e^{\frac{x}{c}} + e^{-\frac{x}{c}})$, (the equation to the catenary),

$$\text{the normal} = \frac{1}{c} y^2.$$

(25) If $\frac{1}{y^n} = \frac{1}{a^n} + \frac{1}{x^n}$ be the equation to a curve, shew that

$$a^n \cdot AT = x^{n+1}.$$

(26) Find that point in a parabola, at which a line from the vertex makes the greatest angle with the curve.

Ans. $x = 4a$.

(27) If A be the extremity of the diameter of a circle, PT a tangent, PN an ordinate, and AP a chord, prove that AP bisects the angle TPN.

(28) If $y^n - (a + bx)\,y^{n-1} + (c + ex + fx^2)\,y^{n-2} - \&c. = 0$, be the equation to a curve of n dimensions, prove that the sum of the ordinates divided by their respective subtangents, is a constant quantity.

(29) If C be the centre of a circle, AQ a chord, and let CNR be drawn cutting AQ in N, draw NP perpendicular to AQ and $= NR$; find the locus of P and draw its asymptote.

If $2c$ be the length of chord, a = radius of circle, origin centre of chord, and the chord be the axis of x, then

$$(y - a)^2 = x^2 + a^2 - c^2.$$

(30) ABD is a semicircle, centre C, and diameter AD, EF is a chord parallel to AD, CQR a radius cutting EF in Q, bisect QR in P; find the locus of P and the position of the asymptote. The curve is the conchoid.

CHAPTER XII.

THE DIFFERENTIALS OF THE AREAS AND LENGTHS OF CURVES: OF THE SURFACES AND VOLUMES OF SOLIDS OF REVOLUTION: SPIRALS.

---

179. ONE of the applications of the Integral Calculus is to find the areas of curves included between given ordinates, the lengths of their arcs, and the surfaces and contents of solids.

The solids of which we shall treat are called solids of revolution, since they may be supposed to be generated by the revolution of a plane figure round a line, thus termed an axis. Hence it follows that every section perpendicular to the axis will be a circle, the radius of which is the revolving ordinate, and every section made by a plane passing through the axis will reproduce the original area.

Considering the areas and lengths of curves, and the contents and surfaces of solids, to be functions of one of the quantities x or y, we can, by the Differential Calculus, find equations between the differential coefficients of these functions, and expressions containing x or y, by which we shall hereafter obtain the values of the functions themselves.

We shall find it useful first to establish the truth of the following Proposition.

180. If $A + Bx$, $A_1 + B_1x$, and $A + B_2x$ be three algebraical expressions taken in order of magnitude, viz., the first greater than the second, and the second greater than the third, then shall

$$A_1 = A, \text{ whatever be the value of } x.$$

For $(A + Bx) - (A + B_2x)$ is $> (A + Bx) - (A_1 + B_1x)$;

$$\therefore \text{ if } (A + Bx) - (A + B_2x) = 0, \text{ or } \frac{A + Bx}{A + B_2x} = 1;$$

$$\therefore \textit{ à fortiori } \text{will } \frac{A + Bx}{A_1 + B_1x} = 1.$$

But as x decreases $\dfrac{A + Bx}{A + B_2x}$ approaches $\dfrac{A}{A}$ or 1; and when x is diminished without limit it actually equals unity, and

$$\therefore \frac{A + Bx}{A_1 + B_1 x} \text{ which becomes } \frac{A}{A_1},$$

by the continued diminution of x, also is equal to unity;

$$\therefore \frac{A}{A_1} = 1, \text{ or } A_1 = A.$$

AREA OF A CURVE.

181. Let AP be a curve, $y = f(x)$, the equation to it, where $AN = x$, $NP = y$; and let A = area ANP.



Then $\frac{dA}{dx} = y.$

Let $NN_1 = h$. Complete the parallelograms QN_1 and PN_1.

Then the area P_1PNN_1 is $> \square PN_1$, $< \square QN_1 \ldots (1)$.

Now A depends upon x; for as x changes, A changes;

$$\therefore A = ANP = \phi(x); \text{ and } \therefore AN_1P_1 = \phi(x + h);$$

$$\therefore PP_1N_1N = \phi(x + h) - \phi(x) = \frac{dA}{dx}h + \frac{d^2A}{dx^2}\frac{h^2}{1.2} + \&c.$$

$$\text{and } \square PN_1 = yh.$$

$$\square QN_1 = h \times P_1N_1 = h.f(x + h) = h\{y + ph + Ph^2\}; \; p = \frac{dy}{dx};$$

therefore, dividing by h, we have by (1),

$$\frac{dA}{dx} + \frac{d^2A}{dx^2}\frac{h}{1.2} + \&c. > y < y + ph + Ph^2;$$

$$\text{i.e. } y + ph + Ph^2, \; \frac{dA}{dx} + \frac{d^2A}{dx^2}\frac{h}{1.2} + \&c. \text{ and } y$$

are in order of magnitude; whence, by the Lemma,

$$\frac{dA}{dx} = y.$$

LENGTH OF A CURVE.

182. If s = length of the curve AP;

$$\frac{ds}{dx} = \sqrt{1 + \frac{dy^2}{dx^2}}.$$

Draw tangent PM, and chord PP_1.



Then arc $PP_1 >$ chord $PP_1 <$ $PM + MP_1$.

But arc $PP_1 = AP_1 - AP = \phi(x+h) - \phi(x) = \frac{ds}{dx}h + \frac{d^2s}{dx^2}\frac{h^2}{1.2} + \&c.$

chord $PP_1 = \sqrt{Pm^2 + (P_1m)^2}$* $= \sqrt{h^2 + (ph + Ph^2)^2}$

$$= h\sqrt{(1+p^2) + 2Pph + P^2h^2},$$

$$PM = \sqrt{Pm^2 + Mm^2}* = \sqrt{h^2 + p^2h^2} = h\sqrt{1+p^2},$$

$$MP_1 = MN_1 - N_1P_1 = (y + ph) - (y + ph + Ph^2) = -Ph^2;$$

whence, dividing by h,

$$\frac{ds}{dx} + \frac{d^2s}{dx^2}\cdot\frac{h}{1.2} + \&c. > \sqrt{1+p^2+2Pph+P^2h^2} < \sqrt{1+p^2} - Ph$$

$$> \sqrt{1+p^2} + \frac{Pp}{\sqrt{1+p^2}}h + \&c. < \sqrt{1+p^2} - Ph;$$

$$\therefore \frac{ds}{dx} = \sqrt{1+p^2} = \sqrt{1 + \frac{dy^2}{dx^2}}.$$

VOLUME OF SOLID.

183. If V be the volume of a solid of revolution APp,

$$\frac{dV}{dx} = \pi y^2.$$

* $P_1m = P_1N_1 - PN = y + ph + Ph^2 - y = ph + Ph^2$,
$Mm = Pm \,.\, \tan MPm = h \,.\, \tan PTN = h \cdot \frac{dy}{dx}$.



Let $\left.\begin{matrix} AN = x \\ NP = y \\ NN_1 = h \end{matrix}\right\}$; $\therefore AP_1p_1 = f(x+h) = V + \frac{dV}{dx}h + \frac{d^2V}{dx^2}\frac{h^2}{1.2} + \&c.$

Then the solid Ppp_1P_1 is $>$ cylinder PMm_1p,

$<$ cylinder RP_1p_1r;

i.e. $\frac{dV}{dx}h + \frac{d^2V}{dx^2}\frac{h^2}{1.2} + \&c. > \pi y^2 h$,

$< \pi (y + ph + Ph^2)^2 h$,

or $\frac{dV}{dx} + \frac{d^2V}{dx^2}\frac{h}{1.2} + \&c. > \pi y^2 < \pi (y + ph + Ph^2)^2$,

or $> \pi y^2 < \pi y^2 + 2\pi pyh + \&c.$

whence $\frac{dV}{dx} = \pi y^2$.

Prob. The surface of a truncated cone, of which the radii of the greater and smaller ends are a, b, and the slant side s, $= \pi s (a + b)$.

Let l = length of cone, radius of the base $= a$,

$l_1 = \ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots = b$;

therefore, surface of frustum

$$= \pi la - \pi l_1 b = \pi \{sa + l_1 (a - b)\},$$

but l or $l_1 + s : l_1 :: a : b$;

$$\therefore s : l_1 :: a - b : b;$$

$$\therefore sb = l_1 (a - b);$$

$\therefore$ surface of frustum $= \pi s . (a + b)$.

SURFACE OF SOLID.

184. If S = surface of the solid of revolution APp,

$$\frac{dS}{dx} = 2\pi y \sqrt{1 + \frac{dy^2}{dx^2}}.$$



$\left.\begin{aligned} AN &= x \\ NP &= y \\ AP &= s \\ NN_1 &= h \end{aligned}\right\}$. Draw the tangent PM, and chord PP_1.

Then, surface generated by arc PP_1, will be

$>$ than that by chord PP_1, $<$ by PM and MP_1.

Now chords PP_1 and PM generate truncated cones, of which the surfaces respectively are

$$\pi \{PN + P_1N_1\} PP_1, \text{ and } \pi \{PN + MN_1\} PM;$$

and MP_1 will generate a circular zone $= \pi (MN_1^2 - N_1P_1^2)$;
and the surface generated by arc PP_1

$$= \frac{dS}{dx} h + \frac{d^2S}{dx^2} \frac{h^2}{1,2} + \&c.$$

But $\{PN + P_1N_1\} PP_1$

$$= (2y + ph + Ph^2) \cdot \sqrt{h^2 + (ph + Ph^2)^2}$$

$$= (2y + ph + Ph^2) h \sqrt{1 + p^2 + Mh},$$

where Mh = terms involving h;

and $(PN + MN_1) PM$

$$= (2y + ph) \cdot \sqrt{h^2 + p^2h^2} = (2y + ph) h \sqrt{1 + p^2},$$

also $MN_1^2 - N_1P_1^2$

$= (y + ph)^2 - (y + ph + Ph^2)^2 = - Ph^2(2y + 2ph + Ph^2)$

$= - Nh^2$, by substitution;

$$\therefore \frac{dS}{dx} + \frac{d^2S}{dx^2}\frac{h}{1.2} + \&c. > \pi(2y + ph + Ph^2)\sqrt{1 + p^2 + Mh}$$

$$< \pi(2y + ph)\sqrt{1 + p^2} - \pi Nh$$

$$> 2\pi y\sqrt{1 + p^2 + Mh} + \text{terms involving } h,$$

$$< 2\pi y\sqrt{1 + p^2} + \pi ph\sqrt{1 + p^2} - \pi Nh$$

$$> 2\pi y\sqrt{1 + p^2} + \pi\frac{yMh}{\sqrt{1 + p^2}} + \&c.$$

$$< 2\pi y\sqrt{1 + p^2} + \pi ph\sqrt{1 + p^2} - \pi Nh;$$

$$\therefore \frac{dS}{dx} = 2\pi y\sqrt{1 + p^2} = 2\pi y\sqrt{1 + \frac{dy^2}{dx^2}} = 2\pi y\frac{ds}{dx}.$$

Cor. 1. Hence it appears, that if in a curve ANP, A = area of ANP; and δA be the differential of the area, and δx the differential of the abscissa; δx being very small, that

$$\frac{\delta A}{\delta x} = \frac{dA}{dx} = y; \quad \therefore \delta A = y\delta x,$$

or the differential of the area of a curve is equal to the rectangle of the ordinate, and the increment of the corresponding abscissa.

Cor. 2. If δs be the increment of the arc AP; corresponding to the increments δx and δy of x and y;

$$\text{then since } \frac{\delta s}{\delta x} = \frac{ds}{dx};$$

$$\therefore \frac{\delta s}{\delta x} = \sqrt{1 + \frac{dy^2}{dx^2}} = \sqrt{1 + \frac{\delta y^2}{\delta x^2}};$$

$$\therefore \delta s = \sqrt{\delta y^2 + \delta x^2};$$

whence it follows that the increment of the arc is the hypothenuse of a right-angled triangle, of which δx and δy are the other two sides.

Hence also it appears that the chord PP_1 = arc PP_1 ultimately.

Cor. 3. If α be the angle which the tangent makes with x;

$$\therefore \frac{\delta s}{\delta x} = \sec \alpha; \quad \delta s = \delta x \,.\, \sec \alpha.$$

Cor. 4. In the same manner, if δV and δS be the differentials of the volume V, and of the surface S of a solid,

$$\frac{\delta V}{\delta x} = \pi y^2; \quad \therefore \delta V = \pi y^2 \,.\, \delta x,$$

$$\text{and } \frac{\delta S}{\delta x} = 2\pi y \,.\, \sqrt{1 + \frac{dy^2}{dx^2}} = 2\pi y \,.\, \frac{\delta s}{\delta x};$$

$$\therefore \delta S = 2\pi y \,.\, \delta s;$$

from the former of which expressions it follows that the differential of a solid is a cylinder, base πy^2, and altitude δx: and that the differential of a surface is the convex surface of a cylinder, the circumference of whose base is $2\pi y$, and altitude δs.

SPIRALS.

185. The expressions just obtained, and those of the preceding Chapter, are only applicable, when the equation to the curve is known in terms of the rectangular co-ordinates; we shall now find corresponding expressions for the perpendicular upon the tangent, the area and length of a curve, when referred to polar co-ordinates; that is, when $r = f(\theta)$, or $p = f(r)$, p being the perpendicular on the tangent, r the radius vector, and θ the angle traced out by r.

186. To find the expression for the perpendicular on the tangent in polar curves.

$SN = x$, $SP = r$,

$NP = y$, $SY = p$, and $\angle ASP = \theta$.



Now, Art. 169, $SY = p = \dfrac{y - x\dfrac{dy}{dx}}{\sqrt{1 + \dfrac{dy^2}{dx^2}}}$.

But $x = SP \cos PSN = -r \cos\theta = f(\theta)$,

$y = SP \sin PSN = +r \sin\theta = \phi(\theta)$,

$$\text{and } \frac{dy}{dx} = \left(\frac{dy}{d\theta}\right)\left(\frac{d\theta}{dx}\right) = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}};$$

$$\therefore p = \frac{y \cdot \frac{dx}{d\theta} - x\frac{dy}{d\theta}}{\sqrt{\frac{dx^2}{d\theta^2} + \frac{dy^2}{d\theta^2}}}.$$

$$\text{But } \frac{dx}{d\theta} = r\sin\theta - \cos\theta \cdot \frac{dr}{d\theta};$$

$$\text{and } \frac{dy}{d\theta} = r\cos\theta + \sin\theta \cdot \frac{dr}{d\theta};$$

$$\therefore y\frac{dx}{d\theta} - x\frac{dy}{d\theta} = r^2\sin^2\theta + r^2\cos^2\theta = r^2,$$

$$\frac{dx^2}{d\theta^2} + \frac{dy^2}{d\theta^2} = \left(r^2 + \frac{dr^2}{d\theta^2}\right)(\sin^2\theta + \cos^2\theta) = r^2 + \frac{dr^2}{d\theta^2};$$

$$\therefore p = \frac{r^2}{\sqrt{r^2 + \frac{dr^2}{d\theta^2}}};$$

whence having given $\theta = f(r)$, p may be found in terms of r, or of r and θ; but the formula may be put under a more convenient form for practice;

$$\therefore \frac{1}{p^2} = \frac{r^2 + \frac{dr^2}{d\theta^2}}{r^4} = \frac{1}{r^2} + \frac{1}{r^4} \cdot \frac{dr^2}{d\theta^2}.$$

$$\text{Let } r = \frac{1}{u}; \quad \therefore u = \frac{1}{r}; \quad \therefore \frac{du}{d\theta} = -\frac{1}{r^2} \cdot \frac{dr}{d\theta};$$

$$\therefore \frac{1}{p^2} = u^2 + \frac{du^2}{d\theta^2}.$$

187. To find the differential of the area.

Let A = area ANP, then will $\frac{dA}{d\theta} = \frac{1}{2}r^2$.

For $ASP = ANP - SNP = ANP - \frac{yx}{2}$.

$$\text{But } \frac{d.ANP}{dx} = y; \quad \frac{d.(ANP)}{d\theta} = y.\frac{dx}{d\theta};$$

$$\therefore \frac{dA}{d\theta} = y.\frac{dx}{d\theta} - \tfrac{1}{2}.\left(y\frac{dx}{d\theta} + x\frac{dy}{d\theta}\right)$$

$$= \tfrac{1}{2}.\left(y\frac{dx}{d\theta} - x\frac{dy}{d\theta}\right)$$

$$= \tfrac{1}{2}r^2.\text{(Art. 186).}$$

Cor. 1. If $\theta = f(r)$, to find $\frac{d\theta}{dr}$.

$$\text{Since } \frac{1}{p^2} = \frac{1}{r^2} + \frac{1}{r^4}.\frac{dr^2}{d\theta^2};$$

$$\therefore \frac{dr^2}{d\theta^2} = r^4\left(\frac{1}{p^2} - \frac{1}{r^2}\right) = r^2.\frac{r^2 - p^2}{p^2};$$

$$\therefore \frac{d\theta}{dr} = \frac{p}{r\sqrt{r^2 - p^2}},$$

a useful equation in dynamics.

188. To find the differential of the length of a spiral.

Let $s = AP$; and let y and x be the co-ordinates of P;

$$\therefore \frac{ds}{dx} = \sqrt{1 + \frac{dy^2}{dx^2}};$$

but s, x, y being functions of θ,

$$\frac{ds}{dx} = \frac{ds}{d\theta}.\frac{d\theta}{dx}; \quad \frac{dy}{dx} = \frac{dy}{d\theta}.\frac{d\theta}{dx};$$

$$\therefore \frac{ds}{d\theta} = \sqrt{\frac{dx^2}{d\theta^2} + \frac{dy^2}{d\theta^2}} = \sqrt{r^2 + \frac{dr^2}{d\theta^2}};$$

189. To find the same, if $s = f(r)$.

$$\frac{ds}{dr} = \frac{ds}{d\theta}.\frac{d\theta}{dr} = \sqrt{r^2\frac{d\theta^2}{dr^2} + 1}$$

$$= \sqrt{\frac{p^2}{r^2 - p^2} + 1} = \frac{r}{\sqrt{r^2 - p^2}}.$$

Cor. Hence $\cos SPY = \frac{\sqrt{r^2 - p^2}}{r} = \frac{dr}{ds}$.

190. To find the tangent of the angle which SP makes with the curve.

$$\text{Tan } SPY = \frac{SY}{PY} = \frac{p}{\sqrt{r^2 - p^2}} = r \cdot \frac{d\theta}{dr},$$

$$\left(\text{or if } r = \frac{1}{u}\right) = -u \cdot \frac{d\theta}{du}.$$

191. To draw a tangent to a spiral.



P the point to which the tangent is to be drawn. S the pole. Join SP. Suppose PT to be the tangent. Draw $SY \perp PT$, and $ST \perp PS$, then ST is called the sub-tangent.

$$\text{And } ST = SP \cdot \frac{SY}{PY} = \frac{pr}{\sqrt{r^2 - p^2}} = \text{also } r^2 \cdot \frac{d\theta}{dr}.$$

Find therefore from the equation to the spiral $\frac{pr}{\sqrt{r^2 - p^2}}$, or $r^2 \cdot \frac{d\theta}{dr}$, according as the equation is $p = f(r)$, or $\theta = f(r)$. Draw ST perpendicular to SP and equal to either of these values. Join TP, it is the tangent.

Cor. Since $ST = \pm r^2 \cdot \frac{d\theta}{dr} = \mp \frac{d\theta}{du}$; $\therefore \frac{1}{ST^2} = \left(\frac{du}{d\theta}\right)^2$;

$$\therefore \frac{1}{SY^2} = u^2 + \left(\frac{du}{d\theta}\right)^2 = \frac{1}{SP^2} + \frac{1}{ST^2}.$$

192. Asymptotes to Spirals.

If ST remain finite when SP is infinite, a tangent may be drawn which will touch the curve at a point infinitely distant from S, and is therefore an asymptote. And since those lines are said to be parallel which coincide only at an infinite distance; the asymptote must be drawn parallel to the infinite line SP.

Hence to construct; find θ and $r^2 \cdot \frac{d\theta}{dr}$ when r is infinite. Draw SP at the angle thus found, ST perpendicular to SP, and TP parallel to the infinite radius vector, TP produced is the asymptote.

193. Asymptotic circle.

If in the equation $\theta = f(r)$, θ becomes infinite when $r = a$; but impossible if r be $> a$; then if we describe a circle

with radius a, the spiral will make an infinite number of revolutions within the circle, and constantly approach the circumference, without exactly reaching it. In this case, the circle is called an exterior asymptotic circle. But if $r = a$ make θ infinite, and $r < a$, make θ impossible, the revolutions of the spiral will be without the circle to which it is always tending; and the circle is an interior asymptote.

Ex. 1. $\theta = \log\left(\frac{a-r}{a+r}\right)$.

Ex. 2. $\theta = \log\left(\frac{r-a}{r+a}\right)$.

194. The preceding expressions may be conveniently obtained by a direct investigation.



Let area $ASP = A$; $SP = r$,

$\angle ASP = \theta$; $SY = p$.

Draw SQ very near to SP,

and draw $PT \perp SQ$.

Then area $PSQ = \delta A$,

$\angle PSQ = \delta\theta$; $QT = \delta r$;

now PT is ultimately = a circular arc;

$$\therefore \delta A = \tfrac{1}{2} SQ \times PT \text{ very nearly}$$
$$= \tfrac{1}{2}(r + \delta r) \cdot r\delta\theta \text{ very nearly}$$
$$= \tfrac{1}{2}r^2\delta\theta + \tfrac{1}{2}r\delta\theta\delta r;$$
$$\therefore \frac{\delta A}{\delta\theta} = \tfrac{1}{2}r^2 + \tfrac{1}{2}r\delta r.$$

But as Q approaches P, δr continually diminishes, and ultimately vanishes; but then $\frac{\delta A}{\delta\theta} = \frac{dA}{d\theta}$,

$$\text{and } \therefore \frac{dA}{d\theta} = \tfrac{1}{2}r^2.$$

195. Again $(\text{chord } PQ)^2 = PT^2 + QT^2 = r^2 \cdot \delta\theta^2 + (\delta r)^2$;

$$\therefore \frac{(\text{chord } PQ)^2}{\delta\theta^2} = \frac{\delta s^2}{\delta\theta^2} = r^2 + \left(\frac{\delta r}{\delta\theta}\right)^2,$$

$$\text{whence } \frac{ds}{d\theta} = \sqrt{1 + \frac{dr^2}{d\theta^2}}.$$

Also from similar triangles, SPY and PQT,

$$\frac{PT}{QT} = \frac{SY}{PY}; \quad r \cdot \frac{\delta\theta}{\delta r} = \frac{p}{\sqrt{r^2 - p^2}};$$

$$\therefore \frac{d\theta}{dr} = \frac{p}{r\sqrt{r^2 - p^2}} \quad \ldots\ldots\ldots\ldots\ldots(1).$$

And $$\frac{PQ}{QT} = \frac{\delta s}{\delta r} = \frac{SP}{PY} = \frac{r}{\sqrt{r^2 - p^2}};$$

$$\therefore \frac{ds}{dr} = \frac{r}{\sqrt{r^2 - p^2}}.$$

196. Since $\tan SPY = \tan PQT = \dfrac{PT}{QT}$;

$$\therefore \tan SPY = r \cdot \frac{\delta\theta}{\delta r} = r \cdot \frac{d\theta}{dr},$$

$$\sin SPY = \frac{PT}{PQ} = r \cdot \frac{\delta\theta}{\delta s} = r \cdot \frac{d\theta}{ds},$$

$$\cos SPY = \frac{QT}{PQ} = \frac{\delta r}{\delta s} = \frac{dr}{ds}.$$

Also $SY : SP :: PT : PQ$;

$$\therefore p = r \cdot \frac{PT}{PQ} = r^2 \cdot \frac{\delta\theta}{\delta s} = r^2 \cdot \frac{d\theta}{ds}.$$

It is necessary that the student should be familiar with these methods of obtaining the above results from the triangle SPY, and the ultimate rectilinear triangle PQT; as he will find their use in dynamics and in the early sections of Newton's Principia.

197. Prob. To find the equation to the curve which is the locus of the intersection of the tangent and the perpendicular from the pole; or to find the locus of Y.

$SP = r$; $SY = p$, and let p_1 be the perpendicular from S on the tangent to the curve traced by Y.

Let $\theta = \angle ASP$; $\phi = \angle ASY$. See fig. Art. 194;

$$\therefore d\theta = \frac{p\,dr}{r\sqrt{r^2 - p^2}}; \quad d\phi = \frac{p_1\,dp}{p\sqrt{p^2 - p_1^2}}.$$

But $$\phi = \theta - PSY = \theta - \cos^{-1} \cdot \frac{p}{r};$$

$$\therefore \frac{p_1 dp}{p\sqrt{p^2 - p_1^2}} = \frac{pdr}{r\sqrt{r^2 - p^2}} + \frac{rdp - pdr}{r\sqrt{r^2 - p^2}}$$

$$= \frac{rdp}{r\sqrt{r^2 - p^2}};$$

$$\therefore \frac{p_1}{\sqrt{p^2 - p_1^2}} = \frac{p}{\sqrt{r^2 - p^2}}; \quad \therefore \frac{p^2}{p_1^2} = \frac{r^2}{p^2};$$

$$\therefore p_1^2 = \frac{p^4}{r^2}; \text{ and } p_1 = \frac{p^2}{r};$$

which is the value of the perpendicular on the tangent in the required curve: and if r_1 be put for p; and since r is a function of $p = f(r_1)$,

$$\therefore p_1 f(r_1) = r_1^2,$$

will be the equation to the locus required.

Ex. Let the spiral be the equiangular where $p = mr$;

$$\therefore r_1 = mr; \quad \therefore p_1 = \frac{r_1^2}{r} = \frac{mr_1^2}{r_1} = mr_1,$$

an equation to a similar spiral.

EXAMPLES.

Example. Find the value of p in the Conic Sections.

$$r = \frac{m}{1 + e\cos\theta}, \text{ where } m = \tfrac{1}{2} \text{ latus rectum};$$

$$\therefore u = \frac{1}{m} + \frac{e}{m}.\cos\theta; \quad \frac{du}{d\theta} = -\frac{e}{m}.\sin\theta;$$

$$\therefore u^2 + \frac{du^2}{d\theta^2} = \frac{1}{m^2}.\{1 + 2e\cos\theta + e^2\}$$

$$= \frac{1}{m^2}(2mu - 1 + e^2)$$

$$= \frac{1}{m^2}.\left\{\frac{2m - r(1 - e^2)}{r}\right\};$$

$$\therefore p^2 = \frac{m^2.r}{2m - r(1 - e^2)}.$$

(1) In parabola, $e = 1$; $\therefore\ p^2 = \frac{mr}{2}$, and $m = 2SA$;

$$\therefore\ SY^2 = SP \,.\, SA.$$

(2) In ellipse, $e < 1$; $m = \frac{b^2}{a}$; $1 - e^2 = \frac{b^2}{a^2}$;

$$\therefore\ p^2 = \frac{\frac{b^4}{a^2} \,.\, r}{2\frac{b^2}{a} - \frac{b^2}{a^2} \,.\, r} = \frac{b^2 r}{2a - r}.$$

(3) In hyperbola, $e^2 > 1$; $e^2 - 1 = \frac{b^2}{a^2}$;

$$\therefore\ p^2 = \frac{m^2 r}{2m + r(e^2 - 1)} = \frac{b^2 r}{2a + r};$$

and therefore in ellipse and hyperbola, $SY^2 = \frac{BC^2 \,.\, SP}{HP}$.

(4) Find the equation between p and r, when $\theta = \frac{a^n}{r^n}$,

$$\theta = \frac{a^n}{r^n} = a^n u^n; \quad \therefore\ u^n = \frac{\theta}{a^n};$$

$$\therefore\ \frac{du}{d\theta} = \frac{1}{na^n \,.\, u^{n-1}} = \frac{r^{n-1}}{b^n} \text{ by substitution;}$$

$$\therefore\ u^2 + \frac{du^2}{d\theta^2} = \frac{1}{r^2} + \frac{r^{2n-2}}{b^{2n}} = \frac{1}{r^2}\left\{\frac{b^{2n} + r^{2n}}{b^{2n}}\right\};$$

$$\therefore\ p = \frac{b^n \,.\, r}{\sqrt{b^{2n} + r^{2n}}}.$$

(5) Draw a tangent and asymptote to the spiral; where $\theta = \frac{a}{r} = au$; $\therefore\ \frac{1}{ST} = \frac{du}{d\theta} = \frac{1}{a}$; $\therefore\ ST = a$; or the locus of T is a circle radius $= a$.

Since ST is constant, and $\theta = 0$ when $r = \infty$. Produce SA indefinitely. Draw $ST \perp SA$ and $= a$. Then a line from T parallel to SP will be the asymptote required.



(6) If $r^2 = a^2 \cos 2\theta$ which is the polar equation to the Lemniscata; find the equation between p and r.

$$\text{Here } u^2 = \frac{1}{a^2 \cos 2\theta}; \quad \cos 2\theta = \frac{1}{u^2 a^2};$$

$$\frac{d\theta}{du} = \frac{1}{u^3 a^2 \sin 2\theta}; \quad \therefore \frac{du}{d\theta} = u^3 a^2 \sin 2\theta;$$

$$\text{and } \sin 2\theta = \sqrt{1 - \frac{1}{a^4 u^4}} = \frac{\sqrt{a^4 u^4 - 1}}{a^2 u^2};$$

$$\therefore \frac{du}{d\theta} = u\sqrt{a^4 u^4 - 1};$$

$$\therefore u^2 + \left(\frac{du}{d\theta}\right)^2 = u^2 + a^4 u^6 - u^2 = a^4 u^6;$$

$$\text{or } \frac{1}{p^2} = \frac{a^4}{r^6}; \quad \therefore p = \frac{r^3}{a^2}.$$

(7) Let $r = a^\theta$, the equation to the logarithmic spiral;

$$\therefore \frac{dr}{d\theta} = Aa^\theta = Ar; \quad \therefore \frac{d\theta}{dr} \text{ or } \frac{p}{r\sqrt{r^2 - p^2}} = \frac{1}{Ar};$$

$$\therefore \frac{p}{\sqrt{r^2 - p^2}} = \frac{1}{A} \text{ or } \frac{SY}{PY} = \tan SPY = \frac{1}{A};$$

that is, $\angle SPY$ is constant, and on this account the curve is called the equiangular spiral.

Cor. 1. Since $\frac{\sqrt{r^2 - p^2}}{p} = A; \quad \therefore \frac{r^2}{p^2} = 1 + A^2;$

$$\therefore \frac{p}{r} = \sin SPY = \frac{1}{\sqrt{1 + A^2}};$$

$$\therefore p = \frac{r}{\sqrt{1 + A^2}} = mr, \text{ by substitution.}$$

Cor. 2. The radii including equal angles are proportional.

Let SP and SP_1 include an $\angle a$,

and SQ and SQ_1 include the same angle.



Let $\angle ASP = \theta$, and $\angle ASQ = \phi$;

$$\therefore SP = a^\theta, \qquad SQ = a^\phi,$$

$$SP_1 = a^{\theta+\alpha}, \quad SQ_1 = a^{\phi+\alpha};$$

$$\therefore \frac{SP_1}{SP} = a^\alpha, \text{ and } \frac{SQ}{SQ} = a^\alpha;$$

$$\therefore \frac{SP}{SP_1} = \frac{SQ}{SQ_1}, \quad \text{or } SP : SP_1 :: SQ : SQ_1.$$

Cor. 3. Given the ratio of SP and SP_1, which include an angle α; find α.

Let $SP : SP_1 :: 1 : 1 + c$.

But $SP = a^\theta$, and $SP_1 = a^{\theta+\alpha}$;

$$\therefore \frac{SP_1}{SP} = 1 + c = a^\alpha;$$

$$\therefore \log(1 + c) = \alpha A = \frac{\alpha}{\tan\beta}; \quad \text{if } \beta = \angle SPY,$$

$$\text{or } \alpha = \tan\beta \,.\, \log(1 + c).$$

(8) If $r = a(1 + \cos\theta)$; find the equation between p and r. Ans. $p^2 = \frac{r^3}{2a}$.

(9) Find the equations between p and r; (1) when $r = a(e^{c+\theta} + e^{c-\theta})$; (2) when $r = a \sec n\theta$.

(1) $p = \frac{r^2}{\sqrt{2r^2 - 4ae^{2c}}}$; (2) $p = \frac{ar}{\sqrt{n^2r^2 - (n^2 - 1)a^2}}$.

(10) In the ellipse, if p be the perpendicular from the centre on the tangent, and r = distance of point from the centre, prove that

$$p^2 = \frac{a^2b^2}{a^2 + b^2 - r^2}.$$

CHAPTER XIII.

SINGULAR POINTS IN CURVES.

198. If in the equation to a curve expressed by $y = f(x)$, where y is the ordinate, and x the abscissa; some value of x as a makes any of the differential coefficients 0, $\frac{1}{0}$, or $\frac{0}{0}$, the point so determined is called a singular point.

(1) Let the values of the first differential coefficient be considered.

Since $\frac{dy}{dx}$ represents the tangent of the angle which the tangent makes with the axis of x, if $\frac{dy}{dx} = 0$, the tangent is parallel to the axis of x, and this circumstance generally indicates a maximum or minimum value of the ordinate.

If $\frac{dy}{dx} = \frac{1}{0}$, the tangent is perpendicular to the axis of x.

If $y = 0$ when $\frac{dy}{dx} = 0$, then the axis of x is a tangent to the curve at the origin.

If $x = 0$ when $\frac{dy}{dx} = \frac{1}{0}$, then the tangent passes through the origin, and is coincident with the axis of y.

When $\frac{dy}{dx} = \frac{0}{0}$. Many branches may pass through the point, as we shall see in the succeeding pages.

If $\frac{d^2y}{dx^2}$ have a real value when $\frac{dy}{dx} = 0$, the ordinate is a maximum or minimum, as in the annexed figures.





Before we proceed to investigate the values of $\frac{d^2y}{dx^2}$ at these points, we must establish the following proposition.

199. Prop. If the ordinate y be reckoned positive, a curve is convex or concave to the axis, according as $\frac{d^2y}{dx^2}$ is positive or negative.

In the annexed figures, let

$AN = x$, $NP = y$, $NN_1 = h$, and $y = f(x)$ be the equation to the curve.



Draw the tangent PM, its equation is

$$y_1 - y = \frac{dy}{dx}(x_1 - x).$$

Now at the point P_1, the equation to the curve becomes $N_1P_1 = f(x + h)$, or

$$N_1P_1 = y + \frac{dy}{dx}h + \frac{d^2y}{dx^2}\frac{h^2}{1.2} + \frac{d^3y}{dx^3}\frac{h^3}{2.3} + \&c.$$



and for the tangent, putting $x + h$ for x_1, and N_1M for y_1,

$$N_1M = y + \frac{dy}{dx}.h;$$

therefore the deflexion from the tangent, or MP_1

in figure (1) $= N_1M - N_1P_1 = -\frac{d^2y}{dx^2}\frac{h^2}{1.2} - \frac{d^3y}{dx^3}\frac{h^3}{2.3} - \&c.$

in figure (2) $= N_1P_1 - N_1M = +\frac{d^2y}{dx^2}\frac{h^2}{1.2} + \frac{d^3y}{dx^3}\frac{h^3}{2.3} + \&c.;$

and since h^2 is positive, and that h may be taken so small, that the first term of the expansion may be made greater than the sum of all the terms that follow it, the algebraical sign of MP_1 will depend upon that of $\frac{d^2y}{dx^2}$.

Therefore when the curve is concave to the axis, $MP_1 = -\frac{d^2y}{dx^2}\frac{h^2}{2} - \&c.$; and when convex to the axis, it

$= + \frac{d^2y}{dx^2}\frac{h^2}{1.2}$ + &c. Hence, (y being positive,) a curve is convex or concave to the axis, according as $\frac{d^2y}{dx^2}$ is positive or negative, or generally according as y and $\frac{d^2y}{dx^2}$ have the *same* or *different* signs.

Cor. If we suppose PT to be drawn $\perp$ to P_1N_1, $PT = h$,

$$\text{and } \frac{MP'}{PT^2} = \mp\tfrac{1}{2}\cdot\frac{d^2y}{dx^2} \mp \frac{d^3y}{dx^3}\cdot\frac{h}{2.3} + \text{&c.}$$

and if h be constantly diminished, the limit of the ratio of $MP' : PT^2$ will be $= \pm\frac{1}{2}\cdot\frac{d^2y}{dx^2}$.

Hence, ultimately, the deflexion from the tangent,

$$\text{or } MP_1 \propto \frac{d^2y}{dx^2}.$$

200. Sometimes the curve after being convex to the axis suddenly changes its curvature, and becomes concave, the point at which the change takes place is called a point of inflexion, or of *contrary flexure.*

If the tangent at this point be produced, one branch of the curve will be above, and the other below it, consequently on one side of the point in question $\frac{d^2y}{dx^2}$ will be positive, and on the other side, negative. Hence at the point itself $\frac{d^2y}{dx^2}$ must $= 0$, or ∞, for no quantity can change its sign without passing through zero or infinity.

There is not however a point of inflexion corresponding to every value of x, that makes $\frac{d^2y}{dx^2} = 0$, for not only must this equation be satisfied, but $\frac{d^2y}{dx^2}$ must change its sign after having passed through the point under consideration.

Also if the same value of x that makes $\frac{d^2y}{dx^2} = 0$, also makes $\frac{d^3y}{dx^3} = 0$, there may not be a point of contrary flexure.

For since $\frac{d^2y}{dx^2}$ is a function of x, write $x+h$ and $x-h$ for x, and then $\frac{d^2y}{dx^2}$ becomes, on these two suppositions, either

$$f(x+h) = \frac{d^2y}{dx^2} + \frac{d^3y}{dx^3}h + \frac{d^4y}{dx^4}\frac{h^2}{2} + \&c.;$$

$$\text{or } f(x-h) = \frac{d^2y}{dx^2} - \frac{d^3y}{dx^3}h + \frac{d^4y}{dx^4}\frac{h^2}{2} - \&c.$$

But at a point of inflexion $\frac{d^2y}{dx^2} = 0$; $\therefore$ the deflexions from the tangent at points $x+h$ and $x-h$ are respectively proportional to

$$+\frac{d^3y}{dx^3}h + \frac{d^4y}{dx^4}\frac{h^2}{2} + \&c., \text{ and } -\frac{d^3y}{dx^3}h + \frac{d^4y}{dx^4}\frac{h^2}{2} - \&c.,$$

which have contrary signs if $\frac{d^3y}{dx^3}$ do not $= 0$; but if $\frac{d^3y}{dx^3} = 0$, and $\frac{d^4y}{dx^4}$ does not vanish, the deflexions before and after the point will have the same algebraical sign, and the branches are both concave, or both convex, to the axis.

And hence in general there may be a point of contrary flexure, when the first differential coefficient which does not vanish is of an odd order.

Hence, to find whether a curve has a point of inflexion, put $\frac{d^2y}{dx^2} = 0$, or $\frac{1}{0}$, and if a be one of the values of x so determined, substitute $a+h$, and $a-h$ for x in the expression for $\frac{d^2y}{dx^2}$. Then if $\frac{d^2y}{dx^2}$ be affected with different signs, $x=a$ gives a point of contrary flexure.

Ex. 1. The cubical parabola $a^2y = x^3$.

$$y = \frac{x^3}{a^2}; \text{ and if } x = 0, \quad y = 0,$$

$$\frac{dy}{dx} = \frac{3x^2}{a^2},$$

$$\frac{d^2y}{dx^2} = \frac{6x}{a^2}.$$



If x be positive or negative, y and $\frac{d^2y}{dx^2}$ are positive or negative; the curve is therefore always convex to the axis.

$$\text{If } x = 0, \frac{d^2y}{dx^2} = 0.$$

$$\text{If } x = h, \frac{d^2y}{dx^2} = \frac{6h}{a^2} \text{ is positive.}$$

$$\text{If } x = -h, \frac{d^2y}{dx^2} = -\frac{6h}{a^2} \text{ is negative.}$$

The origin is therefore a point of contrary flexure; also, since $x = 0$, makes $\frac{dy}{dx} = 0$ and $y = 0$; the axis of x is a tangent to the curve.

Ex. 2. The Witch. $y = \frac{2a}{x}\sqrt{2ax - x^2}$,

$$\frac{dy}{dx} = 2a\left(\frac{\frac{ax - x^2}{\sqrt{2ax - x^2}} - \sqrt{2ax - x^2}}{x^2}\right) = \frac{-2a^2}{x\sqrt{2ax - x^2}},$$

$$\frac{d^2y}{dx^2} = 2a^2\frac{\sqrt{2ax - x^2} + \frac{ax - x^2}{\sqrt{2ax - x^2}}}{x^2(2ax - x^2)} = 2a^2\frac{(3a - 2x)}{x.(2ax - x^2)^{\frac{1}{2}}},$$

which $= 0$ if $x = \frac{3a}{2}$, and changes its algebraical sign, when $\frac{3a}{2} + h$ and $\frac{3a}{2} - h$ are successively put for x.

There are therefore two points of contrary flexure when

$$x = \frac{3a}{2}, \quad \text{and} \quad y = \pm\frac{2a}{\sqrt{3}}.$$

Ex. 3. Find the point of contrary flexure in the trochoid.

$$y = a(1 - e\cos\theta); \quad x = a(\theta - e\sin\theta);$$

$$\therefore \frac{dy}{d\theta} = ae\sin\theta; \quad \frac{dx}{d\theta} = a(1 - e\cos\theta);$$

$$\therefore \frac{dy}{dx} = \frac{e\sin\theta}{1 - e\cos\theta},$$

$$\frac{d^2y}{dx^2} = \frac{e\cos\theta(1 - e\cos\theta) - e\sin^2\theta}{(1 - e\cos\theta)^2} \cdot \frac{d\theta}{dx}$$

$$= \frac{e\cos\theta - e^2}{(1 - e\cos\theta)^2} \times \frac{1}{a(1 - e\cos\theta)}$$

$$= \frac{e(\cos\theta - e)}{a(1 - e\cos\theta)^3} = 0, \text{ if } \cos\theta = e;$$

and $\cos(\theta + h)$ is $< e$, and $\cos(\theta - h) > e$;

$\therefore$ $\cos\theta = e$ gives a point of contrary flexure,

and $y = a(1 - e^2) = a\left(1 - \frac{b^2}{a^2}\right) = \frac{a^2 - b^2}{a}$.

201. Points of contrary flexure in spirals.



Let there be two spirals, one concave and the other convex to the pole. Take two points P and P_1 in each near to each other, and draw SY and $SY_1 \perp$ to the tangents at P and P_1.

Let $SY = p$, $SP = r$, and $SP_1 = r + h$, and $p = f(r)$; therefore if Δ be the difference between SY_1 and SY, we have in figure (1), where the curve is concave to the pole,

$$\Delta = f(r + h) - f(r) = + \frac{dp}{dr} \cdot h + \frac{d^2p}{dr^2} \frac{h^2}{1 \cdot 2} + \&c.;$$

and in figure (2), where the spiral is convex to S,

$$\Delta = f(r) - f(r + h) = - \frac{dp}{dr} \cdot h - \frac{d^2p}{dr^2} \frac{h^2}{1 \cdot 2} - \&c.;$$

and as h may be taken so small that $\frac{dp}{dr} h$ may be greater than all the terms that follow, we see that the spiral is concave or convex to S, according as $\frac{dp}{dr}$ is positive or negative.

NOTE. P_1Y_1 in the figure should be a straight line.

Hence at a point of contrary flexure $\frac{dp}{dr} = 0$, and changes its sign immediately before and after the point.

EXAMPLE.

Let $r = a\theta^n$, find the point of contrary flexure.

$$\frac{d\theta}{dr} = \frac{1}{na\theta^{n-1}} = \frac{\theta}{nr} = \frac{\sqrt[n]{\frac{r}{a}}}{nr} = \frac{r^{\frac{1}{n}}}{na^{\frac{1}{n}}r}.$$

But $\frac{d\theta}{dr} = \frac{p}{r\sqrt{r^2 - p^2}}$; Art. 187, Cor. 1;

$$\therefore \frac{p}{\sqrt{r^2 - p^2}} = \frac{r^{\frac{1}{n}}}{na^{\frac{1}{n}}}; \quad \therefore \frac{r^2}{p^2} = 1 + \frac{n^2a^{\frac{2}{n}}}{r^{\frac{2}{n}}} = \frac{r^{\frac{2}{n}} + n^2a^{\frac{2}{n}}}{r^{\frac{2}{n}}};$$

$$\therefore p = \frac{r^{\frac{n+1}{n}}}{\sqrt{r^{\frac{2}{n}} + n^2a^{\frac{2}{n}}}};$$

$$\therefore \frac{dp}{dr} = \frac{n+1}{n} \cdot r^{\frac{1}{n}}\sqrt{r^{\frac{2}{n}} + n^2a^{\frac{2}{n}}} - r^{\frac{n+1}{n}} \frac{r^{-\frac{n-2}{n}}}{n\sqrt{r^{\frac{2}{n}} + n^2a^{\frac{2}{n}}}} = 0,$$

omitting the denominator;

$$\therefore (n+1)\, r^{\frac{1}{n}}(r^{\frac{2}{n}} + n^2a^{\frac{2}{n}}) - r^{\frac{3}{n}} = 0;$$

$$\therefore r^{\frac{1}{n}}\left\{nr^{\frac{2}{n}} + n^2(n+1)\, a^{\frac{2}{n}}\right\} = 0,$$

whence $r = 0$, and $r = a\left\{-n.(n+1)\right\}^{\frac{n}{2}}$.

If $\frac{n}{2}$ be a fraction with an even denominator, it is obvious that $n.(n+1)$ must be a negative number.

Let $n^2 + n = -p$; $\therefore n + \frac{1}{2} = \sqrt{\frac{1}{4} - p}$;

$\therefore n = -\frac{1}{2} \pm \sqrt{\frac{1}{4} - p}$; $\therefore p$ must never exceed $\frac{1}{4}$.

If $p = \frac{1}{4}$, $n = -\frac{1}{2}$, and $r = \frac{a}{\sqrt{\theta}}$, or $\theta = \frac{a^2}{r^2}$ the equation to the lituus.

MULTIPLE POINTS.

202. Whenever two or more branches of a curve pass through a point, it is called a multiple point; and a double, triple, or quadruple point, according as two, three, or four branches pass through it.

If the branches intersect, as in figure (1), which represents a double point, there will be at P two tangents, inclined at different angles to the axis, and thus $\frac{dy}{dx}$ will have two values corresponding to one of x or y.



Should however the branches pass through P, as in fig. 2. and touch each other, and the contact be only of the first order, there will be but one value of $\frac{dy}{dx}$; but as there are two deflexions from the tangent, there will be two values of $\frac{d^2y}{dx^2}$.



203. Problem. If $u = f(x, y) = 0$ be the equation to a curve, cleared of surds, and there be a point where two or more branches intersect, $\frac{dy}{dx} = \frac{0}{0}$ at that point.

Differentiate the equation, the result will be of the form

$$M \cdot \frac{dy}{dx} + N = 0, \text{ where } M = \frac{du}{dy} \text{ and } N = \frac{du}{dx}.$$

Then since two branches intersect, $\frac{dy}{dx}$ will have two values, but M and N will be the same for both. Let α and β be the two values of $\frac{dy}{dx}$;

$$\therefore M\alpha + N = 0, \text{ and } M\beta + N = 0;$$

$$\therefore M(\alpha - \beta) = 0, \text{ and } \alpha - \beta \text{ does not } = 0;$$

$$\therefore M = 0; \text{ and } \therefore N = 0, \text{ and } \frac{dy}{dx} = -\frac{N}{M} = \frac{0}{0}.$$

Cor. 1. Hence the value of p or $\frac{dy}{dx}$ may be found, by the same method as that by which the values of vanishing fractions are determined.

Thus, since $p = -\frac{N}{M} = \frac{0}{0}$ when $x = a$ and $y = b$; differentiating numerator and denominator,

$$p = -\frac{M_1 p + N_2}{M_1 + M_2 p}, \text{ where } M_1 = \frac{dN}{dy} = \therefore \frac{dM}{dx},$$

$$\text{or } M_2 p^2 + 2M_1 p + N_2 = 0;$$

a quadratic equation, from which two values of p may be found, and which, if possible, shew that the curve has a double point.

Cor. 2. If, however, $M_1 = 0$, $N_1 = 0$, and $M_2 = 0$, when $x = a$ and $y = b$, differentiate numerator and denominator a second time, and putting $q = \frac{d^2y}{dx^2}$, we shall have p of the form

$$p = -\frac{M_1 q + N_2 p^2 + N_3 p + N_4}{M_2 q + M_3 p^2 + M_4 p + M_5}$$

$$= -\frac{N_2 p^2 + N_3 p + N_4}{M_3 p^2 + M_4 p + M_5}, \text{ when } x = a \text{ and } y = b;$$

whence we have a cubic equation of the form

$$p^3 + ap^2 + bp + c = 0 \text{ to determine } p;$$

and if there be three possible roots, there is a triple point.

This process must be continued, if the numerator and denominator again vanish.

Ex. 1. Find the species of point at the origin of the curve,

$$ay^2 - x^3 - bx^2 = 0.$$

Differentiating and putting p for $\frac{dy}{dx}$,

$$2ayp - 3x^2 - 2bx = 0;$$

$$\therefore p = \frac{3x^2 + 2bx}{2ay} = \frac{0}{0}, \text{ if } x = 0 \text{ and } y = 0,$$

therefore there may be a multiple point.

Differentiating both numerator and denominator,

$$p = \frac{6x + 2b}{2ap} = \frac{2b}{2ap}, \text{ when } x = 0;$$

$$\therefore p^2 = \frac{b}{a}, \text{ and } p = \pm\sqrt{\frac{b}{a}};$$

$\therefore$ the origin is a double point, and the tangents cut the axis at angles, $\tan^{-1}\sqrt{\frac{b}{a}}$, and $\tan^{-1}\left(-\sqrt{\frac{b}{a}}\right)$.

This example will be useful in shewing another method by which multiple points may be found. Thus, if there be a surd quantity which disappears from the equation $y = f(x)$ by making $x = a$, but which is found in the equation $\frac{dy}{dx} = \phi(x)$, then $\frac{dy}{dx}$ will have two values, while y has but one, and there is a double point. For resuming the last equation, and solving it with respect to y,

$$y = \pm\frac{x}{\sqrt{a}}\sqrt{x+b}\,;\quad \frac{dy}{dx} = \pm\frac{\sqrt{x+b}}{\sqrt{a}} + \frac{x}{2\sqrt{a}\sqrt{x+b}}.$$

Make $x = 0$. Then $y = 0$, and $\frac{dy}{dx} = \pm\sqrt{\frac{b}{a}}$, as before.

Ex. 2. Find the point at the origin of the Lemniscata.

$$(x^2+y^2)^2 = a^2(x^2-y^2).$$

Here $2(x+py)\,.\,(x^2+y^2) = a^2(x-py)$;

$$\therefore\ p = \frac{a^2x - 2x(x^2+y^2)}{a^2y + 2y(x^2+y^2)} = \frac{0}{0},\ \text{if } x \text{ and } y = 0,$$

$$= \frac{a^2 - 2(x^2+y^2) - 2x(2x+2yp)}{a^2p + 2p(x^2+y^2) + 2y(x+2yp)}$$

$$= \frac{a^2}{a^2p},\ \text{if } x = 0 \text{ and } y = 0;$$

$$\therefore\ p^2 = 1,\ \text{and } p = \pm 1,$$

$$\text{or } \frac{dy}{dx} = \pm 1\,;\ \text{or } \frac{dy}{dx} = \tan 45,\ \text{and } \tan 135.$$

Ex. 3. Find the same, when $x^4 - ayx^2 + by^3 = 0$.

Here $4x^3 - ax^2p - 2ayx + 3by^2p = 0$;

$$\therefore\ p = \frac{2ayx - 4x^3}{3by^2 - ax^2} = \frac{0}{0},\ \text{when } x \text{ and } y = 0,$$

$$= \frac{2apx + 2ay - 12x^2}{6bpy - 2ax} = \frac{0}{0},\ \text{if } x = 0,$$

$$= \frac{2ap + 2axq + 2ap - 24x}{6bp^2 + 6byq - 2a}, \quad q = \frac{d^2y}{dx^2},$$

$$= \frac{4ap}{6bp^2 - 2a}, \text{ if } x = 0;$$

$$\therefore 6bp^3 - 2ap = 4ap, \text{ or } p.\{bp^2 - a\} = 0;$$

$$\therefore p = 0, \text{ and } p = \pm\sqrt{\frac{a}{b}},$$

there is a triple point at the origin, and the axis of x is one of the tangents.

The triple point at A is represented in the annexed figure; TAt is the axis of x, AT_1 and AT_2 are the tangents of the angles,

$$\tan^{-1}\sqrt{\frac{a}{b}}, \text{ and } \tan^{-1}\left(-\sqrt{\frac{a}{b}}\right).$$



Ex. 4. Find the same, when $y^3 - 3axy + x^3 = 0$.

$$y^2p - axp - ay + x^2 = 0;$$

$$\therefore p = \frac{ay - x^2}{y^2 - ax} = \frac{0}{0}, \text{ if } x = 0 \text{ and } y = 0;$$

$$\therefore p = \frac{ap - 2x}{2yp - a} = \frac{ap}{2yp - a}; \text{ if } x = 0;$$

$$\therefore 2yp^2 - ap = ap,$$

or $p(yp - a) = 0;$

$$\therefore p = 0, \text{ and } p = \frac{a}{y} = \frac{a}{0} = \infty.$$



The origin is therefore a double point, and the two axes are the tangents. The curve is represented above.

204. If the branches touch, then $\frac{dy}{dx}$ will have but one value, and yet at the same time be of the form $\frac{0}{0}$.

For supposing the contact to be of the n^{th} order between two branches of the curve; then the values of the differential coefficients, as far as the $(n + 1)^{\text{th}}$ coefficient, when $x = a$, and

$y = b$, will be the same for both branches; but after the n^{th} they will be different.

Let $M\frac{dy}{dx} + N = 0$ be the equation after the first differentiation, the original equation being previously freed from surd quantities.

Then, repeating the differentiation (n) times, we have

$$M \cdot \frac{d^{n+1}y}{dx^{n+1}} + N_1 = 0,$$

M being the same as before, and N_1 being the sum of the differential coefficients below the $(n+1)^{\text{th}}$, together with functions of x or y.

But $\frac{d^{n+1}y}{dx^{n+1}}$ has two values, as α and β, while M and N_1 remain unchanged; $\therefore M.(\alpha - \beta) = 0$; $\therefore M = 0$.

But $M\frac{dy}{dx} + N = 0$; $\therefore N = 0$; and $\therefore \frac{dy}{dx} = \frac{0}{0}$.

The analytical character of *double* points of this description is, that when $\frac{dy}{dx} = \frac{0}{0}$ has but one value, $\frac{d^2y}{dx^2}$, which also $= \frac{0}{0}$, has two.

CONJUGATE OR ISOLATED POINTS.

205. Conjugate or isolated points are those which have a real existence, and are determined by the equation to the curve; but from which no branches extend.

Hence if $x = a$ and $y = b$ give such a point, then $x = a + h$, and $x = a - h$, will make y, $\frac{dy}{dx}$, and the other differential coefficients, impossible.

$$\text{Also } \because y_1 = f(x+h) = y + \frac{dy}{dx}h + \frac{d^2y}{dx^2}\frac{h^2}{1.2} + \&c.$$
$$+ \frac{d^ny}{dx^n}\frac{h^n}{1\ldots n} + \&c.$$

is impossible, and that y and h are possible quantities, it is evident that some one of the differential coefficients is impossible, when $x = a$, and $y = b$.

PROP. At a conjugate point, if the equation be freed of surds, $\frac{dy}{dx} = \frac{0}{0}$.

For differentiating the equation, $u = f(xy) = 0$, we have

$$M\frac{dy}{dx} + N = 0 \ldots\ldots\ldots\ldots\ldots(1),$$

and if $\frac{dy}{dx}$ be not impossible, let $\frac{d^n y}{dx^n}$ be impossible;

$\therefore$ continue the differentiation of (1) $(n-1)$ times;

$$\therefore M\frac{d^n y}{dx^n} + N_1 = 0.$$

$$\text{Let } \frac{d^n y}{dx^n} = \alpha + \beta\sqrt{-1};$$

$$\therefore M\alpha + N_1 + M\beta\sqrt{-1} = 0;$$

$$\therefore M = 0; \quad \therefore \text{ from (1) } N = 0; \quad \therefore \frac{dy}{dx} = \frac{0}{0},$$

and the values of $\frac{dy}{dx}$ may in general be found, if any, by the method used for finding multiple points.

Ex. 1. $ay^2 - x^3 + bx^2 = 0$; $\therefore 2ayp - 3x^2 + 2bx = 0$;

$$\therefore p = \frac{3x^2 - 2bx}{2ay} = \frac{0}{0}, \text{ if } x = 0 \text{ and } \therefore y = 0$$

$$= \frac{6x - 2b}{2ap} = -\frac{b}{ap}, \text{ if } x = 0;$$

$$\therefore p^2 = -\frac{b}{a}; \text{ and } \therefore p = \sqrt{\frac{-b}{a}}.$$

Now $x = 0$ gives $y = 0$, while $p = \sqrt{\frac{-b}{a}}$. Also since $y = x\sqrt{\frac{x-b}{a}}$, if $x = 0 \pm h$, the values of y are impossible, and the origin is therefore a conjugate point. The same result may be obtained by differentiating the equation

$$y = x\sqrt{\frac{x-b}{a}}.$$

For $\frac{dy}{dx} = \sqrt{\frac{x-b}{a}} + \frac{x}{2\sqrt{a}\sqrt{x-b}} = \sqrt{\frac{-b}{a}}$; if $x = 0$.

206. The comparison of this example with Ex. 1. in multiple points will serve to explain the origin of conjugate points. In the curve $ay^2 - x^3 - bx^2 = 0$; two branches pass through the origin and meet at a point $x = -b$, forming an oval, while in the curve $ay^2 - x^3 + bx^2 = 0$, the oval disappears, and no curve exists between the values of $x = 0$, and $x = b$; these cases are represented in the annexed figures.



These two examples will shew that points of this kind arise from the vanishing of certain portions of the curve, owing to the change in the value of the constants.

Ex. 2. $y - b = (x-a)^2\sqrt{x-c}$; $a < c$.

If $x = a$, $y = b$; but if $x = a \pm h$, h being very small, so that $a + h$ is $< c$; y is impossible; $\therefore$ $x = a$, $y = b$, determines a conjugate point.

In this example we shall find $\frac{dy}{dx} = 0$; but if $q = \frac{d^2y}{dx^2}$; and if the equation be freed of surds,

$$q = \frac{4(x-a)^3 + 6(x-a)^2(x-c)}{y-b} = \frac{0}{0}; \quad x = a, \quad y = b;$$

$$= \frac{18(x-a)^2 + 12(x-a)(x-c)}{p} = \frac{0}{0}$$

$$= \frac{48(x-a) + 12(x-c)}{q} = \frac{12(a-c)}{q};$$

$$\therefore q^2 = 12(a-c) = -12(c-a);$$

$$\therefore q = 2\sqrt{3(c-a)}\sqrt{-1};$$

whence we see that $y_1 = f(x+h)$ is impossible.

207. In general we may remark, if there be a surd which vanishes from the equation $y = f(x)$ if $x = a$, but which becomes impossible in $\frac{d^n y}{dx^n} = \phi(x)$, there will be a conjugate point; if, at the same time, the values of y are impossible, both before and after the point.

CUSPS.

208. When two branches of a curve touch each other at a point through which the branches do not extend, the point is called a Cusp.

The branches have at this point but one tangent, and the cusp is said to be of the *first species* when the branches lie on opposite sides of the common tangent, and of the *second species* when they lie upon the same side.

Hence at such a point if $x = a$, $\frac{dy}{dx}$ will have but one value; but if either $a + h$, or $a - h$ be put for x, $\frac{d^2y}{dx^2}$ will have two values.

If the values of $\frac{d^2y}{dx^2}$ be both positive or both negative, the cusp is of the second species; but if one value be positive and the other negative, the cusp is of the first species, for the deflexion of the tangent from the curve is measured by $\frac{d^2y}{dx^2}$.

Since by the definition the branches suddenly stop at the cusp, either $a + h$, or $a - h$, put for x, will make the ordinate and the differential coefficients impossible.

Figures (1) and (2) exhibit cusps of the first and second species.



Sometimes the cusp is of the form below,



in which $(a + h)$ and $(a - h)$ put for x give real values for the ordinate.

These are discoverable by observing, that if $x = a$ and $y = b$ give the point P, that $y = b - k$ makes x impossible. Or we may transform the equation to the axis of y making $x = f^{-1}(y)$; and find the values of x, $\frac{dx}{dy}$, and $\frac{d^2x}{dy^2}$ at and near the point where $y = b$. Ex. $(y-2)^3 = (x-1)^2$.

Ex. 1. The semi-cubical parabola.

$$ay^2 = x^3, \quad y = \frac{x^{\frac{3}{2}}}{\sqrt{a}},$$

$$\frac{dy}{dx} = \frac{3}{2}\sqrt{\frac{x}{a}},$$

$$\frac{d^2y}{dx^2} = \frac{3}{2}\cdot\frac{1}{2}\cdot\frac{1}{\sqrt{ax}}.$$



If $x = 0$, y and $\frac{dy}{dx} = 0$, if $x = -h$, they are both impossible.

But if $x = h$, y and $\frac{d^2y}{dx^2}$ have two values, one positive and the other negative; the axis of x is therefore a tangent: there are two branches to the curve, one above and the other below the axis of x, and both convex to it, but the curve does not extend through the origin to the negative axis of the abscissas. The origin is a cusp of the first species.

Ex. 2. Find the point, when $x = a$ in the curve of which the equation is $y = b + cx^2 + (x-a)^{\frac{5}{2}}$;

$$\therefore \frac{dy}{dx} = 2cx + \frac{5}{2}(x-a)^{\frac{3}{2}},$$

$$\text{and } \frac{d^2y}{dx^2} = 2c + \frac{5\cdot 3}{2\cdot 2}(x-a)^{\frac{1}{2}}.$$

Let $x = a$; $\quad \therefore y = b + ca^2$,

$$\frac{dy}{dx} = 2ca; \quad \frac{d^2y}{dx^2} = 2c.$$



If $x = a + h$; $y = b + c(a+h)^2 + h^{\frac{5}{2}}$,

$$\frac{dy}{dx} = 2c(a+h) + \tfrac{5}{2} \cdot h^{\frac{3}{2}}; \quad \frac{d^2y}{dx^2} = 2c + \frac{15}{4} h^{\frac{1}{2}};$$

whence in consequence of the index $\frac{1}{2}$, y and $\frac{d^2y}{dx^2}$ have each two values, and those of $\frac{d^2y}{dx^2}$ are both positive, and since $a - h$ put for x makes y, $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$ impossible, the point is a cusp, and it is of the second species.

Take $x = a$, $y = b + ca^2$, and draw a tangent through P inclined to the axis of abscissas at an $\angle = \tan^{-1} 2ca$, and then draw two branches above the line through two points Q and Q_1, where

$$MQ = b + c(a+h)^2 + h^{\frac{5}{2}}, \quad \text{and} \quad MQ_1 = b + c(a+h)^2 - h^{\frac{5}{2}},$$

and the curves will be represented.

209. There are also sometimes to be met with, curves, in which a branch having reached a certain point, is suddenly arrested, and extends only on one side of the point, such a point may be called a *stop* point, "*Point d'arrêt.*"

Again, two or more distinct branches sometimes meet at a point, but do not pass through it, nor mutually touch there, each branch having a different tangent. Such points may be named shooting points, *points saillants.*

It is obvious that in the former species of point, $y = f(x)$ suddenly changes its value, or from being real, becomes impossible, and in the latter $\frac{dy}{dx}$ undergoes a similar change, i. e. from one value to another very different value. These are of rare occurrence, but as instances, the following are examples.

Ex. 1. The curve where $y = x \log x$, has a stop point at the origin.

For $x = +0$; $\quad y = 0$;

$x = -0$; $\quad y = \infty$,

Ex. 2. $y = x \tan^{-1}\frac{1}{x}$;

$$\frac{dy}{dx} = \tan^{-1}\frac{1}{x} - \frac{x}{x^2+1}.$$

If $x = +0$; or -0; $y = 0$;

$$\frac{dy}{dx} = \frac{\pi}{2}; \text{ or } -\frac{\pi}{2}.$$

Hence the origin is a shooting point, the tangents being inclined at angles

$$\tan^{-1}(1.5708), \text{ and } \tan^{-1}(-1.5708).$$

210. We shall conclude this Chapter by a few remarks upon the subject of tracing curves by means of their equations.

(1) If it be possible, let the equation be solved with respect to one of the unknown quantities as y, and let it be put under the form $y = f(x)$.

Then give to x all the possible *positive* values the equation admits of, and so determine the branches above and below the axis of *positive* abscissas.

Next put $(-x)$ for x in the equation $y = f(x)$, and in the equation, thus transformed, again substitute for x all its possible *positive* values, and the branches above and below the axis of *negative* abscissas will be determined.

(2) Ascertain whether the curve has asymptotes, and if it has, draw them.

(3) Find whether the branches be concave or convex to the axis, and determine the nature and situation of the singular points.

These remarks refer to curves having rectangular coordinates, but if the equation be between r and θ, give to θ, values from $\theta = 0$ to $\theta = 2\pi$, and draw the corresponding values of r; the positive values of r at the angles denoted by θ; the negative values, in a directly opposite direction, or separated from the positive by the angle π. Sometimes it may be necessary to take the negative values of θ.

Ex. 1. Let $y = x\left(\frac{x-2a}{x-a}\right)$, trace the curve.

A, the origin, Ax and Ay the two axes.



Let $x = 0$; $\therefore\ y = 0$,

$x < a$; $\therefore\ y$ is positive,

$x = a$, $y = \infty$,

$x > a < 2a$, y is negative,

$x = 2a$, $y = 0$,

$x > 2a$, y is positive,

$x = \infty$, y is ∞.

Again, let $-x$ be put for x;

$$\therefore\ y = -x\,\frac{x+2a}{x+a}$$ is always negative.

To draw the asymptote:

$$y = x \cdot \left(\frac{1-\frac{2a}{x}}{1-\frac{a}{x}}\right) = x\left(1-\frac{2a}{x}\right)\cdot\left\{1+\frac{a}{x}+\&c.\right\};$$

$$\therefore\ y = x\left\{1-\frac{a}{x}-\frac{2a^2}{x^2}-\&c.\right\}$$

$$= x - a - \frac{2a}{x} - \&c.;$$

$\therefore\ y = x - a$ is the equation to the asymptote.

Take $\therefore\ AB = a = AD$, and the line BD produced is the

asymptote. Also take $AC = 2a$. Then since $y = 0$, both when $x = 0$ and $x = 2a$, the curve cuts the axis at A and C.

Between A and B the curve is above the axis; at B the ordinate is infinite; from B to C the curve is below, from C to infinity it is above Ax. Again, since if x be negative, y is negative; the branch on the left of A is entirely below the axis.

To find the values of $\frac{dy}{dx}$;

$$\frac{dy}{dx} = \frac{x-2a}{x-a} + x \cdot \frac{(x-a)-(x-2a)}{(x-a)^2}$$

$$= \frac{x^2-3ax+2a^2}{(x-a)^2} + \frac{ax}{(x-a)^2} = \frac{x^2-2ax+2a^2}{(x-a)^2}.$$

Let $x = a$; $\therefore \frac{dy}{dx} = \infty$; and the infinite ordinate through B is an asymptote; if $x = 0$; $\frac{dy}{dx} = 2$, or angle at which the curve cuts the axis at A is $\tan^{-1}(2)$; if $x = 2a$, $\frac{dy}{dx}$ again $= 2$, or $\angle$ at which the curve cuts the axis at C is $= \angle$ at A.

Since if $(x^2 - 2ax + 2a^2)$ or $(x-a)^2 + a^2 = 0$, the values of x are impossible; there is no maximum or minimum ordinate.

Again, $\frac{d^2y}{dx^2} = \frac{2.(x-a)^2 - 2\{(x-a)^2 + a^2\}}{(x-a)^3} = \frac{-2a^2}{(x-a)^3}$;

$$\therefore \frac{d^2y}{dx^2} \text{ is } + \text{ if } x < a, \text{ and is } - \text{ if } x > a.$$

But $x < a$, y is $+$, and $x > a < 2a$, y is $-$; and $x > 2a$, y is $+$; therefore from A to B, and from B to C the curve is convex, and from C concave to the axis.

If x be $-$, $\frac{d^2y}{dx^2} = \frac{2a^2}{(x+a)^3}$ is $+$, but y is $-$; therefore the branch from A to the left hand is concave to the axis.

Ex. 2. Let $y^2 = \frac{x^3+1}{x-1} = \frac{x+1}{x-1}(x^2 - x + 1)$;

$$\therefore y = \pm \sqrt{\frac{x+1}{x-1}} \cdot \sqrt{x^2 - x + 1}.$$

If $x = 0$, $y = \frac{1}{\sqrt{-1}}$ is impossible,

$x < 1$, y is impossible,

$x = 1$, y is $\pm \infty$,

$x > 1$, y is possible $\pm$,

$x = \infty$, y is $\infty \pm$;

therefore there are two infinite branches extending above and below the axis of positive abscissas.

For x put $-x$; $\therefore y = \pm \sqrt{\frac{x-1}{x+1}(x^2 + x + 1)}$, which is impossible, if x be < 1; and $= 0$, if $x = 1$.



If $x > 1$, and increase to infinity, y is possible $\pm$ and increases to infinity; therefore there are two infinite branches which meet the axis Ax_1 in a point C, if $AC = 1$.

To find the asymptote:

$$y = \pm \sqrt{\frac{1 + \frac{1}{x}}{1 - \frac{1}{x}}} \cdot x \sqrt{1 - \frac{1}{x} + \frac{1}{x^2}}$$

$$= \frac{1 + \frac{1}{2x} + \&c.}{1 - \frac{1}{2x} + \&c.} x\left(1 - \frac{1}{2x} + \frac{1}{2x^2} + \&c.\right)$$

$$= \pm \left\{1 + \frac{1}{x} + \&c.\right\} x\left(1 - \frac{1}{2x} + \&c.\right)$$

$$= \pm x \left\{1 + \frac{1}{2x} + \frac{A}{x^2} + \&c.\right\};$$

$\therefore\ y = \pm\left(x + \frac{1}{2}\right)$ gives the two asymptotes.

Take $AD = AD_1 = \frac{1}{2}$, and $Ac = \frac{1}{2}$. Join cD and cD_1, these lines are the asymptotes, and if through B an infinite ordinate be drawn, two branches of the curve will lie within the angular spaces formed by the intersections of this line with cD and cD_1 produced. For these branches of the curve will always lie above the asymptotes, since the ordinate of the asymptote is always less than the ordinate of the curve. This may be thus shewn.

Let y_1 be the ordinate of the asymptote;

$$\therefore\ y^2 = \frac{x^3 + 1}{x - 1}, \quad \text{and} \quad y_1^2 = x^2 + x + \tfrac{1}{4};$$

$$\therefore\ y^2 - y_1^2 = \frac{x^3 + 1 - (x - 1).(x^2 + x + \frac{1}{4})}{x - 1},$$

$$\text{or}\ \ (y + y_1)(y - y_1) = \frac{3x + 1}{4.(x - 1)}.$$

But $y + y_1$ is $+$; $\therefore\ y - y_1$ is $+$; or $y > y_1$.

Similarly it may be shewn that the branches which extend from C, above and below the axis Cx_1, lie between the lines D_1c, and Dc (asymptotes) produced.

To find the values of $\frac{dy}{dx}$.

$$2 \log y = \log (x^3 + 1) - \log (x - 1);$$

$$\therefore\ \frac{dy}{dx} = \frac{y}{2} \cdot \left(\frac{3x^2}{x^3 + 1} - \frac{1}{x - 1}\right) = \frac{2x^3 - 3x^2 - 1}{2(x - 1)^{\frac{3}{2}} \cdot \sqrt{x^3 + 1}},$$

which is ∞, if $x = 1$, or $x = -1$.

Hence the infinite ordinate through B touches the curve at an infinite distance from Ax, or is an asymptote; and the curve at C where $y = 0$ cuts the axis at right angles. Also since the

numerator $2x^3 - 3x^2 - 1$ is $= -2$ when $x = 1$, and is $= 3$ when $x = 2$, there is some value of x between 1 and 2 which will make $\frac{dy}{dx} = 0$, or y a minimum. Take AM this value, and MP and Mp will be minimum ordinates.

Ex. 3. As a last example, let the equation be between polar co-ordinates, and let $r = a(1 + \cos\theta)$.



$$\theta = 0; \quad \therefore\ r = a(1 + 1) = 2a,$$

$$\theta = \frac{\pi}{2}; \quad \therefore\ r = a.$$

Let $\theta = \frac{\pi}{2} + \alpha; \quad \therefore\ \cos\theta = -\sin\alpha,$

and $r = a(1 - \sin\alpha)$, or $r < a$,

$$\alpha = \frac{\pi}{2}, \text{ or } \theta = \pi; \quad \therefore\ r = 0.$$

Again, let $\theta = (\pi + \alpha); \quad \therefore\ \cos\theta = -\cos\alpha,$

and $r = a(1 - \cos\alpha)$, which increases as α increases,

and $r = a$ when $\alpha = 90$.

Again, let $\theta = \frac{3\pi}{2} + \alpha; \quad \therefore\ \cos\left(\frac{3\pi}{2} + \alpha\right) = +\sin\alpha;$

$\therefore\ r = a(1 + \sin\alpha)$, which increases as α increases, and when $\alpha = \frac{\pi}{2}$, or $\theta = 2\pi$, $r = 2a$.

It is obvious that the curves in the first and fourth quadrants are the same, and also those in the second and third quadrants. If $-\theta$ be put for θ, $\because\ \cos(-\theta) = \cos\theta$: precisely the same curve will be produced. Take $AB = 2a$, $AC = AD = a$, and the points at which it cuts the axes are determined.—This curve is the *Cardioid.*

EXAMPLES.

(1) If $y = ax + bx^2 + cx^3$; there is a point of inflexion if $x = -\frac{b}{3c}$.

(2) If $y = x\sqrt{\frac{a^2 - x^2}{a^2 + x^2}}$, trace the curve, find its greatest ordinates, and the angles at which it cuts the axis of x: $x = a\sqrt{\sqrt{2} - 1}$; the angles are 45^0 and 90^0.

(3) Trace the curves defined by the three equations

$$y = x\sqrt{\frac{x^2 - a^2}{x^2 + a^2}}; \quad y = x\sqrt{\frac{x^2 + a^2}{x^2 - a^2}}; \quad y = x\sqrt{\frac{a^2 + x^2}{a^2 - x^2}}.$$

In (1) origin is a conjugate point, two rectilinear asymptotes pass through the origin, and two infinite branches meet the axis of x at $\pm a$. In (2) there are also two rectilinear asymptotes perpendicular to the axis of x: and the branches of the curve are included within the asymptotes. In (3) there are only asymptotes perpendicular to the axis, the branches of the curve pass through the origin and do not extend beyond the asymptotes.

(4) If $y = e^{\cos x}$, there are two points of inflexion corresponding to $x = \cos^{-1}\left(\frac{-1 \pm \sqrt{5}}{2}\right)$.

(5) $y^2 = \frac{a^4}{a^2 + x^2}$; trace the curve: there are inflexions if $x = \pm\frac{a}{\sqrt{2}}$.

(6) $y = (x - 2)\sqrt{\frac{x - 9}{x}}$; trace the curve: there is a conjugate point if $x = 2$; and y is a minimum if $x = -\frac{3}{2}$.

(7) $y^3 = ax^2 - x^3$, a cusp of the first species at the origin, an inflexion if $x = a$; a maximum ordinate if $x = \frac{2a}{3}$.

(8) If $y = \frac{a^2 x}{a^2 + x^2}$, there are two points of flexure when $x = 0$, and $x = a\sqrt{3}$; the curve cuts the axis at 45^0, and the axis of x is an asymptote to the two infinite branches; there are maximum ordinates when $x = \pm a$.

(9) If $y^3 = x^3 - a^3$, trace the curve; there are two points of inflexion, when $x = 0$, and when $x = a$.

(10) Find the points of contrary flexure, in the companion to the cycloid, when $x = a(1 - \cos\theta)$; $y = a\theta$.

Two points when $x = a$; $y = \pm \frac{1}{2}\pi a$.

(11) Find the point of contrary flexure in the lituus.

$$r = a\sqrt{2}; \quad \theta = \tfrac{1}{2}.$$

(12) Shew that the curve defined by $a^2p = r^3$ has a point of inflexion at the origin.

(13) Determine the nature of the points, when $x = a$.

If (1) $(y - b)^3 = (x - a)^5$, (3) $(y - x)^5 = (x - a)^3$,

(2) $(y - b)^2 = (x - a)^5$, (4) $(y - x)^2 = (x - a)^5$,

find also the inclination of the tangents at the points to the axis.

(14) If $r = \dfrac{a\theta^2}{\theta^2 - 1}$, there is a point of contrary flexure when $r = \dfrac{3a}{2}$; there are two rectilinear asymptotes and an asymptotic circle, radius $= a$.

(15) If $x^4 - a^2x^2 + a^3y = 0$, there are two points of contrary flexure.

(16) Trace the curve $r = a(2\cos\theta \pm 1)$. The Trisectrix. Like the Cardioid, with the addition of an interior oval.

(17) Trace the curves $y = \sin x$, $y = \tan x$, $y = \sec x$. The circumference is supposed to coincide with the axis of x, and the ordinate is the sine, tangent, or secant. The resulting locus is called, the curve of sines, tangents, or secants.

(18) Trace the curves $r^2 = a^2 \sin 2\theta$, and $r = a \sin 2\theta$. In the former an oval in the 1st and 3rd quadrants: in the latter an oval in every quadrant.

(19) If $r = a \sin 3\theta$, there are six ovals.

(20) Trace $y^3 - 3axy + x^3 = 0$. Make $x = r\cos\theta, y = r\sin\theta$: an oval in the 1st quadrant, an asymptote cutting the axis of x at 135^0, two infinite branches in the 2nd and 4th quadrants.

(21) Trace $y^2a = x(a \pm x)^2$. The exterior branch is parabolic: the interior has an oval between $x = 0$ and $x = a$.

(22) Trace the curves, (1) $r = a\tan\theta$; (2) $r = 2a\tan\dfrac{\theta}{2}$;

and find the position of the asymptotes. Asymptotic subtangents are a and $4a$; (1) is included between vertical, (2) between horizontal asymptotes.

(23) Trace the curve $r = a\theta$, taking both positive and negative values of θ.

(24) Find the asymptotes, least ordinates, and points of flexure of $y^2x = (x + a)(x - b)^2$.

(25) $(x^2 + y^2)^2 = 4axy$; a double point at the origin, $p = 0$ and $= \infty$.

(26) $(x^2 + y^2)^3 = 4a^2x^2y^2$; a quadruple point at the origin.

(27) $y^4 + 2axy^2 - ax^3 = 0$; a triple point at the origin, $p = \pm \frac{1}{\sqrt{2}}$, and $= \infty$.

(28) Given the base of a triangle and the exterior angle equal to three times the interior and opposite angle at the base, find the equation to the curve, which is the locus of the vertex, draw its asymptote, and find its maximum ordinate.

(29) In the diameter AB of a circle, take a point C; draw a chord AP, and an ordinate PN, and CQ parallel to AP, meeting NP in Q: trace the curve which is the locus of Q.

(30) A rod PQ passes through a fixed point A, find the equation to the curve described by P, when Q moves in the circumference of a circle of given radius, and trace the curve.

(31) Two points start from the opposite extremities of the diameter of a circle in the same direction, the velocities are uniform, and in the ratio of 2 : 1; find the locus of the bisection of the chord which joins the position of the points, and its polar subtangent.

CHAPTER XIV.

CURVATURE AND OSCULATING CURVES.

211. When two curves, as QPQ_1, RPP_1, cut each other in the manner represented in the figure, the values of y and x are the same for both curves at the point of intersection; i.e. if $y = f(x)$ be the equation to the curve RPP_1, and $y = \phi(x)$ the equation to QPQ_1, and $AN = a$, and $NP = b$; the values a and b put for x and y will make the equations $b = f(a)$ and $b = \phi(a)$ true equations, and $\therefore f(a) = \phi(a)$.



212. But if for x, $a + h$, be written, (or as we shall put it, $x + h$,) the values of the ordinates of the two curves no longer become equal, and their difference, which is represented in the figure by P_1Q_1, is equal to the difference between $f(x + h)$ and $\phi(x + h)$, and will therefore be some function of h, and its value will depend upon the relations existing between the differential coefficients of $f(x)$ and $\phi(x)$.

For, let $y_1 = N_1P_1$, $y_2 = N_1Q_1$, $z = f(x)$, and $v = \phi(x)$;

$$\therefore\ y_1 = y + \frac{dz}{dx}h + \frac{d^2z}{dx^2}\frac{h^2}{1.2} + \frac{d^3z}{dx^3}\frac{h^3}{2.3} + \&c.,$$

$$\text{and}\ \ y_2 = y + \frac{dv}{dx}h + \frac{d^2v}{dx^2}\frac{h^2}{1.2} + \frac{d^3v}{dx^3}\frac{h^3}{2.3} + \&c.;$$

$$\therefore\ P_1Q_1 = \left(\frac{dv}{dx} - \frac{dz}{dx}\right)h + \left(\frac{d^2v}{dx^2} - \frac{d^2z}{dx^2}\right)\frac{h^2}{1.2} + \&c.;$$

or putting A_1, A_2, A_3, &c. A_n for the coefficients of h, h^2, h^3, &c., the distance Δ between the curves, or the difference between the ordinates, is represented by a series with ascending powers of h, so that

$$\Delta = A_1h + A_2h^2 + A_3h^3 + A_4h^4 + \&c. + A_nh^n + \&c.$$

213. First, let $A_1 = 0$; $\therefore \frac{dv}{dx} = \frac{dz}{dx}$, or the first differential coefficients are equal. But $\frac{dv}{dx}$ and $\frac{dz}{dx}$ represent

the trigonometrical tangents of the angles which the tangents of the two curves at the point P make with the axis of x. Hence at such a point the ordinates are equal, and the tangents are coincident. This is called a contact of the *first* order.

214. Let not only $A_1 = 0$, but $A_2 = 0$, then

$$f(x) = \phi(x), \quad \frac{df(x)}{dx} = \frac{d\phi(x)}{dx}, \quad \text{and} \quad \frac{d^2f(x)}{dx^2} = \frac{d^2\phi(x)}{dx^2}.$$

This is called a contact of the *second* order.

And in general the curves are said to have a contact of the n^{th} order when the first power of h, in the expression for Δ is h^{n+1}; i. e. when all the differential coefficients as far as the $(n+1)^{\text{th}}$ are respectively equal in both series.

215. To find the degree of contact which a proposed curve of given species has with a given curve of known dimensions.

Let $y = f(x)$ be the equation to the given curve, and $y_1 = \phi(x_1)$ the equation to the proposed curve, which is supposed to contain n arbitrary constants.

Then, to determine these n constants, we must have the n equations,

$$y = y_1, \quad \frac{dy}{dx} = \frac{dy_1}{dx_1}, \quad \frac{d^2y}{dx^2} = \frac{d^2y_1}{dx^2_1}, \quad \text{and} \quad \frac{d^{n-1}y}{dx^{n-1}} = \frac{d^{n-1}y_1}{dx_1{}^{n-1}};$$

or the contact must be of the $(n-1)^{\text{th}}$ order.

Thus, let it be required to find the degree of contact which a straight line may have with a given curve; we observe that the equation to the line is $y_1 = ax_1 + b$, and contains two constants a, b, or the contact may be of the first order.

And to determine the straight line which has a contact of the first order with a curve, $y = f(x)$.

Here $\frac{dy}{dx} = \frac{dy_1}{dx_1} = a$; $y = y_1$, and $x = x_1$;

$$\therefore\ y = ax + b, \quad \text{or } b = y - ax = y - \frac{dy}{dx}x;$$

therefore substituting for a and b, $y_1 = \frac{dy}{dx}x_1 + y - x\frac{dy}{dx}$;

or $y_1 - y = \frac{dy}{dx}(x_1 - x)$ the equation to the tangent; or the

tangent has a contact of the first order with the curve which it touches.

216. In the circle of which the equation is

$$R^2 = (x_1 - \alpha)^2 + (y_1 - \beta)^2$$

there are three arbitrary constants, the radius R and the co-ordinates of the centre α and β. The circle therefore may have a contact of the second order, and the constants may be determined by means of the equations

$$y = y_1, \quad \frac{dy}{dx} = \frac{dy_1}{dx_1}, \quad \text{and} \quad \frac{d^2y}{dx^2} = \frac{d^2y_1}{dx_1^2}.$$

The circle so found is called the *circle of curvature*, and its radius the radius of curvature of any point in a given curve.

For since the curvature in the same circle is uniform, while it varies inversely as the radius in different circles, and that curves are *geometrically* said to have the same curvature, when at a common point, they have the same tangent, and ultimately the same deflexion from the tangent, which conditions are both fulfilled by the circle that has a contact of the second order; this circle is assumed to be the proper measure of curvature, and curves are said to have the same or different curvature, according as the radii of these circles are the same or different, and the curvature in general

$$\propto \frac{1}{\text{radius of curvature}}.$$

The circle of curvature is also called the osculating circle.

217. To find the radius of curvature, and co-ordinates of the centre of the osculating circle to any proposed curve.

Let $y = f(x)$ be the equation to the given curve,

$R^2 = (x_1 - \alpha)^2 + (y_1 - \beta)^2$ the equation to the circle;

$\therefore$ differentiating twice, we have

$$\therefore\ 0 = (x_1 - \alpha) + (y_1 - \beta) \cdot \frac{dy_1}{dx_1} \ldots\ldots\ldots\ldots (1),$$

$$\text{and}\ \ 0 = 1 + \frac{dy_1^2}{dx_1^2} + (y_1 - \beta) \cdot \frac{d^2y_1}{dx_1^2} \ldots\ldots\ldots\ldots (2).$$

But $y = y_1$, $x = x_1$, $\frac{dy}{dx} = \frac{dy_1}{dx_1}$, and $\frac{d^2y}{dx^2} = \frac{d^2y_1}{dx_1^2}$;

$\therefore$ changing x_1 into x, and y_1 into y;

$$\therefore R^2 = (x-\alpha)^2 + (y-\beta)^2$$

$$= (y-\beta)^2 \cdot \left\{1 + \frac{dy^2}{dx^2}\right\} \text{ from.....(1)}$$

$$= \frac{\left(1 + \frac{dy^2}{dx^2}\right)^2}{\left(\frac{d^2y}{dx^2}\right)^2} \cdot \left(1 + \frac{dy^2}{dx^2}\right) \text{.....(2)}$$

$$= \frac{\left(1 + \frac{dy^2}{dx^2}\right)^3}{\left(\frac{d^2y}{dx^2}\right)^2};$$

$$\therefore R = \pm \frac{\left(1 + \frac{dy^2}{dx^2}\right)^{\frac{3}{2}}}{\frac{d^2y}{dx^2}} = \pm \frac{(1+p^2)^{\frac{3}{2}}}{q},$$

where $p = \frac{dy}{dx}$ and $q = \frac{d^2y}{dx^2}$.

This expression has two signs; but if we call the radius positive, when the curve is concave to the axis, or when q is negative; and if, on the contrary, when the curve is convex, or when q is positive, the radius be reckoned negative, we shall always have $R = \frac{(1+p^2)^{\frac{3}{2}}}{-q}$.

The co-ordinates α and β may be found from the equations

$$y - \beta = -\frac{1 + \frac{dy^2}{dx^2}}{\frac{d^2y}{dx^2}} = -\frac{1+p^2}{q},$$

$$\text{and } x - \alpha = -(y-\beta) \cdot \frac{dy}{dx} = \frac{1+p^2}{q} \cdot p;$$

and the circle is thus completely determined.

218. In the annexed figure, let AP be the given curve, PO the radius of curvature, and O therefore the centre of the osculating circle.



Also let $AN = x$; $NP = y$.

Then $An = a$, $nO = -\beta$;

$$\therefore\ PM = y - \beta = -\frac{(1+p^2)}{q},$$

$$OM = a - x = -\frac{(1+p^2)}{q}.p;$$

PM and OM are respectively called the semi-chords perpendicular and parallel to the axis of x; for if we describe the circle, of which the radius is OP and centre O, PM is half the chord of an arc, since OM is perpendicular to it, and OM is equal to half the chord drawn from P parallel to AN.

219. The point O changes its position with the change in the place of P, and traces out a curve, which is called the *evolute* of the original curve. Hence we may define the evolute to be the locus of the centre of the circle of curvature, and its co-ordinates are α and β.

And since from $y = f(x)$; p and q may be found in terms of y or x,

$$\text{and } \therefore \text{ from } y - \beta = -\frac{1+p^2}{q};\quad x - \alpha = \frac{1+p^2}{q}.p;$$

and $y = f(x)$; y and x may be eliminated; therefore there will arise an equation between α, β and constant quantities, which will be that of the evolute.

220. Since $x - \alpha + (y - \beta).\dfrac{dy}{dx} = 0$;

$$\therefore\ y - \beta = -\frac{dx}{dy}.(x - \alpha);$$

but this is the equation to the normal of the original curve, drawn from a point, of which the co-ordinates are x, y, and passing through a point whose co-ordinates are α and β. Hence the normal passes through the centre of the circle of curvature, and therefore the radius is coincident with the normal.

221. The radius of curvature is a tangent to the evolute.

Resuming the equation $(x-a)+(y-\beta).\frac{dy}{dx}=0.$

Differentiate it, considering y, β and a as functions of x;

$$\therefore\ 1-\frac{da}{dx}+(y-\beta).\frac{d^2y}{dx^2}+\frac{dy^2}{dx^2}-\frac{d\beta}{dx}.\frac{dy}{dx}=0.$$

$$\text{But}\ \ 1+\frac{dy^2}{dx^2}+(y-\beta).\frac{d^2y}{dx^2}=0;$$

$$\therefore\ -\frac{da}{dx}-\frac{d\beta}{dx}.\frac{dy}{dx}=0;\quad \therefore\ \frac{dy}{dx}=-\frac{\frac{da}{dx}}{\frac{d\beta}{dx}}=-\frac{da}{d\beta};$$

$$\therefore\ (x-a)-(y-\beta).\frac{da}{d\beta}=0;\quad \text{or}\ (\beta-y)=\frac{d\beta}{da}.(a-x),$$

which is the equation to a tangent drawn to a point, (β, a), and passing through a point, (y, x).

But $(\beta-y)=\frac{d\beta}{da}(a-x)$ is identical with

$$(y-\beta)=-\frac{dx}{dy}(x-a),$$

or with the equation to the normal of the original curve.

Hence the normal to the curve, i.e. the radius of curvature, is the tangent to the evolute.

222. To find the length (s) of the evolute.

Since $R^2=(x-a)^2+(y-\beta)^2.$

Differentiate, considering R, y, x, β as functions of a,

$$\therefore\ R\frac{dR}{da}=(x-a)\frac{dx}{da}+(y-\beta)\frac{dy}{da}-\left\{x-a+(y-\beta)\frac{d\beta}{da}\right\}$$

$$=-\left\{x-a+(y-\beta).\frac{d\beta}{da}\right\}$$

$$=-(x-a)\left\{1+\frac{d\beta^2}{da^2}\right\}.$$

$$\text{But}\ R^2=(x-a)^2\left\{1+\left(\frac{y-\beta}{x-a}\right)^2\right\}$$

$$=(x-a)^2\left\{1+\frac{d\beta^2}{da^2}\right\}\ldots\ldots\ldots\ldots(1).$$

And $R^2 \frac{dR^2}{da^2} = (x-a)^2 \cdot \left(1 + \frac{d\beta^2}{da^2}\right)^2$ .....(2).

Divide (2) by (1);

$$\therefore \frac{dR^2}{da^2} = 1 + \frac{d\beta^2}{da^2} = \frac{ds^2}{da^2};$$

$\therefore R = s + c$, c being some constant length.

Hence, if the equation to the curve be algebraical, R may be found in finite terms, and the length of the evolute found; or the evolutes of algebraic curves are rectifiable.

Cor. Let s_1, s_2, be two arcs of the evolute, from its commencement to the points where the radii are R_1 and R_2;

$$\therefore R_1 - s_1 = c; \quad R_2 - s_2 = c; \quad \therefore R_2 - R_1 = s_2 - s_1:$$

$$\text{let } s_2 - s_1 = a; \quad \therefore a = R_2 - R_1;$$

or the difference between two radii of curvature equals the length of the arc of the evolute intercepted by them.

223. From this property it is, that the curve has derived the name of evolute.

For if we take a string of constant length, one end of which is fastened at B, and the remainder is made to coincide with the curve COO_1B, then if the string be unwrapped or evolved from COO_1B, it will describe the curve APP_1.



COB is called the evolute, and

APP_1..... involute.

From this construction it is obvious,

(1) That the arc OO_1 is equal to $P_1O_1 - PO$.

(2) That O is the centre of a circle of which the radius OP is the radius of curvature to the point P.

(3) That PO is a tangent to the evolute.

(4) That PO is a normal to the involute.

224. Another, geometrical, method of finding the radius of curvature and the co-ordinates of the centre of the osculating circle is to assume that centre to be the limit of the intersections of two consecutive normals.

The truth of this assumption may be thus shewn:

$(y-\beta) = -\frac{dx}{dy}.(x-\alpha)$ is the equation to the normal,

$$\text{or } (x-\alpha) + (y-\beta).\frac{dy}{dx} = 0.$$

Now at the point of intersection, α and β remain the same for the two normals, while x, y and $\frac{dy}{dx}$ vary, since at a consecutive point, x and y become $x+dx$ and $y+dy$; therefore differentiating, considering α and β as constant,

$$1 + \frac{dy^2}{dx^2} + (y-\beta).\frac{d^2y}{dx^2} = 0.$$

The same equation has been before obtained to find the coordinate β of the centre, and α is then known from

$$x - \alpha = -(y-\beta)\frac{dy}{dx}.$$

And R may be found from the equation

$$R^2 = (x-\alpha)^2 + (y-\beta)^2.$$

225. Hence to find the radius of curvature in spirals.



AP the spiral, S the pole. PO a normal, and O the point of ultimate intersection of two consecutive normals. O is the centre of the circle of curvature.

$SP = r$, $PO = R$
$SY = p$, $SO = r_1$ $\Big\}$, $SN \perp$ on $PO = p_1$.

Now $SO^2 = SP^2 + PO^2 - 2PO.PN$,

or $r_1^2 = r^2 + R^2 - 2R.p$; for $PN = SY$.

Then since SO and OP remain constant, while

SP and SY vary, and since $p = f(r)$;

$$\therefore\ 0 = r - R\frac{dp}{dr};\quad \therefore\ R = r.\frac{dr}{dp}.$$

If OM be drawn $\perp$ to PS, or PS produced, then $PM = \frac{1}{2}$ the chord of curvature through S,

$$\text{and } PM = PO \times \frac{SY}{SP} = r.\frac{dr}{dp}.\frac{p}{r} = p.\frac{dr}{dp}.$$

226. To find R in terms of r and θ.

Since $\frac{1}{p^2} = \frac{1}{r^2} + \frac{1}{r^4} \cdot \frac{dr^2}{d\theta^2}$, Art. (186);

$$\therefore \frac{dp}{dr} \cdot \frac{1}{p^3} = \frac{1}{r^3} + \frac{2}{r^5} \cdot \frac{dr^2}{d\theta^2} - \frac{1}{r^4} \cdot \frac{d^2r}{d\theta^2};$$

$$\therefore \frac{dp}{rdr} \cdot \frac{1}{p^3} = \frac{1}{r^6}\left\{r^2 + 2 \cdot \frac{dr^2}{d\theta^2} - r \cdot \frac{d^2r}{d\theta^2}\right\} \ldots\ldots\ldots (1).$$

But $\frac{1}{p^2} = \frac{1}{r^4}\left(r^2 + \frac{dr^2}{d\theta^2}\right)$;

$$\therefore \frac{1}{p^3} = \frac{1}{r^6}\left(r^2 + \frac{dr^2}{d\theta^2}\right)^{\frac{3}{2}} \ldots\ldots\ldots\ldots\ldots\ldots (2);$$

therefore dividing (2) by (1),

$$\frac{rdr}{dp} = R = \frac{\left(r^2 + \frac{dr^2}{d\theta^2}\right)^{\frac{3}{2}}}{r^2 + 2 \cdot \frac{dr^2}{d\theta^2} - r \cdot \frac{d^2r}{d\theta^2}};$$

an expression for the radius of curvature, when $r = f(\theta)$.

Cor. Hence also to find $\frac{pdr}{dp}$ in terms of r and θ.

For $\therefore \frac{dp}{dr} \cdot \frac{1}{p^3} = \frac{1}{r^5}\left(r^2 + 2 \cdot \frac{dr^2}{d\theta^2} - r \cdot \frac{d^2r}{d\theta^2}\right) \ldots (1)$,

$$\frac{1}{p^2} = \frac{1}{r^4}\left(r^2 + \frac{dr^2}{d\theta^2}\right) \ldots\ldots\ldots\ldots (2);$$

therefore dividing (2) by (1),

$$\frac{pdr}{dp} = \tfrac{1}{2} \text{ chord} = \frac{r \cdot \left(\frac{dr^2}{d\theta^2} + r^2\right)}{r^2 + 2 \cdot \frac{dr^2}{d\theta^2} - r \cdot \frac{d^2r}{d\theta^2}}.$$

227. Evolutes to spirals.

The point O will trace out the evolute, and PO is always a tangent to it, and SN is perpendicular to PO, we must therefore find the relation between SO and SN.

$$\text{Now } r_1^2 = r^2 + R^2 - 2Rp \ldots\ldots\ldots(1),$$

$$\text{and } p_1 = PY = \sqrt{r^2 - p^2} \ldots\ldots\ldots(2),$$

$$\text{and } p = f(r) \ldots\ldots\ldots(3), \quad \text{and } R = r.\frac{dr}{dp} \ldots\ldots\ldots(4),$$

between these equations p, r and R may be eliminated, and the resulting equation will involve r_1, p_1, and constant quantities, which will be the equation required.

Ex. Let the spiral be the equiangular.

$$\text{Here } p = mr; \quad \therefore\ R = \frac{rdr}{dp} = \frac{r}{m},$$

$$p_1 = \sqrt{r^2 - p^2} = r\sqrt{1 - m^2};$$

$$\therefore\ r_1^2 = r^2 + R^2 - 2Rp = r^2 + \frac{r^2}{m^2} - 2r^2$$

$$= \frac{r^2}{m^2} - r^2 = \frac{r^2}{m^2}(1 - m^2) = \frac{p_1^2}{m^2};$$

$$\therefore\ p_1 = mr_1,$$

or the evolute is a spiral similar to the original, and described round the same pole S.

228. When two curves intersect, we have seen that the distance between them, measured along the ordinate is, (when x becomes $x + h$) expressed by the equation

$$\Delta = A_1h + A_2h^2 + A_3h^3 + A_4h^4 + A_5h^5 + \&c.$$

If therefore we put $(-h)$ for h, we shall have an expression for the distance between them at a point where the abscissa is $x - h$: let Δ_1 be this distance;

$$\therefore\ \Delta_1 = -A_1h + A_2h^2 - A_3h^3 + A_4h^4 - A_5h^5 + \&c.$$

Now since h may be taken so small that any one term shall exceed the sum of all that follow it; we observe *First*, that if $A_1 = 0$, Δ and Δ_1 have the same sign, or that in a contact of the first order, the curves touch, but do not intersect.

Thus the tangent does not cut the curve, unless $A_2 = 0$, or at a point of contrary flexure.

Secondly. Let both $A_1 = 0$ and $A_2 = 0$, or the contact be of the second order. Then

$$\Delta = A_3h^3 + A_4h^4 + \&c.$$

$$\Delta_1 = -A_3h^3 + A_4h^4 - \&c.,$$

which have different signs, and therefore if the osculating curve be below the given curve at a point where the abscissa is $x + h$, it will be above it at a point where x becomes $x - h$. Hence the circle of curvature both cuts and touches the curve.

There is however an exception to this, when the radius of curvature is a maximum or minimum; for then (as we shall see in the next article) $A_3 = 0$, and the expressions for Δ and Δ_1 have the same sign.

For if the contact be of the third order,

$$\Delta = A_4h^4 + A_5h^5 + \&c. \quad \Delta_1 = A_4h^4 - A_5h^5 + \&c.;$$

that is, Δ and Δ_1 have the same sign, and therefore the osculating curve does not cut the given curve.

Hence it appears that, when the contact is of an even order, the osculating curve both touches and cuts the given curve, but when the contact is of an odd order, it merely touches it.

229. Prop. When the radius of curvature is a maximum or minimum, the contact is of the third order, or $A_3 = 0$.

$$\text{Let } p = \frac{dy}{dx}, \quad q = \frac{d^2y}{dx^2}, \quad \text{and } r = \frac{d^3y}{dx^3}.$$

$$\text{Then } R = \frac{(1+p^2)^{\frac{3}{2}}}{-q}.$$

$$\text{But if } R \text{ be a maximum or minimum, } \frac{dR}{dx} = 0;$$

$$\therefore 3\sqrt{1+p^2}.p - \frac{(1+p^2)^{\frac{3}{2}}}{q^2}r = 0, \quad \text{or } r = \frac{3pq^2}{1+p^2}.$$

But $1 + p^2 + (y - \beta)q = 0$, and if there be a contact of the third order, we must differentiate this equation, and put the co-ordinates of the curve for those of the circle;

$$\therefore 2pq + pq + (y - \beta)r = 3pq - \frac{1+p^2}{q}r = 0;$$

$$\therefore r = \frac{3pq^2}{1+p^2}.$$

The same result as before, and therefore when A_3, or the difference between the third differential coefficient of $y = f(x)$, and of $R^2 = (x - \alpha)^2 + (y - \beta)^2$, equals 0, or when the contact is of the third order, the radius of curvature is either a maximum or minimum.

230. If $q = 0$, while p is not infinite, $R = \infty$; this is the case at a point of contrary flexure; for then the curve changes from convex to concave; the circle of curvature becomes a straight line, namely the tangent, and before and after the point the radius of curvature is measured in opposite directions.

EXAMPLES.

(1) Find the radius of curvature and evolute of the common parabola.

$$y^2 = 4ax, \quad \frac{dy}{dx} = \frac{2a}{y}, \quad \frac{d^2y}{dx^2} = -\frac{2a}{y^2}\frac{dy}{dx} = -\frac{4a^2}{y^3},$$

$$1 + p^2 = 1 + \frac{4a^2}{y^2} = \frac{4a^2 + y^2}{y^2} = \frac{4a\,(a+x)}{y^2}.$$

But $R = \dfrac{(1+p^2)^{\frac{3}{2}}}{-q} = \dfrac{\{4a\,.\,(a+x)\}^{\frac{3}{2}}}{4a^2} = \dfrac{2(a+x)^{\frac{3}{2}}}{\sqrt{a}} = \dfrac{2}{\sqrt{a}}\,.\,SP^{\frac{3}{2}}.$

But $(y - \beta) = \dfrac{1+p^2}{-q} = \dfrac{4a^2 + y^2}{y^2}\dfrac{y^3}{4a^2} = y\,.\,\dfrac{(a^2 + ax)}{a^2}$

$$= y + \frac{yx}{a};$$

$$\therefore\ -\beta = \frac{yx}{a} = \frac{y^3}{4a^2}; \quad \therefore\ -y = (4a^2\beta)^{\frac{1}{3}},$$

$$x - a = -p\,.\,(y - \beta) = -\frac{2a}{y}\frac{y\,(x+a)}{a} = -2\,(x+a);$$

$$\therefore\ 3x = a - 2a, \quad \text{or} \quad x = \left(\frac{a-2a}{3}\right).$$

But $\because\ y^2 = 4ax; \quad \therefore\ (4a^2\beta)^{\frac{1}{3}} = \dfrac{4a}{3}\,(a - 2a);$

$$\therefore\ \beta^2 = \frac{4}{27a}\,.\,(a-2a)^3 = \frac{4}{27a}x_1^3; \quad \text{if } a - 2a = x_1,$$

the equation to the semi-cubical parabola.

(2) In the Conic Sections, the radius of curvature $\propto$ (normal)^3.

$$\text{Normal} = N = y\sqrt{1+p^2}; \quad \therefore\ \sqrt{1+p^2} = \frac{N}{y};$$

$$\therefore\ R = \frac{(1+p^2)^{\frac{3}{2}}}{-q} = \frac{N^3}{-y^2 q}.$$

Now if the vertex be the origin, and the axis the axis of x,

$$y^2 = 2mx + nx^2;\quad \therefore\ yp = m + nx;$$

$$\therefore\ yq + p^2 = n;\quad \therefore\ y^3 q = ny^2 - p^2 y^2$$

$$= n(2mx + nx^2) - (m + nx)^2 = -m^2;$$

$$\therefore\ R = \frac{N^3}{m^2}.$$

(3) Find the radius of curvature of the ellipse.

$$y = \frac{b}{a}\sqrt{a^2 - x^2};\quad \therefore\ p = -\frac{b}{a}\frac{x}{\sqrt{a^2 - x^2}}$$

$$-q = \frac{ba}{(a^2 - x^2)^{\frac{3}{2}}},$$

$$1 + p^2 = 1 + \frac{b^2 x^2}{a^2(a^2 - x^2)} = \frac{a^4 - (a^2 - b^2)x^2}{a^2(a^2 - x^2)} = \frac{a^2 - e^2 x^2}{a^2 - x^2};$$

$$\therefore\ R = \frac{(a^2 - e^2 x^2)^{\frac{3}{2}}}{ba}.$$

Cor. Let R_1 be the radius at the vertex, and R_2 the radius at the extremity of the minor axis;

$$\therefore\ R_1 = \frac{(a^2 - a^2 e^2)^{\frac{3}{2}}}{ba} = \frac{b^2}{a},\quad \text{and}\quad R_2 = \frac{a^3}{ba} = \frac{a^2}{b};$$

therefore, the length of the evolute of the elliptic quadrant

$$= R_2 - R_1 = \frac{a^2}{b} - \frac{b^2}{a} = \frac{a^3 - b^3}{ab};$$

and $\therefore$ the length of the whole evolute $= \frac{4}{ab}.(a^3 - b^3)$.

If R be a maximum or minimum, $\frac{dR}{dx} = 0$;

$$\therefore\ -3\sqrt{a^2 - e^2 x^2}.e^2 x = 0;$$

$\therefore\ x = 0$, and $x^2 = \frac{a^2}{e^2}$; but $x = \frac{a}{e}$ or $> a$ is impossible,

and $\frac{d^2 R}{dx^2} = -3e^2\sqrt{a^2 - e^2 x^2} + 3e^2 x\frac{d}{dx}\sqrt{a^2 - e^2 x^2}$

$$= -3e^2 a,\quad \text{if } x = 0;$$

therefore R is a maximum, when $x = 0$, or $y = \pm b$.

Hence, at the extremities of the minor axis the circle of curvature touches the ellipse.

(4) To find the equation to the evolute.

$$y-\beta=\frac{1+p^2}{-q}=\frac{(a^2-e^2x^2)\sqrt{a^2-x^2}}{ba}=\frac{y(a^2-e^2x^2)}{b^2};$$

$$\therefore\ \beta=-y.\left\{\frac{a^2-e^2x^2}{b^2}-1\right\}=-\frac{ye^2}{b^2}(a^2-x^2)=-\frac{y^3e^2.a^2}{b^4};$$

$$\therefore\ \left(\frac{y}{b}\right)^3=-\frac{b\beta}{(ae)^2};\quad \therefore\ \left(\frac{y}{b}\right)^2=\frac{(b\beta)^{\frac{2}{3}}}{(ae)^{\frac{4}{3}}},$$

$$x-a=-y.\left(\frac{a^2-e^2x^2}{b^2}\right)\frac{dy}{dx}=y.\left(\frac{a^2-e^2x^2}{b^2}\right).\frac{b^2x}{a^2y}$$

$$=\frac{(a^2-e^2x^2)}{a^2}x=x-\frac{e^2x^3}{a^2};$$

$$\therefore\ \left(\frac{x}{a}\right)^3=\frac{-a}{ae^2};\quad \therefore\ \frac{x^2}{a^2}=\frac{(a)^{\frac{2}{3}}}{a^{\frac{2}{3}}e^{\frac{4}{3}}}=\frac{(aa)^{\frac{2}{3}}}{(ae)^{\frac{4}{3}}}.$$

But $\dfrac{x^2}{a^2}+\dfrac{y^2}{b^2}=1$; $\therefore\ \dfrac{(aa)^{\frac{2}{3}}}{(ae)^{\frac{4}{3}}}+\dfrac{(b\beta)^{\frac{2}{3}}}{(ae)^{\frac{4}{3}}}=1$;

$$\therefore\ (aa)^{\frac{2}{3}}+(b\beta)^{\frac{2}{3}}=(ae)^{\frac{4}{3}}=(a^2-b^2)^{\frac{2}{3}}.$$

(5) Radius of curvature and evolute of cycloid.

$$\left.\begin{matrix}AN=x\\NP=y\\AB=2a\end{matrix}\right\};\ \therefore\ \frac{dy}{dx}=\frac{\sqrt{2a-y}}{\sqrt{y}};$$



$$\therefore\ 1+\frac{dy^2}{dx^2}=\frac{2a}{y};$$

$$\therefore\ \frac{dy}{dx}.\frac{d^2y}{dx^2}=\frac{-a}{y^2}\frac{dy}{dx};$$

$$\therefore\ -\frac{d^2y}{dx^2}=\frac{a}{y^2};$$

$$\therefore\ R=\left(\frac{2a}{y}\right)^{\frac{3}{2}}.\frac{y^2}{a}=\frac{2a}{y}\sqrt{\frac{2a}{y}}.\frac{y^2}{a}=2\sqrt{2ay}=2RB.$$

NOTE—*AOD* in the figure should be a curve.

To find the evolute.

$$y - \beta = \frac{2a}{y}\frac{y^2}{a} = 2y; \quad \therefore \beta = -y,$$

$$x - \alpha = -(y-\beta)\frac{dy}{dx} = -2y\frac{\sqrt{2ay-y^2}}{y} = -2\sqrt{2ay-y^2};$$

$$\therefore \alpha = x + 2\sqrt{2ay-y^2};$$

$$\therefore \frac{d\alpha}{dx} = 1 + \frac{2(a-y)}{\sqrt{2ay-y^2}}\cdot\frac{dy}{dx} = 1 + \frac{2(a-y)}{y} = \frac{2a-y}{y},$$

$$\text{and } -\frac{d\beta}{dx} = \frac{dy}{dx} = \sqrt{\frac{2a-y}{y}};$$

$$\therefore -\frac{d\alpha}{d\beta} = \sqrt{\frac{2a-y}{y}} = \sqrt{\frac{2a-(-\beta)}{(-\beta)}}.$$

Take $Am = x_1 = -\beta$, and $mO = \alpha = y_1$;

$$\therefore -\frac{d\alpha}{d\beta} = \frac{dy_1}{dx_1} = \sqrt{\frac{2a-x_1}{x_1}} = \frac{\sqrt{2ax_1 - x_1^2}}{x_1}.$$

The equation to a cycloid, of which the vertex is A, and the diameter of the generating circle $= 2a$.

(6) Find the chords of curvature drawn through the centre and focus of an ellipse.

Since if $CP = r$, and CY, $\perp$ to tangent $= p$,

$$p^2 = \frac{a^2b^2}{a^2+b^2-r^2};$$

$$\therefore 2\log p = \log a^2b^2 - \log(a^2+b^2-r^2);$$

$$\therefore \frac{dp}{pdr} = \frac{r}{a^2+b^2-r^2};$$

$$\therefore \text{chord through centre} = \frac{2pdr}{dp} = \frac{2(a^2+b^2-r^2)}{r} = \frac{2CD^2}{CP},$$

$$\text{diameter} = 2r\frac{dr}{dp} = \frac{2(a^2+b^2-r^2)}{p} = \frac{2CD^2}{PF}.$$

(7) To find the chord through the focus.

Here if $SP = r$, $SY = p$; $p^2 = \frac{b^2r}{2a-r}$;

$$\therefore\ 2\log p = \log b^2 + \log r - \log(2a - r);$$

$$\therefore\ \frac{2dp}{pdr} = \frac{1}{r} + \frac{1}{2a - r} = \frac{2a}{r.(2a - r)};$$

$$\therefore\ \text{chord} = 2p\frac{dr}{dp} = \frac{2r(2a - r)}{a} = \frac{2SP.HP}{AC} = \frac{2CD^2}{AC}.$$

(8) Find the form of the parabola $y = a + bx + cx^2$, which has a contact of the second order, with a given curve at a given point.

Make the given point the origin: then the equation to the parabola becomes $y = bx + cx^2$; and let $y = f(x)$ be the equation to the given curve, from which find p and q.

$$\text{But from above } \frac{dy}{dx} = b + 2cx; \quad \text{and } \frac{d^2y}{dx^2} = 2c;$$

$$\therefore\ p = b + 2cx; \quad \text{and } q = 2c.$$

$$\text{But at the origin } x = 0; \quad \therefore\ b = p, \text{ and } c = \frac{q}{2};$$

$$\therefore\ y = px + \frac{qx^2}{2} = \frac{q}{2}.\left(x^2 + \frac{2p}{q}x + \frac{p^2}{q^2}\right) - \frac{p^2}{2q};$$

$$\therefore\ \left(y + \frac{p^2}{2q}\right) = \frac{q}{2}.\left(x + \frac{p}{q}\right)^2.$$

The equation to a parabola, of which the axis is perpendicular to the axis of x, the co-ordinates of the vertex $-\frac{p^2}{2q}$, and $-\frac{p}{q}$; and the latus rectum $= \frac{2}{q}$.

Cor. The general equation to the second degree, or

$$y^2 + (ax + b)y + cx^2 + ex + f = 0,$$

containing five constants, may have a contact of the fourth order, with a curve. And should there be a point at which $a^2 - 4c = 0$, the osculating curve is a parabola. Immediately before and after this point, a^2 must be greater or less than $4c$; and therefore the osculating parabola is intermediate between an osculating ellipse and hyperbola.

EXAMPLES.

(1) If $y^2 + x^2 = ax - ay$; an equation to the circle, $R = \frac{a}{\sqrt{2}}$.

(2) In the cubical parabola where $a^2y = x^3$;

$$R = -\frac{(a^4 + 9x^4)^{\frac{3}{2}}}{6a^4x};$$

and in the semicubical parabola where $ay^2 = x^3$;

$$R = -\frac{(4a + 9x)^{\frac{3}{2}}}{6a}\sqrt{x}.$$

(3) The equation to the hyperbola being $y^2 = \frac{b^2}{a^2}(x^2 - a^2)$;

$R = \frac{(e^2x^2 - a^2)^{\frac{3}{2}}}{ab}$; and the equation to the evolute is

$$(a\alpha)^{\frac{2}{3}} - (b\beta)^{\frac{2}{3}} = (a^2 + b^2)^{\frac{2}{3}}.$$

(4) In the parabola the chord of curvature through the focus $= 4SP$; and the length of the evolute

$$= \frac{2(SP^{\frac{3}{2}} - SA^{\frac{3}{2}})}{\sqrt{SA}}.$$

(5) If $yx = a^2$, $R = -\frac{(x^4 + a^4)^{\frac{3}{2}}}{2a^2x^3}$, and equation to evolute is

$$(\alpha + \beta)^{\frac{2}{3}} - (\alpha - \beta)^{\frac{2}{3}} = (4a)^{\frac{2}{3}}.$$

(6) The equation to the catenary is $2y = a\left(e^{\frac{x}{a}} + e^{-\frac{x}{a}}\right)$; shew that the radius of curvature is equal, but opposite, to the normal. $R = -\frac{y^2}{a}$.

(7) If $r = a(1 + \cos\theta)$; the radius of curvature $= \frac{2\sqrt{2ar}}{3}$; and chord $= \frac{4r}{3}$.

(8) In the spiral of Archimedes, the radius = the chord of curvature, when $r = \frac{a}{\sqrt{3}}$.

(9) The evolute of the epicycloid, of which the equation is $p^2 = e^2 \cdot \frac{r^2 - a^2}{e^2 - a^2}$, is another epicycloid,

$$p_1^2 = e^2 \cdot \frac{r_1^2 - a_1^2}{e^2 - a^2}; \text{ and } a_1 = \frac{a^2}{e}.$$

(10) Find the chords of curvature drawn through the centre and focus of an hyperbola.

(1°) If $r = CP$; chord $= \frac{2(r^2 - a^2 + b^2)}{r}$.

(2°) If $r = SP$; chord $= \frac{2r(2a + r)}{a}$.

(11) If $y\sqrt{1 + \frac{dx^2}{dy^2}} = c$, be the equation to a curve (the Tractrix); the equation to the evolute is $\frac{d\beta}{da} = \frac{\sqrt{\beta^2 - c^2}}{c}$ (the Catenary).

(12) In the focal distance SP of a parabola, take $SQ = PN$; find the equation to the locus of Q, and the radius of curvature.

$$SQ = r, \quad SA = a, \quad \angle ASQ = \theta; \quad r = 2a \tan \frac{\theta}{2}.$$

$$R = \frac{(r^4 + 24a^2r^2 + 16a^4)^{\frac{3}{2}}}{32a^3(4a^2 + 3r^2)}.$$

(13) Determine that point in a cubical parabola, where the curvature is the greatest: and the point in the common parabola where it is $\frac{1}{8}^{\text{th}}$ of the greatest curvature.

$$(1)\ x = \frac{a}{\sqrt[4]{45}}, \qquad (2)\ x = 3a.$$

(14) If $a\theta = \sqrt{r^2 - a^2} - a \sec^{-1}\left(\frac{r}{a}\right)$, (the involute of the circle), shew that $p = \sqrt{r^2 - a^2}$; and find the equation to the evolute.

(15) In the parabola if D be the point where the axis intersects the directrix, and PN, QM be ordinates of corresponding points in the parabola and evolute, prove that $DM = 3DN$.

(16) If λ be the angle which the normal makes with the axis of x in the ellipse, then

$$R = \frac{a(1-e^2)}{(1-e^2\sin^2\lambda)^{\frac{3}{2}}}.$$

(17) In the curve (hypocycloid) of which the equation is $x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}$; the equation to the evolute is

$$(\alpha+\beta)^{\frac{2}{3}} + (\alpha-\beta)^{\frac{2}{3}} = 2a^{\frac{2}{3}}.$$

(18) Let R and R_1 be the radii of curvature of the extremities of two conjugate diameters of an ellipse, then

$$(R^{\frac{2}{3}} + R_1^{\frac{2}{3}})(ab)^{\frac{2}{3}} = a^2 + b^2.$$

CHAPTER XV.

ENVELOPES TO CURVES. CAUSTICS.

231. When a curve touches a series of curves, all described after a given law, the former is said to be an envelope of the latter; these latter are of a given form, and the problem is to find the touching curve or envelope.

For the better explanation of this application of the Differential Calculus, let us suppose that it was required to find the equation to the curve, touching any number of equal circles, whose centres are in a known curve.

Then if y and x be the co-ordinates of the touching curve, α and β those of the centre of one of the circles

$$(y-\beta)^2+(x-\alpha)^2=r^2.$$

But β and α are the co-ordinates of the known curve;

$$\therefore\ \beta=\phi(\alpha);$$

$$\therefore\ \{y-\phi(\alpha)\}^2+(x-\alpha)^2=r^2.....(1).$$

Now if we suppose α to receive an indefinitely small increment, the equation (1) will belong to an equal circle, the centre of which is indefinitely near to that denoted by equation (1); and the two circles will intersect at a point of which the co-ordinates are ultimately x and y; and similarly proceeding with a third and other circles, we may conceive the touching curve to be formed by the continual intersections of these circles: and to determine its equation, which must be independent of α, α must be eliminated between the equations $\{y-\phi(\alpha)\}^2+(x-\alpha)^2=r^2$, and the equation which indicates that we have passed from the consideration of one circle to the other, that is, the differential of the equation (1), taken with respect to α.

Hence we may conclude, that if $V=f(xy\alpha)=0$ represent the equation of one of the given curves, the touching curve may be found by eliminating α between the equations

$$V=0,\quad \text{and}\quad \frac{dV}{d\alpha}=0.$$

232. That $V=0$, and $\dfrac{dV}{d\alpha}=0$, are simultaneous equa-

tions, may be also thus shewn. Resuming the equation to the circle,

Let $\alpha + \delta\alpha$, and $\beta + \delta\beta$ be the values of α and β in the consecutive circle;

$$\therefore \{x - (\alpha + \delta\alpha)\}^2 + \{y - (\beta + \delta\beta)\}^2 = r^2;$$

therefore by subtraction,

$$(x-\alpha)^2 - \{x - (\alpha + \delta\alpha)\}^2 + (y-\beta)^2 - \{y - (\beta + \delta\beta)\}^2 = 0,$$

$$\text{or } \delta\alpha\{2.(x-\alpha) - \delta\alpha\} + \delta\beta\{2.(y-\beta) - \delta\beta\} = 0,$$

$$\text{or } 2(x-\alpha) + 2.(y-\beta)\frac{\delta\beta}{\delta\alpha} - \{\delta\beta\frac{\delta\beta}{\delta\alpha} + \delta\alpha\} = 0.$$

Now make $\delta\alpha = 0$, and $\delta\beta = 0$, in which case the point of intersection of the two circles becomes a point in the touching curve, and $\frac{\delta\beta}{\delta\alpha}$ becomes the differential coefficient of β with respect to α, or $= \frac{d\beta}{d\alpha}$, and we have $2(x-\alpha) + 2(y-\beta)\frac{d\beta}{d\alpha} = 0$, which is the differential coefficient of $(x-\alpha)^2 + (y-\beta)^2 = r^2$ with respect to α, between which two equations α may be eliminated.

Prob. I. Find the curve which shall touch all the straight lines defined by the equation $y = \alpha x + r\sqrt{\alpha^2 + 1}$, r being a perpendicular of constant length from the origin upon the lines.

Differentiating with respect to α; x and y being constant,

$$x + \frac{r\alpha}{\sqrt{\alpha^2+1}} = 0; \quad \therefore \frac{r}{x} = -\frac{\sqrt{\alpha^2+1}}{\alpha};$$

$$\therefore \frac{1}{\alpha^2} = \frac{r^2}{x^2} - 1 = \frac{r^2 - x^2}{x^2}; \quad \therefore \alpha = \frac{x}{\sqrt{r^2 - x^2}};$$

$$\text{and } \sqrt{\alpha^2+1} = -\alpha\frac{r}{x} = -\frac{r}{\sqrt{r^2-x^2}};$$

$$\therefore y = \alpha x + r\sqrt{\alpha^2+1}$$

$$= \frac{x^2}{\sqrt{r^2-x^2}} - \frac{r^2}{\sqrt{r^2-x^2}} = -\frac{r^2-x^2}{\sqrt{r^2-x^2}} = -\sqrt{r^2-x^2};$$

$$\therefore y^2 + x^2 = r^2, \text{ the equation to a circle.}$$

Prob. II. A given straight line slides between two rectangular axes, find the curve to which it is always a tangent.

Let c be the length of the line, a and b the parts of the axes cut off in any given position of the line;

$$\therefore \frac{x}{a} + \frac{y}{b} = 1, \text{ and } a^2 + b^2 = c^2;$$

$$\therefore \frac{x}{a^2} + \frac{y}{b^2}\frac{db}{da} = 0, \text{ and } a + b\frac{db}{da} = 0; \quad \therefore \frac{db}{da} = -\frac{a}{b};$$

$$\therefore \frac{x}{a^2} - \frac{ya}{b^3} = 0; \quad b = a\sqrt[3]{\frac{y}{x}}; \quad \therefore a^2 + b^2 = a^2\left\{\frac{y^{\frac{2}{3}} + x^{\frac{2}{3}}}{x^{\frac{2}{3}}}\right\} = c^2;$$

$$\therefore a = c\frac{x^{\frac{1}{3}}}{\sqrt{y^{\frac{2}{3}} + x^{\frac{2}{3}}}}; \quad b = c\frac{y^{\frac{1}{3}}}{\sqrt{x^{\frac{2}{3}} + y^{\frac{2}{3}}}};$$

$$\therefore \frac{x}{a} + \frac{y}{b} = \frac{\sqrt{y^{\frac{2}{3}} + x^{\frac{2}{3}}}}{c}\left\{x^{\frac{1}{3}} + y^{\frac{1}{3}}\right\} = 1;$$

$$\therefore (x^{\frac{1}{3}} + y^{\frac{1}{3}})^{\frac{3}{2}} = c, \text{ and } x^{\frac{1}{3}} + y^{\frac{1}{3}} = c^{\frac{2}{3}}.$$

PROB. III. If the equation of condition be $a^n + b^n = c^n$, then the equation to the touching curve will be

$$x^{\frac{n}{n+1}} + y^{\frac{n}{n+1}} = c^{\frac{n}{n+1}}.$$

PROB. IV. Find the curve which touches all the lines defined by $y = mx + \sqrt{m^2a^2 + b^2}$; a and b being constant.

$$a^2y^2 + b^2x^2 = a^2b^2.$$

PROB. V. Find the curve which touches all the ellipses described round the same centre and with coincident axes, the rectangle of the axes being a constant area (m^2).

$$\text{Here } \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \ldots\ldots\ldots(1),$$

$$\text{and } ab = m^2 \ldots\ldots\ldots\ldots(2);$$

$\therefore 2xy = m^2$, the equation to the rectangular hyperbola.

PROB. VI. Find the equation to the curve whose tangent cuts off from the axes two lines the sum of which $= c$.

$$\sqrt{x} + \sqrt{y} = \sqrt{c}.$$

PROB. VII. Find the curve which touches all the curves included under the equation $y = x\tan\theta - \frac{x^2}{4h\cos^2\theta}$, θ being supposed variable.

$$x^2 = 4h(h - y).$$

PROB. VIII. Find the curve when $AD^m = a^{m-1}.AT$.

PROB. IX. Find the curve, when the rectangle contained by two lines, drawn perpendicular to the axis of x, one from the origin, the other from a given point in the axis, to meet the tangent, is $= b^2$.

PROB. X. Find the curve whose tangent cuts off from the axes a constant area; the axes being first rectangular, secondly oblique.

PROB. XI. Find the same as in problem 7, when h and θ both vary; but $m^2 = h^3 \sin^3\theta \cdot \cos\theta$; m^2 being a constant area.

$$xy = \frac{64m^2}{27}.$$

PROB. XII. Two diameters of a circle intersect at right angles, find the locus of the intersections of the chords joining the extremities of the diameters, while the diameters perform a complete revolution. Ans. $x^2 + y^2 = \frac{a^2}{2}$.

CAUSTICS.

233. By the same method as that used in the preceding article, the equations to the curves formed by the intersection of rays reflected by a surface, or refracted through a medium may be found. These curves are called Caustics. Some of them may be practically exhibited by means of a ring of metal, placed on a sheet of paper and held towards the rays of the sun: the curved part of the sugar tongs may be used for the same purpose.

234. PROB. Rays of light fall perpendicularly to the axis of x, find the equation to one of the reflected rays, and the equation to the curve of their intersection, or the Caustic.



QP one of the incident rays,

Pq a reflected ray, making with the normal PG,

$$\angle qPG = \angle GPN.$$

$AN = x$, $\theta = \angle NPG$,

$NP = y$, Y and X the co-ordinates of Pq;

$\therefore\ Y - y = m(X - x)$ is the equation to Pq.

But $m = \tan Pqx = \tan(90 + 2\theta) = -\cot 2\theta$

$$= -\left(\frac{1 - p^2}{2p}\right), \text{ where } p = \frac{dy}{dx};$$

$$\therefore\ Y - y = -\left(\frac{1 - p^2}{2p}\right)(X - x),$$

the equation to the reflected ray.

Now differentiating, Y and X being constant, and $q = \frac{dp}{dx}$;

$$\therefore\ -p = \frac{1 - p^2}{2p} + \tfrac{1}{2}(X - x)\,q\left(\frac{1}{p^2} + 1\right);$$

$$\therefore\ (X - x)\,q\left(\frac{1}{p^2} + 1\right) = -\left(p + \frac{1}{p}\right) = -p\left(1 + \frac{1}{p^2}\right);$$

$$\therefore\ X - x = -\frac{p}{q};\quad X = x - \frac{p}{q} \ldots\ldots\ldots(1),$$

$$Y - y = -\frac{1 - p^2}{2p}(X - x) = \frac{1 - p^2}{2q};$$

$$\therefore\ Y = y + \frac{1 - p^2}{2q} \ldots\ldots\ldots\ldots(2),$$

and from the equation to the curve AP, $y = f(x)$: p and q which are $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$ may be found in terms of y and x: and then between (1), (2), and $y = f(x)$, y and x may be eliminated, and the equation between Y and X which is the required equation to the Caustic, may be found.

Cor. 1. If the incident rays proceed from A, the origin of co-ordinates, we shall find by a similar method that the equation to the reflected ray is

$$Y - y = \frac{2px - y(1 - p^2)}{2py + x(1 - p^2)}(X - x).$$

Cor. 2. To find the length Pp of the reflected ray.

$$Pp^2 = (X-x)^2 + (Y-y)^2$$
$$= \frac{p^2}{q^2} + \frac{(1-p^2)^2}{4q^2} = \frac{(1+p^2)^2}{4q};$$
$$\therefore\ Pp = \tfrac{1}{2}.\frac{1+p^2}{-q},\ \text{or, (Art. 218),}$$

the length of the reflected ray $= \frac{1}{4}^{\text{th}}$ the chord of curvature perpendicular to the axis of x.

Hence, we may construct the caustic: take OP the radius of curvature, draw $Pq = \frac{1}{2}$ the chord $\perp$ to x, and making with OP the same angle that NP does, then p will be a point in the caustic.

Ex. 1. Let the rays fall perpendicularly to the axis of the parabola.

$$y^2 = 4ax;\quad p = \frac{2a}{y};\quad q = -\frac{2a}{y^3}p = -\frac{4a^2}{y^3};$$
$$\therefore\ X = x - \frac{p}{q} = x + \frac{y^2}{2a} = 3x;\quad \therefore\ x = \tfrac{1}{3}X,$$
$$1 - p^2 = 1 - \frac{4a^2}{y^2} = 1 - \frac{a}{x} = \frac{x-a}{x},$$
$$2q = \frac{-8a^3}{y^3} = \frac{-8a^2}{4axy} = \frac{-2a}{xy};$$
$$\therefore\ Y = y + \frac{1-p^2}{2q} = y - y\frac{x-a}{2a} = y\left(\frac{3a-x}{2a}\right);$$
$$\therefore\ Y^2 = \frac{x}{a}.(3a-x)^2 = \frac{X}{27a}(9a-X)^2;$$

whence the caustic cuts the axis at the origin and at a distance $9a$ from the origin.

Ex. 2. Reflecting curve a cycloid; rays parallel to the diameter of the generating circle.

$$AN = x;\quad NP = y;\quad BC = 2a.$$
$$\text{Then } p = \frac{dy}{dx} = \frac{\sqrt{2ay - y^2}}{y},$$
$$q = -\frac{a}{y^2};$$
$$\therefore\ X = x - \frac{p}{q} = x + \frac{y}{a}\sqrt{2ay - y^2}\ \ldots\ldots(1),$$



$$Y = y + \frac{1-p^2}{2q} = \frac{1}{a}(2ay - y^2) \ldots\ldots\ldots (2);$$

$$\therefore\ y = a - \sqrt{a^2 - aY} \ldots\ldots\ldots\ldots\ldots\ldots (3).$$

From (1) $\dfrac{dX}{dy} = \dfrac{2}{a}\sqrt{2ay - y^2} = \dfrac{2}{\sqrt{a}}\sqrt{Y}.$

From (3) $\dfrac{dy}{dY} = \dfrac{a}{2\sqrt{a^2 - aY}} = \dfrac{\sqrt{a}}{2\sqrt{a - Y}};$

$$\therefore\ \frac{dX}{dY} = \frac{\sqrt{Y}}{\sqrt{a - Y}} = \frac{Y}{\sqrt{aY - Y^2}};$$

the equation to a cycloid of which the radius of the generating circle is a, and therefore the base $= AB$.

Ex. 3. Let the rays fall parallel to the diameter of a circle, find the caustic. The Epicycloid.

235. When the pole S of a spiral is the focus of incidence, to find the length Pq of the reflected ray, and the Caustic.



PO the radius of curvature $= R$,

$SP = r$, $\angle SPO = \theta = \angle OPq$,

$SY = p$, $Pq = \rho$;

q being the point of intersection of two consecutive reflected rays, and therefore a point in the caustic.

Now S, O, q may be supposed to be fixed while P moves through an indefinitely small arc.

Also $SO^2 = r^2 + R^2 - 2Rr\cos\theta$.....(1),

$Oq^2 = \rho^2 + R^2 - 2R\rho\cos\theta$.....(2);

$$\therefore r\frac{dr}{d\theta} - R\cos\theta\frac{dr}{d\theta} + Rr\sin\theta = 0,$$

$$\rho\frac{d\rho}{d\theta} - R\cos\theta\frac{d\rho}{d\theta} + R\rho\sin\theta = 0;$$

$$\therefore \frac{dr}{d\theta}(r - R\cos\theta) = -Rr\sin\theta,$$

$$\frac{d\rho}{d\theta}(\rho - R\cos\theta) = -R\rho\sin\theta.$$

But $r + \rho$, for a very small variation of P is constant;

$$\therefore \frac{dr}{d\theta} = -\frac{d\rho}{d\theta};$$ hence by division, we have

$$\frac{\rho}{r} = \frac{R\cos\theta - \rho}{r - R\cos\theta};$$

$$\therefore \rho = \frac{Rr\cos\theta}{2r - R\cos\theta}.$$

But $R\cos\theta = \frac{1}{2}$ chord through $S = p\frac{dr}{dp}$;

$$\therefore \rho = \frac{pr\frac{dr}{dp}}{2r - p\frac{dr}{dp}};$$

$$\therefore \frac{1}{\rho} = \frac{2}{p}\cdot\frac{dp}{dr} - \frac{1}{r} = \frac{d}{dr}\left(\log\frac{p^2}{r}\right),$$

whence ρ the length of the reflected ray may be found.

236. To find the equation to the locus of q.

Join Sq; draw $Sy \perp$ to Pq.

Let $Sq = r_1$; $Sy = p_1$;

$$\therefore\ p_1 = r \sin 2\theta = 2r\sqrt{1 - \frac{p^2}{r^2}} \cdot \frac{p}{r} = \frac{2p\sqrt{r^2 - p^2}}{r} \quad \ldots(1),$$

$$\begin{aligned} r_1^2 &= r^2 + \rho^2 - 2r\rho \cos 2\theta \\ &= (r + \rho)^2 - 4r\rho \cos^2\theta \\ &= (r + \rho)^2 - \frac{4\rho p^2}{r} \quad \ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots(2); \end{aligned}$$

whence from the given equation $p = f(r)$; and from $\rho = \phi(r)$, p and ρ may be found in terms of r; and from (1) and (2) r may be eliminated, and the equation found between p_1 and r_1, which is the equation to the caustic.

Ex. 4. An indefinitely small reflector is placed in a circular ring. Every other point of the ring is luminous, find the caustic.

Here $p = \dfrac{r^2}{2a}$; $2a =$ diameter of ring;

$$\therefore \frac{p^2}{r} = \frac{r^3}{4a^2}; \quad \therefore \log\left(\frac{p^2}{r}\right) = 3\log r - \log(4a^2);$$

$$\therefore \frac{1}{\rho} = \frac{d}{dr} \cdot \log\left(\frac{p^2}{r}\right) = \frac{3}{r}; \quad \therefore \rho = \frac{r}{3}.$$

Hence the caustic will be a circle, the diameter $\dfrac{2a}{3}$.

Ex. 5. Let the reflecting curve be the equiangular spiral.

$$p = mr; \quad \frac{p^2}{r} = m^2 r; \quad \log\left(\frac{p^2}{r}\right) = \log r + \log m^2;$$

$$\therefore \frac{1}{\rho} = \frac{1}{r}; \quad \rho = r,$$

$$p_1 = 2m\sqrt{r^2 - m^2 r^2} = 2mr\sqrt{1 - m^2},$$

$$r_1^2 = 4r^2 - 4p^2 = 4r^2(1 - m^2);$$

$$\therefore p_1 = mr_1;$$

or the spiral is a similar equiangular spiral.

Ex. 6. Let the radiating point be the extremity of a diameter of a circle, find the caustic. The Cardioid.

237. When rays of light fall upon a plane refracting surface, find the equation to the caustic.



QR an incident, RS a refracted ray;

$QA \perp$ to BAC the surface;

Q the origin of co-ordinates;

$AQ = a$; $\angle RQA = \theta$; $\angle RqA = \phi$;

$\therefore y = -x \tan\phi + C$ is equation to Rq.

But $x = 0$; $y = AR = a \tan\theta = C$;

$\therefore y = -x \tan\phi + a \tan\theta$.....(1),

also $\sin\theta = m \sin\phi$.....(2),

since the sines of the angles of incidence and refraction are in a constant ratio.

If now θ and ϕ be supposed to vary slightly, while y and x remain constant, the intersection p, of two of the refracted rays will be found, and p will be a point in the caustic.

From (1) $\dfrac{x d\phi}{\cos^2\phi} = \dfrac{a d\theta}{\cos^2\theta}$,

(2) $\cos\theta\, d\theta = m \cos\phi\, d\phi$;

$$\therefore \sqrt[3]{\frac{x}{ma}} = \frac{\cos\phi}{\cos\theta} = \frac{1}{m} \cdot \frac{\tan\theta}{\tan\phi};$$

$$\therefore 1 + \tan^2\phi = \frac{1}{\cos^2\phi} = 1 + \frac{1}{m^2} \tan^2\theta \sqrt[3]{\frac{m^2a^2}{x^2}}.$$

But $\dfrac{1}{\cos^2\phi} = (1 + \tan^2\theta) \sqrt[3]{\dfrac{m^2a^2}{x^2}}$;

$$\therefore \tan^2\theta \sqrt[3]{\frac{m^2a^2}{x^2}} \cdot \left\{1 - \frac{1}{m^2}\right\} = 1 - \sqrt[3]{\frac{m^2a^2}{x^2}};$$

$$\therefore \tan\theta = \frac{m}{\sqrt{m^2-1}} \sqrt{\frac{x^{\frac{2}{3}}}{(ma)^{\frac{2}{3}}} - 1};$$

$$\therefore \tan\phi = \frac{1}{\sqrt{m^2-1}} \sqrt[3]{\frac{ma}{x}} \sqrt{\frac{x^{\frac{2}{3}}}{(ma)^{\frac{2}{3}}} - 1};$$

$$\therefore y = \left\{\frac{ma}{\sqrt{m^2-1}} - \frac{x^{\frac{2}{3}}\sqrt[3]{ma}}{\sqrt{m^2-1}}\right\} \sqrt{\frac{x^{\frac{2}{3}}}{(ma)^{\frac{2}{3}}} - 1}$$

$$= -\frac{ma}{\sqrt{m^2-1}} \left\{\left(\frac{x}{ma}\right)^{\frac{2}{3}} - 1\right\}^{\frac{3}{2}}.$$

Make $\frac{ma}{\sqrt{m^2-1}} = \beta$, and $ma = \alpha$;

$$\therefore \left(\frac{y}{\beta}\right)^{\frac{2}{3}} - \left(\frac{x}{\alpha}\right)^{\frac{2}{3}} = -1,$$

the evolute of the hyperbola, of which the centre is A and focus Q.

If m be < 1, the caustic will be the evolute of an ellipse.

CHAPTER XVI.

CHANGE OF THE INDEPENDENT VARIABLE. LAGRANGE'S THEOREM.

---

238. In the preceding pages, we have in general assumed, that x is the independent variable, and have derived the differential coefficients, from the equation $y = f(x)$.

We shall now proceed to find what values must be put for $\frac{dy}{dx}$, $\frac{d^2y}{dx^2}$, $\frac{d^3y}{dx^3}$, &c. when y becomes the independent variable; and afterwards, find what must be substituted for the same quantities, when both y and x are functions of a new variable θ.

We have already proved that $\frac{dy}{dx} = \frac{1}{\frac{dx}{dy}}$, and that

$$\frac{dy}{dx} = \frac{dy}{d\theta} \cdot \frac{d\theta}{dx};$$

in the following propositions the same results will be obtained, but in a different manner.

239. Prop. If $y = f(x)$; and $\therefore x = f^{-1}(y)$; find

$\frac{dy}{dx}$, $\frac{d^2y}{dx^2}$, &c. in terms of $\frac{dx}{dy}$, $\frac{d^2x}{dy^2}$, &c.

Let y become $y + k$, when x becomes $x + h$;

$$\therefore k = \frac{dy}{dx}h + \frac{d^2y}{dx^2}\frac{h^2}{1.2} + \frac{d^3y}{dx^3}\frac{h^3}{2.3} + \&c. \ldots . (1).$$

But since $x + h = f^{-1}(y + k)$;

$$\therefore h = \frac{dx}{dy}k + \frac{d^2x}{dy^2}\frac{k^2}{1.2} + \frac{d^3x}{dy^3} \cdot \frac{k^3}{2.3} + \&c.;$$

therefore substituting for h in equation (1), we have

$$k = \frac{dy}{dx}\left\{\frac{dx}{dy}k + \frac{d^2x}{dy^2}\frac{k^2}{1.2} + \frac{d^3x}{dy^3}\frac{k^3}{2.3} + \&c.\right\}$$

$$+ \frac{d^2y}{dx^2}\left\{\frac{dx^2}{dy^2}k^2 + \frac{dx}{dy} \cdot \frac{d^2x}{dy^2}k^3 + \&c.\right\} \cdot \frac{1}{1.2}$$

$$+\frac{d^3y}{dx^3}\left\{\frac{dx^3}{dy^3}k^3+\&c.\right\}\frac{1}{1.2.3}$$
$$+\ \&c.$$
$$=\frac{dy}{dx}\cdot\frac{dx}{dy}k+\left(\frac{dy}{dx}\cdot\frac{d^2x}{dy^2}+\frac{d^2y}{dx^2}\cdot\frac{dx^2}{dy^2}\right)\frac{k^2}{1.2}$$
$$+\left\{\frac{d^3x}{dy^3}\frac{dy}{dx}+3\frac{dx}{dy}\cdot\frac{d^2y}{dx^2}\cdot\frac{d^2x}{dy^2}+\frac{d^3y}{dx^3}\cdot\frac{dx^3}{dy^3}\right\}\frac{k^3}{2.3}$$
$$+\ \&c.$$

equating coefficients,

$$1=\frac{dy}{dx}\cdot\frac{dx}{dy};\quad\therefore\ \frac{dy}{dx}=\frac{1}{\frac{dx}{dy}},$$

$$\frac{dy}{dx}\cdot\frac{d^2x}{dy^2}+\frac{d^2y}{dx^2}\cdot\frac{dx^2}{dy^2}=0;\quad\therefore\ \frac{d^2x}{dy^2}+\frac{d^2y}{dx^2}\cdot\frac{dx^3}{dy^3}=0;$$

$$\therefore\ \frac{d^2y}{dx^2}=\frac{-\frac{d^2x}{dy^2}}{\frac{dx^3}{dy^3}},$$

or putting

p, q, r, &c. for the differential coefficients when $y=f(x)$,

p_1, q_1, r_1, &c..... when $x=f^{-1}(y)$;

$$p=\frac{1}{p_1};\quad q=\frac{-q_1}{p_1^3}.$$

Also, since $\frac{d^3x}{dy^3}\cdot\frac{dy}{dx}+3\frac{dx}{dy}\cdot\frac{d^2y}{dx^2}\cdot\frac{d^2x}{dy^2}+\frac{d^3y}{dx^3}\cdot\frac{dx^3}{dy^3}=0;$

$$\therefore\ r_1p+3p_1qq_1+rp_1^3=0,\ \text{ or }\ \frac{r_1}{p_1}-\frac{3q_1^2}{p_1^2}+rp_1^3=0;$$

$$\therefore\ r=\frac{3q_1^2-r_1p_1}{p_1^5};$$

and similarly may other coefficients be found.

240. Take the expression for the radius of curvature.

$$R=\frac{\left(1+\frac{dy^2}{dx^2}\right)^{\frac{3}{2}}}{-\frac{d^2y}{dx^2}}=\frac{(1+p^2)^{\frac{3}{2}}}{-q},\ \text{ and if }\ x=f^{-1}(y);$$

$$\therefore\ R = \frac{\left(1 + \frac{1}{p_1^2}\right)^{\frac{3}{2}}}{\frac{q_1}{p_1^3}} = \frac{(p_1^2 + 1)^{\frac{3}{2}}}{q_1}.$$

Ex. Let $y^2 = 4mx$; $\therefore\ x = \frac{y^2}{4m^2}$;

$$\therefore\ \frac{dx}{dy} = p_1 = \frac{y}{2m};\quad \frac{d^2x}{dy^2} = q_1 = \frac{1}{2m};$$

$$\therefore\ R = \frac{\left(1 + \frac{y^2}{4m^2}\right)^{\frac{3}{2}}}{\frac{1}{2m}} = \frac{(4m^2 + y^2)^{\frac{3}{2}}}{4m^2} = \frac{2 \cdot (m + x)^{\frac{3}{2}}}{\sqrt{m}}.$$

241. Again, if $y = f(\theta)$, and $x = \phi(\theta)$, to express $\frac{dy}{dx}$, $\frac{d^2y}{dx^2}$, &c. in terms of $\frac{dy}{d\theta}$, $\frac{dx}{d\theta}$, $\frac{d^2y}{d\theta^2}$, $\frac{d^2x}{d\theta^2}$, &c.

Let $y + k$, $\theta + m$, and $x + h$, be corresponding values of y, θ, and x; therefore, by Taylor's Theorem,

$$k = \frac{dy}{d\theta}m + \frac{d^2y}{d\theta^2}\cdot\frac{m^2}{1.2} + \frac{d^3y}{d\theta^3}\cdot\frac{m^3}{2.3} + \&c.$$

$$h = \frac{dx}{d\theta}m + \frac{d^2x}{d\theta^2}\cdot\frac{m^2}{1.2} + \frac{d^3x}{d\theta^3}\cdot\frac{m^3}{2.3} + \&c.$$

also, since $y + k$ is a function of $x + h$,

$$k = \frac{dy}{dx}h + \frac{d^2y}{dx^2}\cdot\frac{h^2}{1.2} + \frac{d^3y}{dx^3}\cdot\frac{h^3}{2.3} + \&c.$$

therefore, substituting for k and h,

$$\frac{dy}{d\theta}m + \frac{d^2y}{d\theta^2}\cdot\frac{m^2}{1.2} + \&c.$$

$$= \frac{dy}{dx}\cdot\left(\frac{dx}{d\theta}m + \frac{d^2x}{d\theta^2}\cdot\frac{m^2}{1.2} + \&c.\right)$$

$$+ \frac{1}{1.2}\cdot\frac{d^2y}{dx^2}\left(\frac{dx^2}{d\theta^2}m^2 + \frac{dx}{d\theta}\cdot\frac{d^2x}{d\theta^2}m^3 + \&c.\right) + \&c.$$

$$= \frac{dy}{dx}\cdot\frac{dx}{d\theta}m + \left(\frac{dy}{dx}\cdot\frac{d^2x}{d\theta^2} + \frac{d^2y}{dx^2}\cdot\frac{dx^2}{d\theta^2}\right)\cdot\frac{m^2}{1.2} + \&c.;$$

$$\therefore \frac{dy}{d\theta} = \frac{dy}{dx} \cdot \frac{dx}{d\theta}, \quad \text{or} \quad \frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}};$$

$$\text{and} \quad \frac{d^2y}{d\theta^2} = \frac{dy}{dx} \cdot \frac{d^2x}{d\theta^2} + \frac{d^2y}{dx^2} \cdot \frac{dx^2}{d\theta^2};$$

$$\therefore \frac{d^2y}{dx^2} = \frac{\frac{d^2y}{d\theta^2} - \frac{dy}{dx} \cdot \frac{d^2x}{d\theta^2}}{\frac{dx^2}{d\theta^2}}$$

$$= \frac{\frac{dx}{d\theta} \cdot \frac{d^2y}{d\theta^2} - \frac{dy}{d\theta} \cdot \frac{d^2x}{d\theta^2}}{\left(\frac{dx}{d\theta}\right)^3},$$

and similarly may $\frac{d^3y}{dx^3}$ be found.

242. The expression for the radius of curvature being

$$R = \frac{\left(1 + \frac{dy^2}{dx^2}\right)^{\frac{3}{2}}}{-\frac{d^2y}{dx^2}},$$

we have, when $x = \phi(\theta)$ and $y = f(\theta)$,

$$\left(1 + \frac{dy^2}{dx^2}\right)^{\frac{3}{2}} = \frac{\left\{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)\right\}^{\frac{3}{2}}}{\left(\frac{dx}{d\theta}\right)^3},$$

$$\text{and} \quad -\frac{d^2y}{dx^2} = \frac{\frac{dy}{d\theta} \cdot \frac{d^2x}{d\theta^2} - \frac{dx}{d\theta} \cdot \frac{d^2y}{d\theta^2}}{\left(\frac{dx}{d\theta}\right)^3};$$

$$\therefore R = \frac{\left\{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2\right\}^{\frac{3}{2}}}{\frac{dy}{d\theta} \cdot \frac{d^2x}{d\theta^2} - \frac{dx}{d\theta} \cdot \frac{d^2y}{d\theta^2}}.$$

243. Let $x = -r\cos\theta$ and $y = r\sin\theta$, find R, r being a function of θ.

$$\frac{dx}{d\theta} = r\sin\theta - \cos\theta\frac{dr}{d\theta};\quad \frac{dy}{d\theta} = r\cos\theta + \sin\theta\,.\frac{dr}{d\theta};$$

$$\frac{d^2x}{d\theta^2} = r\cos\theta + 2\sin\theta\frac{dr}{d\theta} - \cos\theta\,.\frac{d^2r}{d\theta^2},$$

$$\frac{d^2y}{d\theta^2} = -r\sin\theta + 2\cos\theta\frac{dr}{d\theta} + \sin\theta\,.\frac{d^2r}{d\theta^2};$$

$$\therefore \frac{dx^2}{d\theta^2} + \frac{dy^2}{d\theta^2} = r^2 + \frac{dr^2}{d\theta^2};$$

$$\frac{dy}{d\theta}\frac{d^2x}{d\theta^2} - \frac{dx}{d\theta}\cdot\frac{d^2y}{d\theta^2} = r^2 + 2\,.\frac{dr^2}{d\theta^2} - r\,.\frac{d^2r}{d\theta^2};$$

$$\therefore R = \frac{\left(r^2 + \frac{dr^2}{d\theta^2}\right)^{\frac{3}{2}}}{r^2 + 2\,.\frac{dr^2}{d\theta^2} - r\frac{d^2r}{d\theta^2}}.$$

Ex. 1. Let $r = a\sin\theta$, the equation to the circle from a point in the circumference, θ being the angle between the tangent at origin and r, and a the diameter;

$$\therefore \frac{dr}{d\theta} = a\cos\theta = \sqrt{a^2 - r^2},\quad \frac{d^2r}{d\theta^2} = -a\sin\theta = -r;$$

$$\therefore R = \frac{(r^2 + a^2 - r^2)^{\frac{3}{2}}}{r^2 + 2a^2 - 2r^2 + r^2} = \frac{a^3}{2a^2} = \frac{a}{2}.$$

Ex. 2. Let $r^2 = a^2\cos 2\theta$, the equation to the Lemniscata;

$$\therefore \frac{dr}{d\theta} = -\frac{a^2}{r}\,.\sin 2\theta = -\frac{a^2}{r}\,.\sqrt{1 - \frac{r^4}{a^4}};$$

$$\therefore r^2 + \frac{dr^2}{d\theta^2} = r^2 + \frac{a^4 - r^4}{r^2} = \frac{a^4}{r^2},$$

$$\frac{d^2r}{d\theta^2} = \frac{a^2}{r^2}\,.\sin 2\theta\,.\frac{dr}{d\theta} - \frac{2a^2}{r}\,.\cos 2\theta$$

$$= -\frac{a^4}{r^3}\left(1 - \frac{r^4}{a^4}\right) - \frac{2a^2}{r}\cdot\frac{r^2}{a^2};$$

$$\therefore -r\frac{d^2r}{d\theta^2} = \frac{a^4 - r^4}{r^2} + 2r^2 = \frac{a^4 + r^4}{r^2};$$

$$\therefore\ r^2 + 2\frac{dr^2}{d\theta^2} - r\frac{d^2r}{d\theta^2} = r^2 + 2\cdot\frac{a^4 - r^4}{r^2} + \frac{a^4 + r^4}{r^2} = \frac{3a^4}{r^2};$$

$$\therefore\ R = \frac{\left(\frac{a^4}{r^2}\right)^{\frac{3}{2}}}{\frac{3a^4}{r^3}} = \frac{a^6}{3a^4r} = \frac{a^2}{3r}.$$

Ex. 3. Let the equation

$$\frac{d^2y}{dx^2} - \frac{x}{1-x^2}\cdot\frac{dy}{dx} + \frac{y}{1-x^2} = 0,$$

be transformed into one where $\theta = \cos^{-1}x$ shall be the independent variable.

$$x = \cos\theta;\quad \therefore\ \frac{dx}{d\theta} = -\sin\theta;\quad \frac{d^2x}{d\theta^2} = -\cos\theta = -x,$$

$$\frac{dy}{dx} = \frac{dy}{d\theta}\cdot\frac{d\theta}{dx} = -\frac{1}{\sin\theta}\cdot\frac{dy}{d\theta},$$

$$\text{and}\ \frac{d^2y}{dx^2} = \frac{\frac{d^2y}{d\theta^2} - \frac{dy}{dx}\cdot\frac{d^2x}{d\theta^2}}{\frac{dx^2}{d\theta^2}} = \frac{\frac{d^2y}{d\theta^2} - \frac{1}{\sin\theta}\cdot\cos\theta\frac{dy}{d\theta}}{\sin^2\theta};$$

$$\therefore\ \frac{1}{\sin^2\theta}\cdot\frac{d^2y}{d\theta^2} - \frac{\cos\theta}{\sin^3\theta}\cdot\frac{dy}{d\theta} + \frac{\cos\theta}{\sin^3\theta}\cdot\frac{dy}{d\theta} + \frac{y}{\sin^2\theta} = 0;$$

$$\therefore\ \frac{d^2y}{d\theta^2} + y = 0,$$

which is satisfied by making $y = A\cos(\theta + B)$.

244. Find the radius of curvature, when the arc is the independent variable.

$$R = \frac{\left(1 + \frac{dy^2}{dx^2}\right)^{\frac{3}{2}}}{-\frac{d^2y}{dx^2}},\ \text{and}\ \frac{ds^2}{dx^2} = 1 + \frac{dy^2}{dx^2}.$$

But if x and y be functions of s,

$$\frac{ds^2}{dx^2} = 1 + \frac{\left(\frac{dy}{ds}\right)^2}{\left(\frac{dx}{ds}\right)^2} = \left(\frac{ds}{dx}\right)^2\left\{\left(\frac{dx}{ds}\right)^2 + \left(\frac{dy}{ds}\right)^2\right\};$$

$$\therefore \quad \left(\frac{dx}{ds}\right)^2 + \left(\frac{dy}{ds}\right)^2 = 1,$$

$$\text{and} \quad \left(1 + \frac{dy^2}{dx^2}\right)^{\frac{1}{2}} = \left(\frac{ds}{dx}\right)^3,$$

$$\text{and} \quad -\frac{d^2y}{dx^2} = \frac{\frac{dy}{ds}\cdot\frac{d^2x}{ds^2} - \frac{dx}{ds}\cdot\frac{d^2y}{ds^2}}{\left(\frac{dx}{ds}\right)^3},$$

$$\therefore \quad R = \frac{1}{\frac{dy}{ds}\cdot\frac{d^2x}{ds^2} - \frac{dx}{ds}\cdot\frac{d^2y}{ds^2}},$$

whence by multiplying the numerator and denominator by ds^3,

$$R = \frac{ds^3}{dy\,d^2x - dx\,d^2y},$$

where dy, dx, d^2y, and d^2x are the first and second differentials of y and x with respect to s.

245. Again, $\because \quad \frac{1}{R} = \frac{dy}{ds}\cdot\frac{d^2x}{ds^2} - \frac{dx}{ds}\cdot\frac{d^2y}{ds^2}$;

$$\therefore \quad \frac{1}{R^2} = \frac{dy^2}{ds^2}\cdot\left(\frac{d^2x}{ds^2}\right)^2 - 2\frac{dy}{ds}\cdot\frac{dx}{ds}\cdot\frac{d^2x}{ds^2}\cdot\frac{d^2y}{ds^2} + \frac{dx^2}{ds^2}\cdot\left(\frac{d^2y}{ds^2}\right)^2$$

$$= \left(\frac{d^2x}{ds^2}\right)^2 + \left(\frac{d^2y}{ds^2}\right)^2 - \left\{\frac{dx}{ds}\cdot\frac{d^2x}{ds^2} + \frac{dy}{ds}\cdot\frac{d^2y}{ds^2}\right\}^2.$$

But $\because \quad \frac{dx^2}{ds^2} + \frac{dy^2}{ds^2} = 1$; $\quad \therefore \quad \frac{dx}{ds}\cdot\frac{d^2x}{ds^2} + \frac{dy}{ds}\cdot\frac{d^2y}{ds^2} = 0$;

$$\therefore \quad \frac{1}{R} = \sqrt{\left(\frac{d^2x}{ds^2}\right)^2 + \left(\frac{d^2y}{ds^2}\right)^2};$$

$$\therefore \quad R = \frac{1}{\sqrt{\left(\frac{d^2x}{ds^2}\right)^2 + \left(\frac{d^2y}{ds^2}\right)^2}}.$$

Cor. If d^2x, and d^2y be put for the second differentials of x and y, and we multiply the numerator and denominator by ds^2,

$$R = \frac{ds^2}{\sqrt{(d^2x)^2 + (d^2y)^2}}.$$

Ex. Find the radius of curvature to the catenary.

Here $x = \sqrt{c^2 + s^2}$; $y = c \log\left(\frac{s + \sqrt{s^2 + c^2}}{c}\right)$,

$$\frac{dx}{ds} = \frac{s}{\sqrt{c^2 + s^2}}; \quad \frac{d^2x}{ds^2} = \frac{c^2}{(c^2 + s^2)^{\frac{3}{2}}},$$

$$\frac{dy}{ds} = \frac{c}{\sqrt{c^2 + s^2}}; \quad \frac{d^2y}{ds^2} = \frac{-cs}{(c^2 + s^2)^{\frac{3}{2}}};$$

$$\therefore \left(\frac{d^2y}{ds^2}\right)^2 + \left(\frac{d^2x}{ds^2}\right)^2 = \frac{c^4 + c^2s^2}{(c^2 + s^2)^3} = \frac{c^2}{(c^2 + s^2)^2};$$

$$\therefore R = \frac{c^2 + s^2}{c} = \frac{x^2}{c}.$$

246. Next, let $u = f(x, y)$, to find $\frac{du}{dx}$, and $\frac{du}{dy}$ in terms of r, and θ, when x and y are functions of r and θ, so that $x = \phi(r, \theta)$; $y = \psi(r, \theta)$.

$$\frac{du}{dr} = \frac{du}{dx}\cdot\frac{dx}{dr} + \frac{du}{dy}\cdot\frac{dy}{dr},$$

$$\frac{du}{d\theta} = \frac{du}{dx}\cdot\frac{dx}{d\theta} + \frac{du}{dy}\cdot\frac{dy}{d\theta};$$

$$\therefore \frac{du}{dr}\cdot\frac{dy}{d\theta} - \frac{du}{d\theta}\cdot\frac{dy}{dr} = \frac{du}{dx}\cdot\left(\frac{dx}{dr}\cdot\frac{dy}{d\theta} - \frac{dy}{dr}\cdot\frac{dx}{d\theta}\right),$$

$$\frac{du}{dr}\cdot\frac{dx}{d\theta} - \frac{du}{d\theta}\cdot\frac{dx}{dr} = -\frac{du}{dy}\cdot\left(\frac{dx}{dr}\cdot\frac{dy}{d\theta} - \frac{dy}{dr}\cdot\frac{dx}{d\theta}\right);$$

$$\therefore \frac{du}{dx} = \frac{\frac{du}{dr}\cdot\frac{dy}{d\theta} - \frac{du}{d\theta}\cdot\frac{dy}{dr}}{\frac{dx}{dr}\cdot\frac{dy}{d\theta} - \frac{dy}{dr}\cdot\frac{dx}{d\theta}},$$

$$\frac{du}{dy} = -\frac{\frac{du}{dr}\cdot\frac{dx}{d\theta} - \frac{du}{d\theta}\cdot\frac{dx}{dr}}{\frac{dx}{dr}\cdot\frac{dy}{d\theta} - \frac{dy}{dr}\cdot\frac{dx}{d\theta}}.$$

247. These values are much simplified, when

$$x = r\cos\theta\,; \quad \text{and } y = r\sin\theta.$$

$$\text{For } \frac{dx}{dr} = \cos\theta\,; \quad \frac{dy}{dr} = \sin\theta,$$

$$\frac{dx}{d\theta} = -r\sin\theta\,; \quad \frac{dy}{d\theta} = r\cos\theta\,;$$

$$\therefore \frac{dx}{dr}\cdot\frac{dy}{d\theta} - \frac{dy}{dr}\cdot\frac{dx}{d\theta} = r(\cos^2\theta + \sin^2\theta) = r\,;$$

$$\therefore \frac{du}{dx} = \frac{1}{r}\left(\frac{du}{dr}\cdot r\cos\theta - \frac{du}{d\theta}\cdot\sin\theta\right)$$

$$= \frac{du}{dr}\cdot\cos\theta - \frac{du}{d\theta}\cdot\frac{\sin\theta}{r} \quad \ldots\ldots\ldots\ldots(1),$$

$$\frac{du}{dy} = \frac{du}{dr}\cdot\sin\theta + \frac{du}{d\theta}\cdot\frac{\cos\theta}{r} \quad \ldots\ldots\ldots\ldots(2).$$

Ex. 1. Transform $x\dfrac{du}{dy} - y\dfrac{du}{dx}$, to variables θ and r: when $x = r\cos\theta$; $y = r\sin\theta$.

$$(2) \times x = r\frac{du}{dr}\cdot\cos\theta\cdot\sin\theta + \frac{du}{d\theta}\cdot\cos^2\theta\,;$$

$$(1) \times y = r\frac{du}{dr}\cdot\cos\theta\cdot\sin\theta - \frac{du}{d\theta}\cdot\sin^2\theta\,;$$

$$\therefore x\frac{du}{dy} - y\frac{du}{dx} = \frac{du}{d\theta}\cdot(\sin^2\theta + \cos^2\theta) = \frac{du}{d\theta}.$$

Ex. 2. If $\dfrac{d^2u}{dx^2} + \dfrac{d^2u}{dy^2} = 0$; transform it when $x = r\cos\theta$, and $y = r\sin\theta$.

$$\text{From (2), } \frac{du}{dy} = \frac{du}{dr}\cdot\sin\theta + \frac{du}{d\theta}\cdot\frac{\cos\theta}{r} = V, \text{ suppose;}$$

$$\therefore \frac{d^2u}{dy^2} = \frac{dV}{dy} = \frac{dV}{dr}\cdot\sin\theta + \frac{dV}{d\theta}\cdot\frac{\cos\theta}{r}.$$

$$\text{But } \frac{dV}{dr} = \frac{d^2u}{dr^2}\cdot\sin\theta + \frac{d^2u}{d\theta\,dr}\cdot\frac{\cos\theta}{r} - \frac{du}{d\theta}\cdot\frac{\cos\theta}{r^2},$$

$$\frac{dV}{d\theta} = \frac{d^2u}{dr\,d\theta}\cdot\sin\theta + \frac{du}{dr}\cdot\cos\theta + \frac{d^2u}{d\theta^2}\cdot\frac{\cos\theta}{r} - \frac{du}{d\theta}\cdot\frac{\sin\theta}{r^2};$$

$$\therefore \frac{d^2u}{dy^2} = \sin^2\theta\cdot\frac{d^2u}{dr^2} + \frac{\cos^2\theta}{r^2}\cdot\frac{d^2u}{d\theta^2} + \frac{\cos^2\theta}{r}\cdot\frac{du}{dr}$$

$$+ \frac{2\sin\theta\cdot\cos\theta}{r}\cdot\left\{\frac{d^2u}{d\theta\,dr} - \frac{1}{r}\cdot\frac{du}{d\theta}\right\}.$$

Similarly, or by changing θ into $\frac{\pi}{2} - \theta$.

$$\frac{d^2u}{dx^2} = \cos^2\theta\cdot\frac{d^2u}{dr^2} + \frac{\sin^2\theta}{r^2}\cdot\frac{d^2u}{d\theta^2} + \frac{\sin^2\theta}{r}\cdot\frac{du}{dr}$$

$$- \frac{2\sin\theta\cdot\cos\theta}{r}\cdot\left\{\frac{d^2u}{d\theta\,dr} - \frac{1}{r}\cdot\frac{du}{d\theta}\right\};$$

$$\therefore \frac{d^2u}{dx^2} + \frac{d^2u}{dy^2} = \frac{d^2u}{dr^2} + \frac{1}{r^2}\cdot\frac{d^2u}{d\theta^2} + \frac{1}{r}\cdot\frac{du}{dr} = 0.$$

248. Transform the double integral $\iint V\,dx\,dy$ into one where r and θ are the variables, x and y being the same as before.

$$\because\ x = r\cos\theta;\quad y = r\sin\theta;$$

$$\therefore\ dx = \cos\theta\,.\,dr - r\sin\theta\,.\,d\theta,$$

$$dy = \sin\theta\,.\,dr + r\cos\theta\,.\,d\theta.$$

Now since in integrating, one of the quantities y or x is supposed to vary, while the other is constant, let $dx = 0$;

$$\therefore\ 0 = \cos\theta\,.\,dr - r\sin\theta\,.\,d\theta,$$

$$dy = \sin\theta\,.\,dr + r\cos\theta\,.\,d\theta;$$

$$\therefore\ \sin\theta\,.\,dy = dr;\ \text{eliminating } d\theta;$$

$$\therefore\ \text{if } dy = 0;\quad dr = 0;\quad \therefore\ dx = -r\sin\theta\,.\,d\theta;$$

$$\therefore\ dx\,dy = -r\sin\theta\,.\,d\theta\times\frac{dr}{\sin\theta} = -r\,dr\,d\theta;$$

$$\therefore\ \iint V\,dx\,dy = -\iint V_1 r\,dr\,d\theta.$$

Thus if $V = e^{x^2+y^2}$; $\iint e^{x^2+y^2}\,dx\,dy = -\iint e^{r^2} r\,dr\,d\theta.$

Ex. 3. If $\frac{d^2u}{dx^2} + \frac{d^2u}{dy^2} = 0$, and $x^2 + y^2 = r^2$, transform to an equation in which r is the independent variable.

$$\text{Ans. } \frac{d^2u}{dr^2} + \frac{1}{r} \cdot \frac{du}{dr} = 0.$$

Ex. 4. If $\frac{d^2u}{dx^2} + \frac{d^2u}{dy^2} + \frac{d^2u}{dz^2} = 0$; $r^2 = x^2 + y^2 + z^2$,

$$\text{then } \frac{d^2u}{dr^2} + \frac{2}{r} \cdot \frac{du}{dr} = 0.$$

Ex. 5. If $\frac{d^2u}{dx^2} + \frac{d^2u}{dy^2} + \frac{d^2u}{dz^2} = 0$; and

$$x = r\cos\theta; \quad y = r\sin\theta \cdot \sin\phi; \quad z = r\sin\theta \cdot \cos\phi;$$

transform into a function of r, θ, ϕ;

assume $\rho = r\sin\theta$; and use Ex. 2;

$$\therefore r\frac{d^2(ru)}{dr^2} + \frac{1}{\sin^2\theta} \cdot \frac{d^2u}{d\phi^2} + \frac{1}{\sin\theta} \cdot \frac{d}{d\theta} \cdot \left(\sin\theta \cdot \frac{du}{d\theta}\right) = 0.$$

See *Camb. Math. Journal*, Vol. I, p. 121; and O'Brien's *Tracts*.

Ex. 6. Transform $\iiint V dx dy dz$, to a function of r, θ, ϕ,

$$\iiint V dx dy dz = \iiint V r^2 dr \sin\theta \cdot d\theta \cdot d\phi.$$

LAGRANGE'S THEOREM.

249. Let $u = f(y)$, where $y = z + x\phi(y)$, and z is independent of x; required u or $f(y)$ in terms of x.

By Maclaurin,

$$u = U_0 + U_1 x + U_2 \frac{x^2}{1.2} + U_3 \frac{x^3}{2.3} + \&c. + \frac{U_n x^n}{1.2.3 \ldots n} + \&c.$$

where U_0, U_1, U_2, &c. are the values of u, $\frac{du}{dx}$, $\frac{d^2u}{dx^2}$, &c.; when $x = 0$.

First, if $x = 0$, $y = z$; $\therefore U_0 = f(z)$.

$$\text{Now } \frac{du}{dx} = \frac{du}{dy}\cdot\frac{dy}{dx}, \text{ and } \frac{du}{dz} = \frac{du}{dy}\cdot\frac{dy}{dz}.$$

$$\text{But } \frac{dy}{dx} = x\,.\,\phi'(y)\,.\,\frac{dy}{dx} + \phi(y), \text{ where } \phi' y = \frac{d\phi(y)}{dy};$$

$$\therefore \frac{dy}{dx} = \frac{\phi(y)}{1 - x\phi'(y)},$$

$$\frac{dy}{dz} = 1 + x\,.\,\phi'(y)\,.\,\frac{dy}{dz}; \quad \therefore \frac{dy}{dz} = \frac{1}{1 - x\phi'(y)};$$

$$\therefore \frac{dy}{dx} = \phi(y)\,.\,\frac{dy}{dz};$$

$$\therefore \frac{du}{dx} = \frac{du}{dy}\,.\,\phi(y)\,.\,\frac{dy}{dz} = \phi(y)\,.\,\frac{du}{dy}\,.\,\frac{dy}{dz} = \phi(y)\,.\,\frac{du}{dz}.$$

$$\text{Make } x = 0; \quad \therefore U_1 = \phi(z)\,.\,\frac{d\,.\,f(z)}{dz}.$$

$$\text{Next, let } \phi(y)\frac{du}{dz} = \frac{du_1}{dz}; \quad \therefore \frac{du}{dx} = \frac{du_1}{dz};$$

$$\therefore \frac{d^2u}{dx^2} = \frac{d^2u_1}{dx\,dz} = \frac{d^2u_1}{dz\,dx} = \frac{d\,.\left(\frac{du_1}{dx}\right)}{dz} = \frac{d\,.\left\{\phi(y)\,.\left(\frac{du_1}{dz}\right)\right\}}{dz}$$

$$= \frac{d\,.\left\{\{\phi(y)\}^2\,.\,\frac{du}{dz}\right\}}{dz}$$

$$U_2 = \frac{d\left\{\{\phi(z)\}^2\,.\,\frac{d\,.\,f(z)}{dz}\right\}}{dz}.$$

And so may U_3 be found; but to find U_n,

$$\text{assume } \frac{d^{n-1}u}{dx^{n-1}} = \frac{d^{n-2}\left\{\{\phi(y)\}^{n-1}\,.\,\frac{du}{dz}\right\}}{dz^{n-2}};$$

$$\text{let } \{\phi(y)\}^{n-1}\,.\,\frac{du}{dz} = \frac{du_{n-1}}{dz}; \quad \therefore \frac{d^{n-1}u}{dx^{n-1}} = \frac{d^{n-1}u_{n-1}}{dz^{n-1}};$$

$$\therefore \frac{d^nu}{dx^n} = \frac{d^nu_{n-1}}{dx\,.\,dz^{n-1}} = \frac{d^nu_{n-1}}{dz^{n-1}\,dx} = \frac{d^{n-1}\,.\left(\frac{du_{n-1}}{dx}\right)}{dz^{n-1}}$$

$$= \frac{d^{n-1}.\left\{\{\phi(y)\}.\frac{du_{n-1}}{dx}\right\}}{dx^{n-1}} = \frac{d^{n-1}.\left\{\{\phi(y)\}^n.\frac{du}{dx}\right\}}{dx^{n-1}};$$

$$\therefore U_n = \frac{d^{n-1}.\left\{\{\phi(x)\}^n.\frac{d.f(x)}{dx}\right\}}{dx^{n-1}}.$$

Hence if the assumption be true for $n-1$, it is true for n; and it is true for $n=1$ and $n=2$; therefore it is universally true, and writing Z for $\frac{d.f(x)}{dx}$, we have

$$u = f(x) + \{\phi(x).Z\}.\frac{x}{1} + \frac{d.\{[\phi(x)]^2.Z\}}{dx}.\frac{x^2}{1.2}$$

$$+ \frac{d^2.\{[\phi(x)]^3.Z\}}{dx^2}.\frac{x^3}{2.3} + \&c. + \frac{d^{n-1}.\{[\phi(x)]^n.Z\}}{dx^{n-1}}.\frac{x^n}{1.2.3.n}$$

$$+ \&c. \ldots\ldots (1),$$

which is the theorem required.

Cor. If $f(y) = y$ or y be required, then

$$f(x) = x, \text{ and } Z = \frac{df(x)}{dx} = 1;$$

$$\therefore y = x + \phi(x).\frac{x}{1} + \frac{d.\{\phi(x)\}^2}{dx}.\frac{x^2}{1.2}$$

$$+ \frac{d^2\{\phi(x)\}^3}{dx^2}.\frac{x^3}{2.3} + \&c. \ldots\ldots (2).$$

Ex. 1. $y^3 - ay + b = 0$; find y or the root of the cubic equation.

Here $y = \frac{b}{a} + \frac{1}{a}.y^3$, and taking series (2),

$$\therefore x = \frac{b}{a}, \quad x = \frac{1}{a}, \quad \phi(y) = y^3;$$

$$\therefore \phi(x) = x^3, \ \{\phi(x)\}^2 = x^6, \ \{\phi(x)\}^3 = x^9, \ \{\phi(x)\}^4 = x^{12};$$

$$\therefore \frac{d.\{\phi(x)\}^2}{dx} = 6x^5, \ \frac{d^2\{\phi(x)\}^3}{dx^2} = 8.9x^7,$$

$$\frac{d^3\{\phi(x)\}^4}{dx^3} = 10.11.12x^9;$$

$$\therefore\ y = x + x^3 . \frac{x}{1} + 6 . x^5 . \frac{x^2}{1 . 2} + 8 . 9 x^7 . \frac{x^3}{2 . 3}$$

$$+ \frac{10 . 11 . 12}{2 . 3 . 4} . x^9 . x^4 + \&c.$$

$$= \frac{b}{a}\left\{1 + \frac{b^2}{a^3} + 3 . \frac{b^4}{a^6} + 12 \frac{b^6}{a^9} + 55 . \frac{b^8}{a^{12}} + \&c.\right\}.$$

Ex. 2. In the same example, find y^n.

Here $Z = n x^{n-1}$, $\phi(x) = x^3$, and using series (1);

$$\therefore\ \phi(x) . Z = n x^{n+2},$$

$$\{\phi(x)\}^2 Z = n . x^{n+5};\quad \therefore\ \frac{d\{\phi(x)\}^2 . Z}{dx} = n . (n+5) . x^{n+4},$$

$$\{\phi(x)\}^3 Z = n . x^{n+8};\quad \therefore\ \frac{d^2\{\phi(x)\}^2 . Z}{dx^2} = n . (n+8) . (n+7) . x^{n+6};$$

$$\therefore\ y^n = x^n + n x^{n+2} . \frac{x}{1} + \frac{n(n+5)}{1 . 2} . x^{n+4} . x^2$$

$$+ \frac{n . (n+8) . (n+7)}{1 . 2 . 3} x^{n+6} . x^3 + \&c.$$

$$= \frac{b^n}{a^n}\left\{1 + n . \frac{b^2}{a^2} . \frac{1}{a} + \frac{n(n+5)}{1 . 2} . \frac{b^4}{a^4} . \frac{1}{a^2}\right.$$

$$\left. + \frac{n . (n+8)(n+7)}{1 . 2 . 3} . \frac{b^2}{a^6} . \frac{1}{a^3} + \&c.\right\}$$

Ex. 3. Find $\log y$, when $1 - y + a^y = 0$.

$$y = 1 + a^y, \text{ and } u = \log y;$$

$$\therefore\ x = 1,\quad x = 1,\quad \phi(y) = a^y,\quad f(x) = \log x;\quad \therefore\ Z = \frac{1}{x};$$

$$\therefore\ f(x) = 0;\quad \phi(x) . Z = a^x . \frac{1}{x} = a;\quad \{\phi(x)\}^2 Z = a^{2x} \frac{1}{x};$$

$$\therefore\ \frac{d\{\phi(x)\}^2 . Z}{dx} = 2A . a^{2x} . \frac{1}{x} - \frac{a^{2x}}{x^2} = 2A a^2 - a^2;$$

$$\text{and } \{\phi(x)\}^3 . Z = a^{3x} . \frac{1}{x} . \frac{d . \{\phi(x)\}^3 Z}{dx} = 3A a^{3x} . \frac{1}{x} - \frac{a^{3x}}{x^2},$$

$$\frac{d^2.\{\phi(x)\}^3.Z}{dx^2} = 9A^2a^{3x}.\frac{1}{x} - \frac{6A.a^{3x}}{x^2} + \frac{2a^{3x}}{x^3}$$

$$= 9A^2a^3 - 6Aa^3 + 2a^3\,;\quad x = 1,$$

$$= a^3(9A^2 - 6A + 2)\,;$$

$$\therefore \log y = a + (2A-1)\frac{a^2}{1.2} + (9A^2 - 6A + 2).\frac{a^3}{1.2.3} + \&c.$$

Ex. 4. Let $y = m + e\sin y$, find y.

Here $x = m$, $\alpha = e$, $\phi(y) = \sin y$, and $f(x) = x$, $Z = 1$;

$$\therefore y = x + \phi(x)\frac{\alpha}{1} + \frac{d\{\phi(x)\}^2}{dx}.\frac{\alpha^2}{1.2} + \frac{d^2\{\phi(x)\}^3}{dx^2}\frac{\alpha^3}{2.3} + \&c.$$

$\phi(x) = \sin x = \sin m$ if $\alpha = 0$; $\therefore \{\phi(x)\}^2 = \sin^2 x$;

$$\therefore \frac{d\{\phi(x)\}^2}{dx} = 2\sin x.\cos x = \sin 2x = \sin 2m \text{ if } \alpha = 0,$$

$$\{\phi(x)\}^3 = \sin^3 x\,;\quad \therefore \frac{d\{\phi(x)\}^3}{dx} = 3\sin^2 x\cos x,$$

$$\frac{d^2\{\phi(x)\}^3}{dx^2} = 6\sin x\cos^2 x - 3\sin^3 x$$

$$= 6\sin x - 9\sin^3 x = \tfrac{3}{4}(3\sin 3x - \sin x)\,;$$

$$\therefore y = m + \sin m.\frac{e}{1} + \sin 2m.\frac{e^2}{1.2} + \tfrac{3}{4}(3\sin 3m - \sin m)\frac{e^3}{2.3}$$

$$+ \&c.$$

Ex. 5. Let $x_1 = ay + by^2 + cy^3 + ey^4 + \&c.$, find y in term of x_1.

Here $y = \frac{x_1}{a} - \frac{b}{a}.(y^2 + \frac{c}{b}y^3 + \frac{e}{b}y^4 + \&c.)$;

$$\therefore x = \frac{x_1}{a};\quad \alpha = -\frac{b}{a};\quad \phi(y) = y^2 + \frac{c}{b}y^3 + \frac{e}{b}y^4 + \&c.;$$

$$\therefore y = \frac{x_1}{a} - \frac{b}{a^3}x_1^2 + \frac{2b^2 - ac}{a^5}x_1^3 - \frac{5b^3 - 5abc + a^2e}{a^7}x_1^4 + \&c.,$$

a general formula for the inversion of series.

Ex. 6. Let $u + \frac{du}{dx}h + \frac{d^2u}{dx^2}\frac{h^2}{1.2} + \frac{d^3u}{dx^3}\frac{h^3}{2.3} + \&c. = 0$;

find h in terms of u, and its differential coefficients.

Put p, q, r, &c. for $\frac{du}{dx}$, $\frac{d^2u}{dx^2}$, $\frac{d^3u}{dx^3}$, &c.;

$$\therefore\ h = -\frac{u}{p} - \frac{1}{p}\left(\frac{qh^2}{2} + \frac{rh^3}{2.3} + \&c.\right);$$

$$\therefore\ s = -\frac{u}{p};\quad x = -\frac{1}{p};\quad \phi(y) = \frac{qh^2}{2} + \frac{rh^3}{2.3} + \&c.;$$

$$\therefore\ h = -\left\{\frac{u}{p} + \frac{q}{p^3}\cdot\frac{u^2}{2} + \frac{3q^2 - pr}{p^5}\frac{u^3}{2.3} + \&c.\right\}.$$

If a be a root of an equation $u = 0$; and x an approximate value of a, so that $x + h = a$; the preceding series may be used to find a near value of the root; and it has been thus used by Lagrange. Thus if $u = x^4 - 2x^2 + 4x - 8$,

$$h = -\left\{\frac{u}{2^2.(x^3 - x + 1)} + \frac{3x^2 - 1}{2^4.(x^3 - x + 1)^3}\cdot\frac{u^2}{2}\right.$$
$$\left. + \frac{21x^4 - 12x^2 - 6x + 3}{2^6.(x^3 - x + 1)^5}\frac{u^3}{2.3} + \&c.\right\};$$

whence if $x = \frac{3}{2}$; $\therefore\ u = -\frac{23}{16}$; $a = x + h = 1.61$ nearly;

and if 1.61 be put for x, a more correct value may be obtained.

END OF THE DIFFERENTIAL CALCULUS.

THE

INTEGRAL CALCULUS.

CHAPTER I.

1. THE Integral Calculus is the inverse of the Differential, its object being to discover the original function from a given relation between the differential coefficients and functions of x and u. At present we shall only consider the case in which the first differential coefficient $\frac{du}{dx}$ is an *explicit* function of x, as $\phi'(x)$, and $u = \phi(x)$ is required.

2. The process by which u is found from $\frac{du}{dx}$ is called *integration*, and when to be performed is expressed by prefixing the symbol $\int_x$.

Thus if $\frac{du}{dx} = \phi(x)$, $u = \int_x . \phi(x) + C$.

Also since if $\frac{du}{dx} = \phi(x)$;

$$\therefore du = \phi(x) . dx,$$

u is found by prefixing the symbol $\int$, thus $u = \int \phi(x) . dx + C$, or since $\int$ is the initial letter of *summa*, the integral has been said to be the sum of the differentials of the function.

Hence $\int \phi(x) . dx$, and $\int_x \phi(x)$ mean the same thing; also since $\int du = u$, we see that $\int$ and d indicate inverse operations.

The letter C, representing a constant quantity, is added, since constant quantities connected with the original function by the sign $\pm$ disappear in differentiation: and therefore, when we return to the original value u, an arbitrary quantity as C must be added, the value of which will be determined by the nature of the Problem.

3. The simplest case is when $\frac{du}{dx} = ax^m$.

$$\text{Let } u = Ax^n + C; \quad \therefore \frac{du}{dx} = nAx^{n-1} = ax^m;$$

$$\therefore a = nA, \text{ and } m = n - 1; \quad \therefore n = m + 1;$$

$$\text{and } A = \frac{a}{n} = \frac{a}{m+1}; \quad \int_x ax^m = \frac{a}{m+1} \cdot x^{m+1} + C;$$

or to integrate a monomial, *add unity to the index, divide by the index so increased, and add a constant.*

Cor. 1. Thus also if $\frac{du}{dx} = ax^{-m} = \frac{a}{x^m}$, $u = -\frac{a}{m-1} \cdot \frac{1}{x^{m-1}} + C$, which is derived from the preceding by writing $-m$ for m.

Cor. 2. The general formula fails when $m = -1$, for then

$$u = \frac{a \cdot x^{1-1}}{1-1} + C = \frac{a}{0} + C.$$

$$\text{But if } m = -1, \quad \frac{du}{dx} = \frac{a}{x} = a \cdot \frac{1}{x}.$$

$$\text{Now } \frac{1}{x} = \frac{d \cdot \log x}{dx}; \quad \therefore \frac{a}{x} = a \cdot \frac{d \cdot (\log x)}{dx};$$

$$\text{and } \therefore a \cdot \int_x \frac{1}{x} = a \cdot \log x + C;$$

however, the true value of $\frac{du}{dx}$ may be derived from the general expression, if C be first determined.

For, suppose $u = 0$ when $x = b$;

$$\therefore 0 = \frac{ab^{m+1}}{m+1} + C, \text{ or } C = -\frac{ab^{m+1}}{m+1};$$

$$\therefore u = a \cdot \frac{x^{m+1} - b^{m+1}}{m+1}, \text{ a fraction of the form } \frac{0}{0},$$

when $m = -1$, and of which the real value is

$$a \log \frac{x}{b} = a \log x - a \log b = a \log x + C;$$

the same value as that which has been just obtained.

4. Since if $u = \log \{f(x)\} = \log (z)$, where $z = f(x)$,

$$\frac{du}{dx} = \frac{\frac{dz}{dx}}{z}; \quad \therefore \int_x \frac{\frac{dz}{dx}}{z} = \log (z) + C.$$

Hence, if there be a fractional expression, in which the numerator is the differential coefficient of the denominator, the integral is the logarithm of the denominator.

Ex. 1. Let $\frac{du}{dx} = \frac{x}{1+x^2} = \frac{1}{2}.\frac{2x}{1+x^2}$;

$$\therefore\ u = \tfrac{1}{2}.\log(1+x^2) = \log\sqrt{1+x^2}.$$

Ex. 2. Let $\frac{du}{dx} = \frac{2x-1}{x^2-x+1}$; $\therefore\ u = \log(x^2-x+1)$.

5. Again, since

$$\frac{dp}{dx} + \frac{dq}{dx} + \frac{dr}{dx} + \&c. = \frac{d}{dx}(p+q+r+\&c.);$$

$$\therefore \int_x \left\{\frac{dp}{dx} + \frac{dq}{dx} + \frac{dr}{dx} + \&c.\right\} = \int_x \frac{d}{dx}(p+q+r+\&c.)$$

$$= p+q+r+\&c.$$

or the integral of the sum of any number of differential coefficients = sum of the integrals of each differential coefficient taken separately.

Ex. 3. Let $\frac{du}{dx} = Ax^m + Bx^n + Cx^p + \&c.$;

$$\therefore\ u = A\int_x x^m + B\int_x x^n + C\int_x x^p + \&c.$$

$$= \frac{A}{m+1}x^{m+1} + \frac{B}{n+1}x^{n+1} + \frac{C}{p+1}x^{p+1} + \&c.$$

6. If $\frac{du}{dx} = z^m.\frac{dz}{dx}$, where z is a function of x, find u.

Since if $u = z^{m+1} + C$, $\frac{du}{dx} = (m+1)z^m.\frac{dz}{dx}$;

$$\therefore \int_x z^m.\frac{dz}{dx} = \frac{z^{m+1}}{m+1} + C;$$

or to integrate a function of this description, *increase the index by unity, divide by the index so increased,* and by the *differential coefficient of the quantity under the index.*

EXAMPLES OF SIMPLE INTEGRATION.

(1) Let $\frac{du}{dx} = ax^3$; $\therefore u = \frac{ax^4}{4}$.

(2) Let $\frac{du}{dx} = \frac{a}{x^2} = ax^{-2}$; $\therefore u = \frac{ax^{-1}}{-1} = -\frac{a}{x}$.

(3) Let $\frac{du}{dx} = ax^{\frac{m}{n}}$; $\therefore u = \frac{n}{m+n} . ax^{\frac{m+n}{n}}$.

(4) Let $\frac{du}{dx} = (ax^n + b)^m x^{n-1}$.

Let $z = ax^n + b$; $\therefore \frac{dz}{dx} = nax^{n-1}$;

$$\therefore \frac{du}{dz} = \frac{du}{dx} . \frac{dx}{dz} = z^m . x^{n-1} . \frac{1}{nax^{n-1}} = \frac{1}{na} z^m;$$

$$\therefore u = \frac{1}{na} \int_z z^m = \frac{z^{m+1}}{na . (m+1)} = \frac{(ax^n + b)^{m+1}}{na . (m+1)}.$$

(5) $\frac{du}{dx} = (ax + b)^m$; $\therefore u = \frac{(ax + b)^{m+1}}{a . (m+1)}$.

(6) $\frac{du}{dx} = (ax^n + b)^m . x^r$, m being a whole number.

Expand $(ax^n + b)^m$ by the binomial, and after having multiplied each term by x^r, integrate them separately.

(7) $\frac{du}{dx} = \frac{x^m}{(a + bx)^n}$, m and n being whole numbers.

Let $a + bx = z$; $\therefore x = \frac{z - a}{b}$;

$$\therefore \frac{x^m}{(a + bx)^n} = \frac{(z - a)^m}{b^m . z^n}; \text{ and } \frac{dx}{dz} = \frac{1}{b};$$

$$\text{but } \frac{du}{dz} = \frac{du}{dx} . \frac{dx}{dz} = \frac{1}{b} . \frac{du}{dx} = \frac{1}{b^{m+1}} . \frac{(z - a)^m}{z^n};$$

$$\therefore u = \frac{1}{b^{m+1}} . \int_z \frac{(z - a)^m}{z^n}.$$

Expand $(x-a)^m$ by the binomial, and integrate each term separately, first dividing by x^n.

(8) $\dfrac{du}{dx} = \dfrac{1}{x^m(a+bx)^n}$, m and n being integers.

For x put $\dfrac{1}{z}$; $\therefore \dfrac{dx}{dz} = -\dfrac{1}{z^2}$,

$$\therefore \frac{du}{dz} = \frac{du}{dx}\cdot\frac{dx}{dz} = -\frac{1}{z^2}\cdot\frac{du}{dx} = -\frac{z^{m+n-2}}{(az+b)^n};$$

$$\therefore u = -\int_z \frac{z^{m+n-2}}{(az+b)^n},$$

which resolves itself into the preceding case.

(9) $$\frac{du}{dx} = \frac{1}{a+bx^2} = \frac{1}{a}\cdot\frac{1}{1+\frac{b}{a}x^2}.$$

Let $z^2 = \dfrac{b}{a}\cdot x^2$; $\therefore z = \sqrt{\dfrac{b}{a}}\cdot x$, and $\dfrac{dz}{dx} = \sqrt{\dfrac{b}{a}}$;

$$\therefore \frac{du}{dz} = \frac{du}{dx}\cdot\frac{dx}{dz} = \frac{1}{a}\cdot\frac{1}{1+z^2}\cdot\sqrt{\frac{a}{b}} = \frac{1}{\sqrt{ab}}\cdot\frac{1}{1+z^2},$$

$$\text{or } u = \frac{1}{\sqrt{ab}}\int_z \cdot\frac{1}{1+z^2}.$$

But if $u = \tan^{-1}z$, $\dfrac{du}{dz} = \dfrac{1}{1+z^2}$;

$$\therefore u = \frac{1}{\sqrt{ab}}\cdot\tan^{-1}z = \frac{1}{\sqrt{ab}}\cdot\tan^{-1}\left(x\sqrt{\frac{b}{a}}\right).$$

(10) $\dfrac{du}{dx} = \dfrac{1}{2+3x^2}$; $\therefore u = \dfrac{1}{\sqrt{6}}\tan^{-1}\left(x\sqrt{\dfrac{3}{2}}\right)$.

(11) $\dfrac{du}{dx} = \dfrac{1}{a+x}$; $\therefore u = \log(a+x)$.

(12) $\dfrac{du}{dx} = \dfrac{x^2}{1+x^3}$; $\therefore u = \log\sqrt[3]{1+x^3}$.

(13) $\dfrac{du}{dx} = (a+bx+cx^2)^m\cdot(b+2cx)$; $\therefore u = \dfrac{(a+bx+cx^2)^{m+1}}{m+1}$.

(14) $\dfrac{du}{dx} = \dfrac{x^6}{1+x^2}$,

when the index of x in the numerator is not less than that in the denominator, divide by the denominator, thus

$$\frac{x^6}{1+x^2} = x^4 - x^2 + 1 - \frac{1}{1+x^2};$$

$$\therefore\ u = \frac{x^5}{5} - \frac{x^3}{3} + x - \tan^{-1}x.$$

7. Integrate the following differential coefficients.

(1) $ax^{\frac{5}{2}}$. (2) $ax^3 + bx^7 + cx^9$. (3) $(ax^3+b)^2 . x^2$.

(4) $(2ax+x^2)^7 . (a+x)$. (5) $\dfrac{x^2}{(a+bx)}$. (6) $\dfrac{3x^2+2x+1}{x^3+x^2+x+2}$.

(7) $\dfrac{1}{1+5x^2}$. (8) $\dfrac{1}{x^2(a+bx)}$. (9) $\left(\dfrac{a}{x}+\dfrac{b}{x^2}+\dfrac{c}{x^3}+\dfrac{e}{x^4}\right)$.

(10) $(ax+bx^2)^2$. (11) $\left(a+bx+\dfrac{c}{x}\right)^2$.

(12) $\left(ax^2+\dfrac{b}{x^2}\right)^2$. (13) $(1+x^2)(1+x)^2 . x^2$.

(14) $\dfrac{(1+x)^2 . (1-x)}{x^2}$. (15) $\dfrac{x^4}{1+x^2}$. (16) $\dfrac{x^3}{(2+x)^2}$.

(17) $\dfrac{x^5}{1+x^2}$. (18) $\dfrac{5}{2x^4+3x^2}$.

(19) $\dfrac{x\sqrt{x}}{1+\sqrt{x}}$. (20) $\dfrac{x^2\sqrt{x}}{1+x}$.

In (19) and (20) for $\sqrt{x}$ put z; the accuracy of the results may be tested by differentiation.

8. These integrals being found, it will be convenient to classify the remaining functions in the following order.

(1) Rational fractions of the form

$$\frac{Ax^{m-1} + Bx^n + Cx^p + \&c.}{A_1x^m + B_1x^n{}_1 + C_1x^p{}_1 + \&c.}.$$

(2) Irrational quantities.

(3) Exponential and logarithmic functions which are of the forms

$$a^x f(a^x), \quad \log(x), \quad \log(p), \quad p^m \log(q).$$

(4) Circular functions which are of the form

$$\sin p, \quad f(\sin p), \quad (\tan x)^m.$$

The methods for the integration of such functions will be given in the four succeeding chapters, and the integrals thus found, will be then applied to find the areas and lengths of curves, and the volumes and surfaces of solids of revolution.

CHAPTER II.

RATIONAL FRACTIONS.

9. EVERY rational fraction may be represented by

$$\frac{Ax^{m-1} + Bx^{m-2} + Cx^{m-3} + \&c.}{A_1x^m + B_1x^{m-1} + C_1x^{m-2} + \&c.},$$

for it is manifest that the index of x in the numerator can by division be made less by unity at least, than that of x in the denominator.

To integrate this fraction we first separate it into others of a more simple form. Now the denominator may be composed 1st of simple factors all different. 2nd. Some of the factors may be equal. 3rd. It may contain quadratic factors, the roots of which are impossible. 4th. It may be an assemblage of all these.

10. First let $\frac{U}{V}$ be a fraction where V is the product of n factors all different, so that

$$V = (x - a_1)(x - a_2)(x - a_3)\ldots(x - a_n).$$

Assume $\frac{U}{V} = \frac{A_1}{(x-a_1)} + \frac{A_2}{(x-a_2)} + \frac{A_3}{(x-a_3)} + \&c. + \frac{A_n}{(x-a_n)}$;

$$\therefore U = A_1(x-a_2).(x-a_3)\ldots(x-a_n) + A_2.(x-a_1).(x-a_3).(x-a_4)$$
$$+ \&c. + A_n.(x-a_1)(x-a_2)\ldots(x-a_{n-1})\ldots\ldots$$

Successively make $x = a_1$, a_2, a_3, &c.; and let U_{a_1}, U_{a_2}, U_{a_3}, &c. be the corresponding values of U;

$$\therefore U_{a_1} = A_1(a_1 - a_2)(a_1 - a_3)\ldots(a_1 - a_n),$$

$$\text{or } A_1 = \frac{U_{a_1}}{(a_1 - a_2)(a_1 - a_3)\ldots\ldots}.$$

Similarly, $A_2 = \frac{U_{a_2}}{(a_2-a_1)(a_2-a_3)}$, &c. and $A_3 = \frac{U_{a_3}}{(a_3-a_1)(a_3-a_2)\ldots}$,

$$\int_x \left(\frac{U}{V}\right) = A_1 \int_x \frac{1}{x - a_1} + A_2 \int_x \frac{1}{x - a_2} + A_3 \int_x \frac{1}{x - a_3} + \&c.$$

$$= A_1 \log(x - a_1) + A_2 \log(x - a_2) + A_3 \log(x - a_3) + \&c.$$

$$= \log (x - a_1)^{A_1} (x - a_2)^{A_2} (x - a_3)^{A_3} \ldots (x - a_n)^{A_n}.$$

11. Let some of the roots be equal, viz. m of them $= a$, or let $(x-a)^m$ be a factor of V.

Let $V = (x-a)^m Q$.

Assume $\frac{U}{V} = \frac{A}{(x-a)^m} + \frac{B}{(x-a)^{m-1}} + \frac{C}{(x-a)^{m-2}} + \&c. + \frac{P}{Q}$;

$\therefore U = AQ + \{B.(x-a) + C.(x-a)^2 + \&c.\}\, Q + P(x-a)^m$.

Let $x = a$, and let U_a, Q_a be the values of U and Q;

$$\therefore U_a = AQ_a, \text{ and } A = \frac{U_a}{Q_a};$$

$$\therefore U - \frac{U_a}{Q_a}.Q = (x-a)\{[B + C.(x-a) + D(x-a)^2 + \&c.]\, Q + P(x-a)^{m-1}\}.$$

Hence, as the right-hand side of the equation is divisible by $(x-a)$, the left-hand side is also, let the division be effected, and let U^1 be the quotient;

$\therefore U^1 = \{B + C.(x-a) + D(x-a)^2 + \&c.\}\, Q + P.(x-a)^{m-1}$.

Again, make $x = a$, and we have $B = \frac{U^1{}_a}{Q_a}$, and proceeding in the same manner we at length arrive at P, which is either constant, or a function of x; if the latter, the case is reduced to that of the preceding article.

To illustrate these methods, we will take two examples*.

Ex. 1. Integrate $\frac{du}{dx} = \frac{x^2 - 7x + 1}{x^3 - 6x^2 + 11x - 6}$

$$= \frac{x^2 - 7x + 1}{(x-1)(x-2)(x-3)}.$$

Let $\frac{x^2 - 7x + 1}{x^3 - 6x^2 + 11x - 6} = \frac{A}{x-1} + \frac{B}{x-2} + \frac{C}{x-3}$;

$\therefore x^2 - 7x + 1 = A(x-2)(x-3) + B(x-1)(x-3) + C.(x-1)(x-2)$.

Let $x = 1$; $\therefore 1 - 7 + 1 = -5 = A(1-2)(1-3) = 2A$; $\therefore A = -\frac{5}{2}$.

$x = 2$; $\therefore 4 - 14 + 1 = -9 = B(2-1)(2-3) = -B$; $\therefore B = 9$;

$x = 3$; $\therefore 9 - 21 + 1 = -11 = C(3-1)(3-2) = 2C$; $\therefore C = -\frac{11}{2}$;

$$\therefore \int_x \frac{U}{V} = -\frac{5}{2} \cdot \int_x \frac{1}{x-1} + 9 \int_x \frac{1}{x-2} - \frac{11}{2} \cdot \int_x \frac{1}{x-3}$$

* In these and the following examples the constant will be omitted.

$$= -\frac{5}{2}\log(x-1) + 9\log(x-2) - \frac{11}{2}\log(x-3)$$

$$= \log\frac{(x-2)^9}{\sqrt{(x-1)^5(x-3)^{11}}}.$$

Ex. 2. Integrate $\frac{du}{dx} = \frac{2x-5}{(x+3)(x+1)^2}$.

Let $\frac{2x-5}{(x+3)(x+1)^2} = \frac{A}{(x+1)^2} + \frac{B}{x+1} + \frac{P}{x+3}$;

$\therefore\ 2x-5 = A.(x+3) + B(x+1)(x+3) + P.(x+1)^2.$

Let $x=-1$; $\therefore\ -7 = A(3-1) = 2A$; $\therefore\ A = -\frac{7}{2}$;

$$\therefore\ 2x-5+\frac{7}{2}(x+3) = \frac{11x+11}{2} = \frac{11}{2}(x+1)$$

$$= B(x+1)(x+3) + P(x+1)^2;$$

$$\therefore\ \frac{11}{2} = B(x+3) + P(x+1).$$

Let $x+1=0$; $\therefore\ \frac{11}{2} = 2B$; $\therefore\ B = \frac{11}{4}$;

$x+3=0$; $\therefore\ \frac{11}{2} = -2P$; $\therefore\ P = -\frac{11}{4}$;

$$\therefore\ \int_x \frac{U}{V} = -\frac{7}{2}\int_x \frac{1}{(x+1)^2} + \frac{11}{4}\int_x \frac{1}{x+1} - \frac{11}{4}\int_x \frac{1}{x+3}$$

$$= \frac{7}{2}.\frac{1}{x+1} + \frac{11}{4}\log(x+1) - \frac{11}{4}\log(x+3)$$

$$= \frac{7}{2}.\frac{1}{x+1} + \frac{11}{4}\log\left(\frac{x+1}{x+3}\right).$$

12. Next, let V contain quadratic factors having impossible roots.

(1) Let V contain two impossible roots only, and let $(x-\alpha)^2+\beta^2$ be the quadratic factor;

$$\therefore\ V = Q.\{(x-\alpha)^2+\beta^2\}.$$

Assume $\therefore\ \frac{U}{V} = \frac{Mx+N}{(x-\alpha)^2+\beta^2} + \frac{P}{Q}$;

$$\therefore\ U = (Mx+N)Q + P\{(x-\alpha)^2+\beta^2\}.$$

Put $x = \alpha + \beta\sqrt{-1}$; $\therefore\ (x-\alpha)^2 + \beta^2 = 0$.

Then U becomes $U_1 + U_2\sqrt{-1}$, and Q becomes $Q_1 + Q_2\sqrt{-1}$.

Substituting and making the sum of the possible quantities = 0, and also the coefficient of $\sqrt{-1} = 0$, M and N may be found. Or if P be first found, subtract $P\{(x-\alpha)^2 + \beta^2\}$ from each side of the equation;

$$\therefore\ U - P\{(x-\alpha)^2 + \beta^2\} = (Mx + N).Q;$$

$$\therefore\ Mx + N = \frac{U - P.\{(x-\alpha)^2 + \beta^2\}}{Q} \text{ is known;}$$

$$\therefore\ \int_x \frac{U}{V} = \int_x \frac{Mx+N}{(x-\alpha)^2+\beta^2} + \int_x \frac{P}{Q}.$$

To integrate $\dfrac{du}{dx} = \dfrac{Mx+N}{(x-\alpha)^2+\beta^2}$, let $x - \alpha = z$;

$$\therefore\ \frac{du}{dz} = \frac{du}{dx} = \frac{Mz + M\alpha + N}{z^2+\beta^2} = \frac{Mz}{z^2+\beta^2} + \frac{M\alpha+N}{z^2+\beta^2};$$

$$\therefore\ u = M\int_z \frac{z}{z^2+\beta^2} + (M\alpha+N)\int_z \frac{1}{z^2+\beta^2}$$

$$= M\log\sqrt{z^2+\beta^2} + \frac{M\alpha+N}{\beta}\tan^{-1}\left(\frac{z}{\beta}\right)$$

$$= M\log\sqrt{(x-\alpha)^2+\beta^2} + \frac{M\alpha+N}{\beta}\tan^{-1}\left(\frac{x-\alpha}{\beta}\right).$$

Cor. If $\alpha = 0$, or $\dfrac{du}{dx} = \dfrac{Mx+N}{x^2+\beta^2}$,

$$u = M\log\sqrt{x^2+\beta^2} + \frac{N}{\beta}\tan^{-1}\frac{x}{\beta}.$$

Ex. 3. Let $\dfrac{du}{dx} = \dfrac{x-3}{x^3+1} = \dfrac{x-3}{(x+1)(x^2-x+1)}$.

Let $\dfrac{x-3}{x^3+1} = \dfrac{A}{x+1} + \dfrac{Mx+N}{x^2-x+1}$;

$$\therefore\ x - 3 = A(x^2 - x + 1) + (x+1)(Mx+N),$$

$$x = -1;\quad \therefore\ -4 = 3A,\ \text{or } A = -\frac{4}{3};$$

$$\therefore\ x - 3 + \frac{4}{3}(x^2 - x + 1) = \frac{4x^2 - x - 5}{3} = \frac{(4x-5)(x+1)}{3}$$

$$= (x+1)(Mx+N);$$

$$\therefore \frac{4x-5}{3} = (Mx+N);$$

$$\therefore \int_x \frac{x-3}{x^3+1} = -\frac{4}{3}\int_x \frac{1}{x+1} + \frac{1}{3}\int_x \frac{4x-5}{x^2-x+1}.$$

To integrate $\frac{du}{dx} = \frac{4x-5}{x^2-x+1} = \frac{4x-5}{\left(x-\frac{1}{2}\right)^2+\frac{3}{4}}$.

Let $x-\frac{1}{2} = z$; $\therefore \frac{du}{dx} = \frac{du}{dz}$; and $4x-5 = 4z-3$;

$$\therefore u = \int_z \frac{4z-3}{z^2+\frac{3}{4}} = 4\int_z \frac{z}{z^2+\frac{3}{4}} - 3\int_z \frac{1}{z^2+\frac{3}{4}}$$

$$= 2\log\left(z^2+\frac{3}{4}\right) - 2\sqrt{3}\tan^{-1}\frac{2z}{\sqrt{3}};$$

$$\therefore \int_x \frac{x-3}{x^3+1} = -\frac{4}{3}\log(x+1) + \frac{2}{3}\log(x^2-x+1) - \frac{2}{\sqrt{3}}\tan^{-1}\frac{2x-1}{\sqrt{3}}$$

$$= \log\left(\frac{\sqrt{x^2-x+1}}{x+1}\right)^{\frac{4}{3}} - \frac{2}{\sqrt{3}}\tan^{-1}\frac{2x-1}{\sqrt{3}}.$$

Ex. 4. Let $\frac{du}{dx} = \frac{1}{(x+1)(x+2)^2(x^2+1)}$.

Let $\frac{U}{V} = \frac{A}{x+1} + \frac{B}{(x+2)^2} + \frac{C}{x+2} + \frac{Mx+N}{x^2+1}$.

$$1 = A.(x+2)^2(x^2+1) + \{B + C.(x+2)\}(x^2+1)(x+1)$$
$$+ (Mx+N).(x+1)(x+2)^2,$$

$x = -2$; $\therefore 1 = B.5.(1-2) = -5B$, i.e. $B = -\frac{1}{5}$,

$x = -1$; $\therefore 1 = A.2 = 2A$; $\therefore A = \frac{1}{2}$.

$$1 - \frac{(x+2)^2(x^2+1)}{2} + \frac{(x^2+1).(x+1)}{5} = C.(x+2)(x^2+1)(x+1)$$
$$+ (Mx+N).(x+1)(x+2)^2,$$

$$\text{or } -\frac{(5x^4+18x^3+23x^2+18x+8)}{10} = (x+2).(x+1)$$

$$\{C.(x^2+1)+(Mx+N)(x+2)\}.$$

Divide both sides by $(x+2).(x+1)$, or x^2+3x+2;

$$\therefore\ -\frac{5x^2+3x+4}{10} = C(x^2+1)+(Mx+N)(x+2).$$

$$\text{Let } x=-2;\ \ \therefore\ -\frac{9}{5}=5C;\ \ \therefore\ C=-\frac{9}{25};$$

$$\therefore\ \frac{9(x^2+1)}{25}-\frac{5x^2+3x+4}{10}=-\frac{(7x^2+15x+2)}{50}=(Mx+N)(x+2),$$

$$\text{or } -\frac{(7x+1).(x+2)}{50} = (Mx+N)(x+2);$$

$$\therefore\ Mx+N=-\frac{7x+1}{50};$$

$$\therefore\ \int_x\frac{U}{V}=\frac{1}{2}\cdot\int_x\frac{1}{x+1}-\frac{1}{5}\int_x\frac{1}{(x+2)^2}-\frac{9}{25}\int_x\frac{1}{x+2}-\frac{1}{50}\cdot\int_x\frac{7x+1}{x^2+1}$$

$$=\frac{1}{2}\log(x+1)+\frac{1}{5}\frac{1}{x+2}-\frac{9}{25}\log(x+2)$$

$$-\frac{7}{50}\log\sqrt{x^2+1}-\frac{1}{50}\tan^{-1}x.$$

13. If there be m quadratic factors, each $=(x-\alpha)^2+\beta^2$, assume

$$\frac{U}{V}=\frac{Mx+N}{\{(x-\alpha)^2+\beta^2\}^m}+\frac{M_1x+N_1}{\{(x-\alpha)^2+\beta^2\}^{m-1}}+\&c.+\frac{P}{Q};$$

$$\therefore\ U=\{Mx+N+(M_1x+N_1)[(x-\alpha)^2+\beta^2]+\&c.\}$$

$$Q+P\{(x-\alpha)^2+\beta^2\}^m;$$

and after determining $(Mx+N)$, by putting $(x-\alpha)^2+\beta^2=0$, and subtracting $(Mx+N).Q$ from U; divide both sides by the factor $(x-\alpha)^2+\beta^2$, and then proceed in a similar manner to find M_1 and N_1.

Ex. 5. Let $\dfrac{U}{V}=\dfrac{1}{(x^2+1)^2(x+1)}$, resolve it into its partial fractions.

$$\frac{U}{V} = \frac{Mx+N}{(x^2+1)^2} + \frac{M_1x+N_1}{x^2+1} + \frac{P}{x+1};$$

$$\therefore U=1=\{(Mx+N)+(M_1x+N_1)(x^2+1)\}(x+1)+P(x^2+1)^2.$$

Let $x=\sqrt{-1}$;

$$\therefore 1=(M\sqrt{-1}+N).(\sqrt{-1}+1)=-M+M\sqrt{-1}+N\sqrt{-1}+N;$$

$$\therefore N-M=1, \text{ and } N+M=0; \quad \therefore N=\tfrac{1}{2}=-M; \quad \therefore M=-\tfrac{1}{2},$$

$$1+\tfrac{1}{2}.(x+1)(x-1)=\frac{x^2+1}{2}=(M_1x+N_1).(x^2+1).(x+1)+P(x^2+1)^2;$$

$$\therefore \tfrac{1}{2}=(M_1x+N_1)(x+1)+P(x^2+1).$$

Let $x=\sqrt{-1}$; $\quad \therefore +\tfrac{1}{2}=(M_1\sqrt{-1}+N_1)(\sqrt{-1}+1)$

$$=-M_1+M_1\sqrt{-1}+N_1\sqrt{-1}+N_1;$$

$$\therefore N_1-M_1=+\tfrac{1}{2}, \text{ and } N_1+M_1=0; \quad \therefore N_1=\tfrac{1}{4}, \text{ and } M_1=-N_1=-\tfrac{1}{4},$$

$$x=-1; \quad \therefore \tfrac{1}{2}=P\times 2; \quad \therefore P=\tfrac{1}{4};$$

$$\therefore \frac{U}{V}=-\tfrac{1}{2}\cdot\frac{x-1}{(x^2+1)^2}-\tfrac{1}{4}\frac{x-1}{x^2+1}+\tfrac{1}{4}\frac{x}{x-1}.$$

14. To integrate the fraction $\frac{x-1}{(x^2+1)^2}$, divide it into two others, $\frac{x}{(x^2+1)^2}$, and $\frac{-1}{(x^2+1)^2}$. From the former we have,

$$\int_x \frac{x}{(x^2+1)^2} = \int_x (x^2+1)^{-2}.x = -\frac{(x^2+1)^{-1}}{2} = -\tfrac{1}{2}\cdot\frac{1}{(x^2+1)};$$

but $\int_x \frac{1}{(x^2+1)^2}$ which is a particular case of $\int_x \frac{1}{(x^2+1)^n}$ must be deduced from that integral, which has not yet been found.

15. Now to integrate $\frac{du}{dx}=\frac{1}{(x^2+1)^n}$, we may assume

$$\int_x \frac{1}{(x^2+1)^n} = \frac{Ax}{(x^2+1)^{n-1}} + B\int_x \frac{1}{(x^2+1)^{n-1}};$$

$$\therefore \frac{1}{(x^2+1)^n} = \frac{A(x^2+1)-(2n-1)Ax^2}{(x^2+1)^n} + \frac{B}{(x^2+1)^{n-1}}$$

$$= \frac{A+B+\{B-(2n-3)A\}x^2}{(x^2+1)^n};$$

$$\therefore A + B = 1,\ B - (2n-3).A = 0;\ A = \frac{1}{2n-2},\ B = \frac{2n-3}{2n-2};$$

$$\therefore \int_x \frac{1}{(x^2+1)^n} = \frac{1}{2n-2}\cdot\frac{x}{(x^2+1)^{n-1}} + \frac{2n-3}{2n-2}\cdot\int_x \frac{1}{(x^2+1)^{n-1}};$$

by this process $\int_x \frac{1}{(x^2+1)^n}$ is made to depend upon $\int_x \frac{1}{(x^2+1)^{n-1}}$, and by substituting $n-1$, $n-2$, and $n-(n-1)$ for n, it will be reduced to $\int_x \frac{1}{x^2+1} = \tan^{-1}x$.

Ex. 6. Let $n = 4$, or let $\int_x \frac{1}{(x^2+1)^4}$ be required.

$$\int_x \frac{1}{(x^2+1)^4} = \frac{1}{6}\cdot\frac{x}{(x^2+1)^3} + \frac{5}{6}\cdot\int_x \frac{1}{(x^2+1)^3},$$

$$\int_x \frac{1}{(x^2+1)^3} = \frac{1}{4}\cdot\frac{x}{(x^2+1)^2} + \frac{3}{4}\cdot\int_x \frac{1}{(x^2+1)^2},$$

$$\int_x \frac{1}{(x^2+1)^2} = \frac{1}{2}\cdot\frac{x}{x^2+1} + \frac{1}{2}\cdot\int_x \frac{1}{x^2+1}$$

$$= \frac{1}{2}\cdot\frac{x}{x^2+1} + \frac{1}{2}\tan^{-1}x;$$

$$\therefore \int_x \frac{1}{(x^2+1)^4} = \frac{1}{6}\cdot\frac{x}{(x^2+1)^3} + \frac{5}{4.6}\cdot\frac{x}{(x^2+1)^2} + \frac{3.5}{2.4.6}\cdot\frac{x}{x^2+1}$$

$$+ \frac{3.5}{2.4.6}\tan^{-1}x.$$

16. To the $\int_x \frac{1}{(x^2+1)^n}$ may also be referred the

$$\int_x \frac{Mx+N}{\{(x-\alpha)^2+\beta^2\}^m}.$$

For, let $x - \alpha = z$; $\therefore \frac{dx}{dz} = 1$; $\therefore \frac{du}{dz} = \frac{du}{dx}$;

$$\therefore \int_x \frac{Mx+N}{\{(x-\alpha)^2+\beta^2\}^m} = \int_z \frac{Mz + (M\alpha+N)}{(z^2+\beta^2)^m}$$

$$= M\int_z \frac{z}{(z^2+\beta^2)^m} + (M\alpha+N)\int_z \frac{1}{(z^2+\beta^2)^m},$$

and $M\int_z \frac{z}{(z^2+\beta^2)^m} = -\frac{M}{2m-2}\cdot\frac{1}{(z^2+\beta^2)^{m-1}}$,

$$\text{and } (Ma+N)\int_x \frac{1}{(x^2+\beta^2)^m} = \frac{Ma+N}{\beta^{2m}}\int_x \frac{1}{\left(\frac{x^2}{\beta^2}+1\right)^m}$$

$$= \frac{Ma+N}{\beta^{2m}} \cdot \int_y \frac{\beta}{(y^2+1)^m} = \frac{Ma+N}{\beta^{2m-1}}\int_y \frac{1}{(y^2+1)^m},$$

by putting $\frac{x}{\beta} = y$, or $x = \beta y$, which can be integrated by the preceding method.

INTEGRATION BY PARTS.

17. The method which has been given for the integration of $\int_x \frac{1}{(x^2+1)^n}$ is commonly superseded by another, called the integration by parts, which is very general in its application, and which we now proceed to explain.

$$\text{Since } \frac{d}{dx}(pq) = p\frac{dq}{dx} + q\frac{dp}{dx};$$

$$\therefore p\frac{dq}{dx} = \frac{d}{dx}(pq) - q\cdot\frac{dp}{dx};$$

$$\therefore \int_x p\frac{dq}{dx} = pq - \int_x q\cdot\frac{dp}{dx},$$

or if any differential coefficient can be divided into two parts, one of which is a function of x as p, and the other is the differential coefficient of a known function q, then u, the required function, is equal to the product of p and q, *minus* the integral of q multiplied by $\frac{dp}{dx}$. The utility of this method depends upon $q\frac{dp}{dx}$ being less complicated then the original function $p\frac{dq}{dx}$.

Ex. 1. Let $\frac{du}{dx} = x^3(1+x^2)^5 = x^2\, x(1+x^2)^5$;

$$\therefore p = x^2; \quad \frac{dp}{dx} = 2x; \quad \frac{dq}{dx} = x(1+x^2)^5; \quad q = \frac{(1+x^2)^6}{12};$$

$$\therefore \int_x x^3(1+x^2)^5 = \frac{x^2(1+x^2)^6}{12} - \frac{1}{6}\int_x x(1+x^2)^6$$

$$= \frac{x^2(1+x^2)^6}{12} - \frac{(1+x^2)^7}{84}$$

$$= \frac{(1+x^2)^6}{12}\left\{x^2 - \frac{1+x^2}{7}\right\}$$

$$= \frac{(1+x^2)^6}{12}\left(\frac{6x^2-1}{7}\right).$$

Ex. 2. Integrate $\frac{du}{dx} = \frac{1}{(x^2+1)^n}$.

$$\frac{1}{(x^2+1)^{n-1}} = \frac{x^2+1}{(x^2+1)^n} = \frac{x^2}{(x^2+1)^n} + \frac{1}{(x^2+1)^n};$$

$$\therefore \int_x \frac{1}{(x^2+1)^n} = \int_x \frac{1}{(x^2+1)^{n-1}} - \int_x \frac{x^2}{(x^2+1)^n}.$$

But $\int_x \frac{x^2}{(x^2+1)^n} = \int_x x \cdot \frac{x}{(x^2+1)^n}$.

$$p = x; \quad \frac{dp}{dx} = 1; \quad \frac{dq}{dx} = \frac{x}{(x^2+1)^n}; \quad q = -\frac{1}{(2n-2)(x^2+1)^{n-1}};$$

$$\therefore \int_x \frac{x^2}{(x^2+1)^n} = \frac{-x}{(2n-2)(x^2+1)^{n-1}} + \frac{1}{2n-2}\cdot\int_x \frac{1}{(x^2+1)^{n-1}};$$

$$\therefore \int_x \frac{1}{(x^2+1)^n} = \frac{x}{(2n-2)(x^2+1)^{n-1}} + \left(1 - \frac{1}{2n-2}\right)\int_x \frac{1}{(1+x^2)^{n-1}}$$

$$= \frac{x}{(2n-2)(x^2+1)^{n-1}} + \frac{2n-3}{2n-2}\cdot\int_x \frac{1}{(1+x^2)^{n-1}};$$

the formula of reduction which was obtained in Art. 15.

Ex. 3. To integrate $\frac{du}{dx} = \frac{x^m}{(1+x^2)^n}$.

$$\int_x \frac{x^m}{(1+x^2)^n} = \int_x x^{m-1}\frac{x}{(1+x^2)^n};$$

$$\therefore p = x^{m-1}; \quad \frac{dp}{dx} = (m-1)\cdot x^{m-2}; \quad q = \frac{-1}{(2n-2)(1+x^2)^{n-1}};$$

$$\therefore \int_x \frac{x^m}{(1+x^2)^n} = \frac{-x^{m-1}}{(2n-2)(1+x^2)^{n-1}} + \frac{m-1}{2n-2}\int_x \frac{x^{m-2}}{(1+x^2)^{n-1}};$$

a formula of reduction by which the original integral may be made to depend upon

$$\int_x \frac{x}{(1+x^2)^{n_1}} \text{ or } \int \frac{1}{(1+x^2)^{n_1}},$$

according as m is an odd or even integer.

Ex. 4. Integrate $\frac{du}{dx} = \frac{x^5}{(1+x^2)^4}$;

$$\therefore \int_x \frac{x^5}{(1+x^2)^4} = \frac{-x^4}{8(1+x^2)^3} + \frac{4}{8} \cdot \int_x \frac{x^3}{(1+x^2)^3};$$

$$\int_x \frac{x^3}{(1+x^2)^3} = \frac{-x^2}{6(1+x^2)^2} + \frac{2}{6} \cdot \int \frac{x}{(1+x^2)^2}$$

$$= \frac{-x^2}{6(1+x^2)^2} - \frac{1}{6} \frac{1}{1+x^2};$$

$$\therefore \int_x \frac{x^5}{(1+x^2)^4} = -\left\{\frac{x^4}{8(1+x^2)^3} + \frac{4x^2}{6.8(1+x^2)^2} + \frac{4}{6.8.(1+x^2)}\right\}$$

$$= -\frac{1}{8(1+x^2)} \cdot \left\{\frac{x^4}{(1+x^2)^2} + \frac{2x^2}{3(1+x^2)} + \frac{2}{3}\right\}$$

$$= -\frac{4+6x^2+5x^4}{24 \,.\, (1+x^2)^3}.$$

18. Next, to integrate functions of the form

$$\frac{x^m}{(a+bx+cx^2)^n} \text{ and } \frac{1}{x^m(a+bx+cx^2)^n}.$$

In these cases the trinomial $a+bx+cx^2$ must be reduced to a binomial; and then the integration may be effected by methods already given: we will first however shew how the function may be integrated when $m=0$ and $n=1$.

19. Integrate $\frac{du}{dx} = \frac{1}{a+bx+cx^2}$.

$$\frac{1}{a+bx+cx^2} = \frac{1}{c} \frac{1}{\left(\frac{a}{c} + \frac{b}{c}x + x^2\right)}.$$

Let $x + \frac{b}{2c} = z$; $\therefore \frac{dx}{dz} = 1$,

$$\text{and } x^2 + \frac{bx}{c} + \frac{a}{c} = z^2 + \frac{a}{c} - \frac{b^2}{4c^2};$$

$$\therefore \frac{du}{dz} = \frac{du}{dx} = \frac{1}{c\left(z^2 + \frac{a}{c} - \frac{b^2}{4c^2}\right)}.$$

(1) Let $\frac{a}{c} > \frac{b^2}{4c^2}$, or $4ac > b^2$;

$$\therefore u = \frac{1}{c} \cdot \int_z \frac{1}{z^2 + \frac{4ac - b^2}{4c^2}}.$$

But $\because \int_z \frac{1}{z^2 + a^2} = \frac{1}{a} \cdot \tan^{-1}\frac{z}{a}$;

$$\therefore = \frac{1}{c} \cdot \frac{2c}{\sqrt{4ac - b^2}} \cdot \tan^{-1}\left(\frac{2cz}{\sqrt{4ac - b^2}}\right)$$

$$= \frac{2}{\sqrt{4ac - b^2}} \cdot \tan^{-1}\frac{2cx + b}{\sqrt{4ac - b^2}}.$$

(2) Let $\frac{a}{c} < \frac{b^2}{4c^2}$, make $a^2 = \frac{b^2 - 4ac}{4c^2}$;

$$\therefore u = \frac{1}{c} \cdot \int_z \frac{1}{z^2 - a^2} = \frac{1}{2ca} \cdot \int_z \left(\frac{1}{z - a} - \frac{1}{z + a}\right)$$

$$= \frac{1}{2ca} \cdot \log\left(\frac{z - a}{z + a}\right)$$

$$= \frac{1}{\sqrt{b^2 - 4ac}} \cdot \log \frac{2cx + b - \sqrt{b^2 - 4ac}}{2cx + b + \sqrt{b^2 - 4ac}}.$$

Ex. 1. Let $a = b = c = 1$; $\therefore \sqrt{4ac - b^2} = \sqrt{3}$,

$$\int_x \frac{1}{1 + x + x^2} = \frac{2}{\sqrt{3}} \tan^{-1}\frac{2x + 1}{\sqrt{3}}.$$

Ex. 2. Let $c = b = 1$, and $a = -1$; $\therefore \sqrt{b^2 - 4ac} = \sqrt{5}$;

$$\therefore \int_x \frac{1}{x^2 + x - 1} = \frac{1}{\sqrt{5}} \log\left(\frac{2x + 1 - \sqrt{5}}{2x + 1 + \sqrt{5}}\right).$$

20. To integrate $\frac{x^m}{(a + bx + cx^2)^n}$.

$$\frac{x^m}{(a+bx+cx^2)^n} = \frac{1}{c^n} \cdot \frac{x^m}{\left(x^2 + \frac{b}{c}x + \frac{a}{c}\right)^n}.$$

Let $x + \frac{b}{2c} = z$, or $x + a = z$, if $a = \frac{b}{2c}$;

$$\therefore x^2 + \frac{b}{c}x + \frac{a}{c} = z^2 + \frac{a}{c} - \frac{b^2}{4c^2} = (z^2 \pm \beta^2);$$

$$\therefore \int_x \frac{x^m}{(a+bx+cx^2)^n} = \frac{1}{c^n} \int_z \frac{(z-\alpha)^m}{(z^2 \pm \beta^2)^n}.$$

Here are two cases:

(1) Let $\frac{a}{c} > \frac{b^2}{4c^2}$; then $\int_z \frac{(z-\alpha)}{(z^2+\beta)^n}$ may be found by the method used in Art. 15, or Art. 17,

(2) Let $\frac{a}{c} < \frac{b^2}{4c^2}$; $\therefore \int_z \frac{(z-\alpha)^m}{(z^2-\beta^2)^n} = \int_z \frac{(z-\alpha)^m}{(z+\beta)^n (z-\beta)^n}$

must be integrated by the method of partial fractions.

21. Again, to integrate $\frac{1}{x^m(a+bx+cx^2)^n}$.

Let $x = \frac{1}{z}$; $\therefore \frac{dx}{dz} = -\frac{1}{z^2}$;

$$\therefore \frac{du}{dz} = -\frac{1}{z^2} \cdot \frac{du}{dx} = -\frac{1}{z^2} \cdot \frac{z^{m+2n}}{(az^2+bz+c)^n};$$

$$\therefore u = -\int_z \frac{z^{m+2n-2}}{(az^2+bz+c)^n},$$

which is a case of the preceding article.

22. To integrate $\frac{du}{dx} = \frac{1}{x^n - 1}$; and $\frac{du}{dx} = \frac{1}{x^n + 1}$.

Since when n is an even number,

$$x^n - 1 = (x-1)(x+1)\left(x^2 - 2x\cos\frac{2\pi}{n} + 1\right)\left(x^2 - 2x\cos\frac{4\pi}{n} + 1\right)\ldots$$

continued to the factor $x^2 - 2x\cos\left(\frac{n-2}{n}\right)\pi + 1$;

and when n is odd,

$$x^n - 1 = (x-1)\left(x^2 - 2x\cos\frac{2\pi}{n} + 1\right)\left(x^2 - 2x\cos\frac{4\pi}{n} + 1\right)\ldots$$

continued to the factor $x^2 - 2x\cos\dfrac{n-1}{n}\pi + 1$;

and $\because$ the factors of $x^n + 1 = 0$ are contained in

$$x^2 - 2x\cos\frac{2m+1}{n}\pi + 1,$$

we may integrate these differential coefficients by resolving them into partial fractions, having simple and quadratic factors for their denominators: to effect this the following process may be used.

Case 1. Let n be even, then since

$$x^n - 1 = (x-1)(x+1)\left(x^2 - 2x\cos\frac{2m\pi}{n} + 1\right),$$

where $x^2 - 2x\cos\dfrac{2m\pi}{n} + 1$ represents all the quadratic factors;

$$\therefore \log(x^n - 1) = \log(x-1) + \log(x+1) + \log\left(x^2 - 2x\cos\frac{2m\pi}{n} + 1\right);$$

$$\therefore \frac{nx^{n-1}}{x^n - 1} = \frac{1}{x-1} + \frac{1}{x+1} + \frac{2x - 2\cos\dfrac{2m\pi}{n}}{x^2 - 2x\cos\dfrac{2m\pi}{n} + 1};$$

$$\therefore \frac{nx^n}{x^n - 1} = \frac{x}{x-1} + \frac{x}{x+1} + \frac{2x^2 - 2x\cos\dfrac{2m\pi}{n}}{x^2 - 2x\cos\dfrac{2m\pi}{n} + 1}.$$

Now subtract n from the left-hand side of the equation, and on the right side, *unity* from each simple factor, and *two* from each quadratic factor;

$$\therefore \frac{n}{x^n - 1} = \frac{1}{x-1} + \frac{1}{x+1} - \frac{2 - 2x\cos\dfrac{2m\pi}{n}}{x^2 - 2x\cos\dfrac{2m\pi}{n} + 1};$$

$$\therefore \int_x \frac{1}{x^n - 1} = \frac{1}{n}\log(x^2 - 1) - \frac{2}{n}\int_x \frac{1 - x\cos\dfrac{2m\pi}{n}}{x^2 - 2x\cos\dfrac{2m\pi}{n} + 1}.$$

The last integral is of the form $\int_z \frac{1-\beta x}{x^2-2\beta x+1}$, and is, if we make $x-\beta=z$, $1-\beta^2=\delta^2$

$$=\delta\tan^{-1}\frac{z}{\delta}-\beta\log\sqrt{z^2+\delta^2},\text{ or since }\delta=\sin\frac{2m\pi}{n},$$

$$\int_x\frac{1}{x^n-1}=\frac{1}{n}\log(x^2-1)-\frac{2}{n}\left\{\sin\frac{2m\pi}{n}\tan^{-1}\left(\frac{x-\cos\frac{2m\pi}{n}}{\sin\frac{2m\pi}{n}}\right)\right.$$

$$\left.-\cos\frac{2m\pi}{n}\log\sqrt{x^2-2x\cos\frac{2m\pi}{n}+1}\right\}.$$

The method is the same when n is odd.

The same method applies to $\frac{du}{dx}=\frac{1}{x^n+1}$; and n odd,

$$x^n+1=(x+1)\left(x^2-2x\cos\frac{2m+1}{n}\pi+1\right);$$

$$n\text{ even, }x^n+1=\left(x^2-2x\cos\frac{2m+1}{n}\pi+1\right);$$

whence giving proper values to m, u may be found.

23. To integrate the functions $\frac{x^r}{x^n-1}$ and $\frac{x^r}{x^n+1}$.

$$\text{Let }\frac{x^r}{x^n-1}=\frac{Mx+N}{x^2-2x\cos\left(\frac{2m\pi}{n}\right)+1}+\frac{P}{Q};$$

$$\therefore\ x^r=(Mx+N).Q+P\left\{x^2-2x\cos\left(\frac{2m\pi}{n}\right)+1\right\}.$$

$$\text{Let }x=\cos\frac{2m\pi}{n}+\sqrt{-1}\sin\frac{2m\pi}{n},\text{ and}$$

let Q_1 and z be corresponding values of Q and x;

$$\therefore\ z^r=(Mz+N)\,Q_1.$$

But $x^n-1=Q\left(x^2-2x\cos\frac{2m\pi}{n}+1\right)$;

$$\therefore\ nx^{n-1} = Q\left(2x - 2\cos\frac{2m\pi}{n}\right) + (x^2 - 2x\cos\frac{2m\pi}{n} + 1)\cdot\frac{dQ}{dx};$$

$$\therefore\ nx^{n-1} = Q_1\left(2x - 2\cos\frac{2m\pi}{n}\right);$$

$$\therefore\ nx^n = n = Q_1\left(2x^2 - 2x\cos\frac{2m\pi}{n}\right) = Q_1(x^2 - 1);$$

$$\therefore\ Q_1 = \frac{n}{x^2 - 1};\quad \text{and } x^r = (Mx + N)\cdot\frac{n}{x^2 - 1},$$

$$\text{or } n(Mx + N) = x^r(x^2 - 1) = x^{r+1}\cdot\left(x - \frac{1}{x}\right);$$

whence substituting for x,

$$Mn\cdot\cos\frac{2mp}{n} + Mn\sqrt{-1}\cdot\sin\frac{2m\pi}{n} + Nn$$

$$= \left\{\cos\left(\frac{2m\cdot(r+1)\pi}{n}\right) + \sqrt{-1}\cdot\sin\left(\frac{2m(r+1)\pi}{n}\right)\right\}2\sqrt{-1}\cdot\sin\frac{2m\pi}{n}$$

$$= 2\sqrt{-1}\cos\frac{2m\cdot(r+1)\pi}{n}\cdot\sin\frac{2m\pi}{n} - 2\sin\frac{2m(r+1)\pi}{n}\cdot\sin\frac{2m\pi}{n};$$

$$\therefore\ M = \frac{2}{n}\cdot\cos\left(\frac{2m\cdot(r+1)\pi}{n}\right),$$

$$N = -\frac{2}{n}\cdot\sin\frac{2m\cdot(r+1)\pi}{n}\cdot\sin\frac{2m\pi}{n}$$

$$-\frac{2}{n}\cdot\cos\frac{2m(r+1)\pi}{n}\cdot\cos\frac{2m\pi}{n}$$

$$= -\frac{2}{n}\cdot\cos\frac{2mr\cdot\pi}{n};$$

$$\therefore\ \frac{Mx + N}{x^2 - 2x\cdot\cos\frac{2m\pi}{n} + 1} = \frac{2}{n}\cdot\frac{x\cdot\cos\frac{2m\cdot(r+1)\pi}{n} - \cos\frac{2mr\cdot\pi}{n}}{x^2 - 2x\cdot\cos\frac{2m\pi}{n} + 1}.$$

CASE 1. Let n be odd; $\therefore Q = x - 1$, and $P = A$;

$$\text{also } x^r = f(Mx + N)\cdot(x - 1) + A\cdot\frac{x^n - 1}{x - 1}.$$

$$\text{Let } x = 1\,; \quad \therefore \frac{x^n - 1}{x - 1} = n, \text{ and } A = \frac{1}{n}\,;$$

$$\therefore \int_x \frac{x^r}{x^r - 1} = \frac{1}{n} \cdot \int_x \frac{1}{x - 1} + \frac{2}{n} \cdot \int_x \frac{x\cos\frac{2m.(r+1)\pi}{n} - \cos\frac{2mr.\pi}{n}}{x^2 - 2x\cos\frac{2m\pi}{n} + 1}\,;$$

the latter integral is of the form $\int_x \frac{Mx + N}{x^2 - 2ax + 1}$

$$= M\log\sqrt{x^2 - 2ax + 1} + \frac{Ma + N}{\sqrt{1 - a^2}} \cdot \tan^{-1}\left(\frac{x - a}{\sqrt{1 - a^2}}\right);$$

$$\text{but } a = \cos\frac{2m\pi}{n}\,; \quad \therefore \sqrt{1 - a^2} = \sin\frac{2m\pi}{n}.$$

$$\text{Also } \frac{Ma + N}{\sqrt{1 - a^2}} = -\frac{2}{n} \cdot \sin\frac{2m(r+1) \cdot \pi}{n},$$

$$\therefore \int_x \frac{x^r}{x^n - 1} = \frac{1}{n} \cdot \int_x \frac{1}{(x - 1)} + \frac{2}{n}\left\{\cos \cdot \frac{2m(r+1)\pi}{n}\right.$$

$$\log\sqrt{x^2 - 2x\cos\frac{2m\pi}{n} + 1} - \sin\frac{2m(r+1)\pi}{n}$$

$$\left.\tan^{-1}\left(\frac{x - \cos\frac{2m\pi}{n}}{\sin\frac{2m\pi}{n}}\right)\right\},$$

where m must be taken from $m = 1$ to $m = \frac{n-1}{2}$.

Cor. 1. If $r = 0$, we have $\int_x \frac{1}{x^n - 1}$

$$= \frac{1}{n} \cdot \log(x - 1) + \frac{2}{n} \cdot \cos\frac{2m\pi}{n} \log\sqrt{x^2 - 2x\cos\frac{2m\pi}{n} + 1}$$

$$- \frac{2}{n} \cdot \sin\frac{2m\pi}{n} \cdot \tan^{-1}\frac{x - \cos\frac{2m\pi}{n}}{\sin\frac{2m\pi}{n}}.$$

24. If we add to the integral the constant quantity

$$-\sin\left(\frac{2m(r+1)\cdot\pi}{n}\right)\tan^{-1}\left(\frac{-\cos\left(\frac{2m\pi}{n}\right)}{\sin\frac{2m\pi}{n}}\right).$$

Since $\tan^{-1}A-\tan^{-1}B=\tan^{-1}\left(\frac{A-B}{1+AB}\right)$;

$$\therefore\int_s\frac{x^r}{x^n-1}=\frac{1}{n}\log(x-1)+\frac{2}{n}\cdot\cos\left(\frac{2m\cdot(r+1)\pi}{n}\right)$$

$$\log\sqrt{x^2-2x\cos\frac{2m\pi}{n}+1}-\frac{2}{n}\cdot\sin\frac{2m(r+1)\pi}{n}$$

$$\tan^{-1}\left\{\frac{x\sin\frac{2m\pi}{n}}{1-x\cos\frac{2m\pi}{n}}\right\}.$$

25. And $\therefore\int_s\frac{1}{x^n-1}=\frac{1}{n}\log(x-1)+\frac{2}{n}\cos\left(\frac{2m\pi}{n}\right)$

$$\log\sqrt{x^2-2x\cos\frac{2m\pi}{n}+1}-\frac{2}{n}\cdot\sin\frac{2m\pi}{n}$$

$$\tan^{-1}\left\{\frac{x\sin\frac{2m\pi}{n}}{1-x\cos\frac{2m\pi}{n}}\right\}.$$

Ex. $\int\frac{1}{x^5-1}=\frac{1}{5}\log(x-1)+\frac{2}{5}\cos\frac{2\pi}{5}$

$$\log\sqrt{x^2-2x\cos\frac{2\pi}{5}+1}-\frac{2}{5}\sin\frac{2\pi}{5}\tan^{-1}\left\{\frac{x\sin\frac{2\pi}{5}}{1-x\cos\frac{2\pi}{5}}\right\}$$

$$+\frac{2}{5}\cos\frac{4\pi}{5}\log\sqrt{x^2-2x\cos\frac{4\pi}{5}+1}$$

$$-\frac{2}{5}\sin\frac{4\pi}{5}\tan^{-1}\left\{\frac{x\sin\frac{4\pi}{5}}{1-x\cos\frac{4\pi}{5}}\right\}+C.$$

26. If n be even, there will be two terms of the form $\frac{A}{x-1}$ and $\frac{B}{x+1}$; and A and B may be found by putting $+1$ and -1 for x in the equation

$$x^r = f(Mx+N)(x-1)+A\frac{x^n-1}{x-1},$$

$$\text{and } x^r = f(Mx+N)(x+1)+B\frac{x^n-1}{x+1};$$

$$\therefore A=\frac{1}{n}, \text{ and } B=\pm\frac{1}{n}.$$

27. Also the function $\frac{x^r}{x^n+1}$, since the quadratic factors of the denominator are included in the general formula, $x^2-2x\cos\left(\frac{2m+1}{n}\right)\pi+1$, may be integrated in a similar manner, and will be found to depend upon the terms

$$\frac{2}{n}\left\{\cos\frac{(n-r-1)\{(2m+1)\}\pi}{n}\log\sqrt{x^2-2x\cos\frac{2m+1}{n}\pi+1}\right.$$

$$\left.+\sin\left\{\frac{(n-r-1)(2m+1)\pi}{n}\right\}\tan^{-1}\frac{x\sin\left(\frac{2m+1}{n}\pi\right)}{1-x\cos\left(\frac{2m+1}{n}\pi\right)}\right\}+C.$$

If n be even, m must be taken from $m=0$ to $m=\frac{n}{2}$.

If n be odd.....$m=1$ to $m=\frac{n-1}{2}$,

there will be a simple factor $(x+1)$, and a fraction $\frac{A}{x+1}$;

where $A=\frac{(-1)^r}{n}$.

Cor. Hence $\int_a \frac{1}{x^n+1}$ will depend upon

$$\frac{2}{n}.\sin\left\{\frac{(n-1).(2m+1).\pi}{n}\right\}.\log\sqrt{x^2-2x\cos\left(\frac{2m+1}{n}\right)\pi+1}$$

$$-\frac{2}{n}.\sin\frac{(n-1).(2m+1).\pi}{n}\tan^{-1}\left\{\frac{x.\sin\frac{(2m+1)}{n}\pi}{1-x\cos\left(\frac{2m+1}{n}\pi\right)}\right\}+C.$$

28. To integrate $\frac{x^r}{x^{2n}-2x^n\cos\alpha+1}$.

The quadratic factors of the denominator will be all comprised in the term $x^2-2x\cos\theta+1$, where $\theta=\frac{2m\pi+\alpha}{n}$.

Let $\frac{x^r}{x^{2n}-2x^n.\cos\alpha+1}=\frac{Mx+N}{x^2-2x\cos\theta+1}+\frac{P}{Q}$;

$\therefore\ x^r=(x^2-2x\cos\theta+1)P+(Mx+N).Q.$

Let $\cos\theta+\sqrt{-1}\sin\theta=z$, and Q_1 be the value of Q;

$$\therefore\ z^r=(Mz+N).Q_1.$$

Now $x^{2n}-2x^n\cos\alpha+1=(x^2-2x\cos\theta+1).Q$;

$$\therefore\ 2n.x^{2n-1}-2n.x^{n-1}\cos\alpha=(2x-\cos\theta)Q+(x^2-2x\cos\theta+1)\frac{dQ}{dx}$$

$$2nz^{2n-1}-2nz^{n-1}\cos\alpha=(2z-2\cos\theta).Q_1,$$

$$\text{or } 2nz^{2n}-2nz^n\cos\alpha=n(z^{2n}-1)=(2z^2-2z\cos\theta).Q_1$$

$$=(z^2-1).Q_1;$$

$$\therefore\ Q_1=\frac{n(z^{2n}-1)}{z^2-1};$$

$$\therefore\ z^r=(Mz+N).n\frac{z^{2n}-1}{z^2-1}=n.(Mz+N).\frac{z^{2n-1}-\frac{1}{z}}{z-\frac{1}{z}};$$

$$\therefore\ z^{r-n+1}=n(Mz+N).\frac{\left(z^n-\frac{1}{z^n}\right)}{z-\frac{1}{z}},$$

or $\cos\{(r-n+1)\}\theta + \sqrt{-1}\sin(r-n+1)\theta$

$$= n\{M(\cos\theta + \sqrt{-1}\sin\theta) + N\} \times \frac{\sin n\theta}{\sin\theta}$$

$$= n.\left(M\frac{\cos\theta\sin n\theta}{\sin\theta} + M\sqrt{-1}.\sin n\theta + N.\frac{\sin n\theta}{\sin\theta}\right);$$

$$\therefore M = \frac{1}{n}.\frac{\sin(r-n+1)\theta}{\sin n\theta} = \frac{1}{n}\frac{\sin\left\{\frac{(r-n+1).(2m\pi+a)}{n}\right\}}{\sin a},$$

and $\frac{N.\sin n\theta}{\sin\theta}$, or $\frac{N.\sin a}{\sin\theta} = \frac{1}{n}\cos(r-n+1)\theta - \frac{M.\cos\theta\sin a}{\sin\theta}$

$$= \left\{\frac{1}{n}.\cos(r-n+1)\theta.\sin\theta - \frac{1}{n}.\sin(r-n+1)\theta.\cos\theta\right\}\frac{1}{\sin\theta}$$

$$= -\frac{1}{n}.\frac{\sin(r-n)\theta}{\sin\theta} = \frac{1}{n}.\frac{\sin(n-r)\theta}{\sin\theta};$$

$$\therefore N = \frac{1}{n}\frac{\sin(n-r)\theta}{\sin n\theta} = \frac{1}{n}\frac{\sin\left\{\frac{(n-r)(2m\pi+a)}{n}\right\}}{\sin a},$$

and the integral is reduced to that of

$$\frac{1}{n\sin a}.\int_x \frac{x.\sin(r-n+1)\theta + \sin(n-r)\theta}{x^2 - 2x\cos\theta + 1},$$

the form of which is known.

EXAMPLES.

(1) $\int_x \frac{1}{x-x^2} = \log\left(\frac{x}{1-x}\right)$.

(2) $\int_x \frac{1}{x-x^3} = \log\frac{x}{\sqrt{1-x^2}}$.

(3) $\int_x \frac{x}{(x+a)(x+b)} = \frac{1}{b-a}\log\left\{\frac{(x+b)^b}{(x+a)^a}\right\}$.

(4) $\int_x \frac{x^2}{(x+a)(x+b)(x+c)} = \frac{a^2}{(b-a)(c-a)}\log(x+a)$

$$+ \frac{b^2}{(a-b)(c-b)}\log(x+b) + \frac{c^2}{(a-c)(b-c)}\log(x+c).$$

(5) $\int_x \frac{5x-2}{x^3+6x^2+8x} = \frac{1}{4}\log\frac{(x+2)^{12}}{x(x+4)^{11}}$.

(6) $\int_x \frac{3x+1}{x^3+2x^2+x} = \log\left(\frac{x}{x+1}\right) - \frac{2}{x+1}$.

(7) $\int_x \frac{1}{1+3x+2x^2} = \log\left(\frac{2x+1}{x+1}\right)$.

(8) $\int_x \frac{x}{(x+2)(x+3)^2} = -\frac{3}{x+3} + \log\left(\frac{x+3}{x+2}\right)^2$.

(9) $\int_x \frac{x}{(x-2)(x+3)^2} = -\frac{3}{5}\frac{1}{x+3} + \frac{2}{25}\log\left(\frac{x-2}{x+3}\right)$.

(10) $\int_x \frac{x^2}{(x+2)^2(x+4)^2} = -\frac{5x+12}{x^2+6x+8} + \log\left(\frac{x+4}{x+2}\right)^2$.

(11) $\int_x \frac{x^2+3x+1}{x^3+x^2-2x} = \frac{1}{3}\log\frac{(x-1)^5}{\sqrt{x^4+2x^3}}$.

(12) $\int_x \frac{2x}{(x^2+1).(x^2+3)} = \log\sqrt{\frac{x^2+1}{x^2+3}}$.

(13) $\int_x \frac{x^2}{(x^2+1).(x^2+4)} = \frac{1}{3}\left\{2\tan^{-1}\frac{x}{2} - \tan^{-1}x\right\}$.

(14) $\int_x \frac{x}{(x+1)(x+2)(x^2+1)} = \frac{2}{5}\log(x+2)$
$+\frac{1}{10}\left\{\log\sqrt{\frac{x^2+1}{x+1}} + 3\tan^{-1}x\right\}$.

(15) $\int_x \frac{1}{x^4+4x+3} = -\frac{1}{6}\frac{1}{x+1} + \frac{1}{9}\log\left(\frac{x+1}{\sqrt{x^2-2x+3}}\right)$
$+\frac{1}{18\sqrt{2}}\tan^{-1}\left(\frac{x-1}{\sqrt{2}}\right)$.

(16) $\int_x \frac{3x^2+x-2}{(x-1)^3.(x^2+1)} = -\frac{1}{2}\frac{1}{(x-1)^2} - \frac{5}{2}.\frac{1}{x-1}$
$+\frac{3}{2}\log\frac{\sqrt{x^2+1}}{x-1} - \tan^{-1}x$.

(17) $$\int_x \frac{1-x+x^2}{1+x+x^2+x^3} = \frac{1}{2}\left\{\log\frac{(1+x)^3}{\sqrt{1+x^2}} - \tan^{-1}x\right\}.$$

(18) $$\int_x \frac{1}{x\,(1+x)^2.(1+x+x^2)} = \frac{1}{1+x} + \log\frac{\sqrt{x^2+x^3+x^4}}{(1+x)^2} - \frac{1}{\sqrt{3}}\tan^{-1}\left(\frac{2x+1}{\sqrt{3}}\right).$$

(19) $$\int_x \frac{1}{x^8+x^7-x^4-x^3} = \frac{2-2x-5x^2}{4\,(1+x)\,x^2} + \log\sqrt[8]{\frac{(x-1)\,(x+1)^9}{(1+x^2)\,x^8}} - \frac{1}{4}\tan^{-1}x.$$

(20) $$\int_x \frac{x^4}{2+3x^2} = \frac{x^3}{9} - \frac{2x}{9} + \frac{2\sqrt{2}}{9\sqrt{3}}\tan^{-1}x\sqrt{\frac{3}{2}}.$$

(21) $$\int_x \frac{x}{a^4+x^4} = \frac{1}{2a^2}\tan^{-1}\left(\frac{x^2}{a^2}\right).$$

(22) $$\int_x \frac{x^5}{1+2x^2} = \frac{x^4}{8} - \frac{x^2}{8} + \frac{1}{8}\log\sqrt{1+2x^2}.$$

(23) $$\int_x \frac{x^4}{(1+x^2)^2} = \left(x^3+\frac{3x}{2}\right)\frac{1}{1+x^2} - \frac{3}{2}\tan^{-1}x.$$

(24) $$\int_x \frac{x^5}{(1+x^2)^3} = \left(x^2+\frac{3}{4}\right)\frac{1}{(1+x^2)^2} + \log\sqrt{1+x^2}.$$

(25) $$\int_x \frac{1}{x^6.(1+x^2)} = -\frac{1}{5x^5} + \frac{1}{3x^3} - \frac{1}{x} - \tan^{-1}x.$$

(26) $$\int_x \frac{1}{x^4(1+x^2)^2} = -\left(\frac{1}{3x^3} - \frac{5}{3x} - \frac{5x}{2}\right)\frac{1}{1+x^2} + \frac{5}{2}\tan^{-1}x.$$

(27) $$\int_x \frac{x^2}{a+bx+cx^2} = \frac{x}{c} - \frac{b}{c^2}.\log\sqrt{a+bx+cx^2} + \left(\frac{b^2}{2c^2} - \frac{a}{c}\right).\int_x \frac{1}{a+bx+cx^2}.$$

(28) $$\int_x \frac{x}{(1+x+x^2)^2} = -\frac{x+2}{3(1+x+x^2)} - \frac{2}{3\sqrt{3}}\tan^{-1}\left(\frac{2x+1}{\sqrt{3}}\right).$$

(29) $$\int_x \frac{x^5}{x^3+1} = \frac{x^3}{3} - \log\sqrt[3]{x^3+1}.$$

(30) $$\int_x \frac{x^4}{x^3+1} = \frac{x^2}{2} + \frac{1}{3}\log\frac{\sqrt{x^2-x+1}}{(x+1)} + \frac{1}{\sqrt{3}}\tan^{-1}\frac{x\sqrt{3}}{2-x}.$$

(31) $$\int_x \frac{x^2}{x^4-a^4} = \frac{1}{4a}\log\frac{x-a}{x+a} + \frac{1}{2}\tan^{-1}\frac{x}{a}.$$

(32) $$\int_x \frac{x^2}{x^4+1} = \frac{1}{4\sqrt{2}}\log\frac{x^2-x\sqrt{2+1}}{x^2+x\sqrt{2+1}} + \frac{1}{2\sqrt{2}}\tan^{-1}\frac{x\sqrt{2}}{1-x^2}.$$

(33) $$\int_x \frac{1}{x^6-1} = \frac{1}{6}\log\frac{(x-1)\sqrt{x^2-x+1}}{(x+1)\sqrt{x^2+x+1}} - \frac{1}{2\sqrt{3}}\tan^{-1}\frac{x\sqrt{3}}{1-x^2}.$$

(34) $$\int_x \frac{x^6}{1+x^3} \text{ being } = 0, \text{ when } x = 0, \text{ is when } x = 1$$

$$= \frac{1}{3}\log 2 - \frac{3}{4} + \frac{\pi}{3\sqrt{3}}.$$

(35) $$\int_x \frac{1}{x(1+x^3)} = \log\sqrt[3]{\frac{x^3}{1+x^3}}.$$

(36) $$\int_x \frac{1}{x(1+x^4)} = \log\sqrt{\frac{x^4}{1+x^4}}.$$

(37) $$\int_x \frac{1}{x^4(a+bx^3)} = -\frac{1}{3ax^3} + \frac{b}{3a^2}\log\left(\frac{a+bx^3}{x^3}\right).$$

(38) $$\int_x \frac{1}{x(1+x^3)^2} = \frac{1}{3(1+x^3)} - \log\sqrt[3]{\frac{1+x^3}{x^3}}.$$

CHAPTER III.

IRRATIONAL QUANTITIES.

---

29. The functions of this class will be treated of in the following order:

(1) Those which are the differential cofficients of known functions.

(2) Those which may be reduced to rational functions by means of obvious substitutions.

(3) Those which must be referred by means of *Formulas of Reduction* to known integrals.

30. Class I. Since if $u = \sin^{-1}x$, $\frac{du}{dx} = \frac{1}{\sqrt{1-x^2}}$,

and if $u = \cos^{-1}x$, $\frac{du}{dx} = -\frac{1}{\sqrt{1-x^2}}$,

..... $u = V\sin^{-1}x$, $\frac{du}{dx} = \frac{1}{\sqrt{2x-x^2}}$,

..... $u = \sec^{-1}x$, $\frac{du}{dx} = \frac{1}{x\sqrt{x^2-1}}$;

$$\therefore \int_x \frac{1}{\sqrt{1-x^2}} = \sin^{-1}x,$$

$$\int_x \frac{-1}{\sqrt{1-x^2}} = \cos^{-1}x,$$

$$\int_x \frac{1}{\sqrt{2x-x^2}} = V\sin^{-1}x,$$

$$\int_x \frac{1}{x\sqrt{x^2-1}} = \sec^{-1}x.$$

Cor. Since $\int_x \frac{1}{\sqrt{a^2-x^2}} = \int_x \frac{\frac{1}{a}}{\sqrt{1-\frac{x^2}{a^2}}} = \int_z \frac{1}{\sqrt{1-z^2}}$; $z = \frac{x}{a}$;

$$\therefore \int_x \frac{1}{\sqrt{a^2-x^2}} = \sin^{-1}(z) = \sin^{-1}\left(\frac{x}{a}\right),$$

and $\int_x \frac{1}{x\sqrt{x^2-a^2}} = \frac{1}{a}\cdot\int_x \frac{\frac{1}{a}}{\frac{x}{a}\sqrt{\frac{x^2}{a^2}-1}} = \frac{1}{a}\cdot\int_z \frac{1}{z\sqrt{z^2-1}}$

$$= \frac{1}{a}\sec^{-1}z = \frac{1}{a}\cdot\sec^{-1}\left(\frac{x}{a}\right).$$

31. Class II. Next, to integrate:

$$\frac{1}{\sqrt{x^2\pm a^2}},\quad \frac{1}{\sqrt{x^2\pm 2ax}},\quad \frac{1}{x\sqrt{x^2+a^2}},\quad \frac{1}{x\sqrt{a^2-x^2}}.$$

(1) If $\frac{du}{dx} = \frac{1}{\sqrt{x^2+a^2}}$,

Let $\sqrt{x^2+a^2} = xz$; $\quad\therefore\ x^2 = \frac{a^2}{z^2-1}$,

$$2\log x = \log a^2 - \log(z^2-1);$$

$$\therefore\ \frac{dx}{dz} = -\frac{xz}{z^2-1};$$

$$\therefore\ \frac{du}{dz} = \frac{du}{dx}\cdot\frac{dx}{dz} = \frac{1}{xz}\cdot\frac{dx}{dz} = -\frac{1}{z^2-1};$$

and $u = -\int_z \frac{1}{z^2-1} = \frac{1}{2}\cdot\int_z\left\{\frac{1}{z+1}-\frac{1}{z-1}\right\} = \frac{1}{2}\log\frac{z+1}{z-1}$

$$= \frac{1}{2}\cdot\log\frac{\sqrt{a^2+x^2}+x}{\sqrt{a^2+x^2}-x} = \frac{1}{2}\cdot\log\frac{(\sqrt{a^2+x^2}+x)^2}{a^2}$$

$$= \log\frac{\sqrt{a^2+x^2}+x}{a} = \log(x+\sqrt{x^2+a^2}) + C.$$

(2) Since $\frac{1}{\sqrt{x^2+2ax}} = \frac{1}{\sqrt{(x+a)^2-a^2}}$;

$$\therefore\ \int_x \frac{1}{\sqrt{x^2+2ax}} = \log(x+a+\sqrt{x^2+2ax}).$$

(3) If $\frac{du}{dx} = \frac{1}{x\sqrt{x^2+a^2}}$.

Let $\sqrt{x^2+a^2} = z$; $\quad\therefore\ x^2 = z^2-a^2$;

$$\therefore\ 2\log x = \log(x^2 - a^2);$$

$$\therefore\ \frac{dx}{dz} = \frac{xz}{x^2 - a^2};$$

$$\therefore\ \frac{du}{dz} = \frac{du}{dx}\cdot\frac{dx}{dz} = \frac{1}{x^2 - a^2};$$

$$\therefore\ u = \int_z \frac{1}{x^2 - a^2} = \frac{1}{2a}\cdot\int_z\left(\frac{1}{x-a} - \frac{1}{x+a}\right) = \frac{1}{2a}\cdot\log\frac{x-a}{x+a}$$

$$= \frac{1}{2a}\cdot\log\frac{\sqrt{x^2+a^2}-a}{\sqrt{x^2+a^2}+a} = \frac{1}{2a}\cdot\log\frac{x^2}{(\sqrt{x^2+a^2}+a)^2}$$

$$= \frac{1}{a}\cdot\log\left(\frac{x}{\sqrt{x^2+a^2}+a}\right).$$

(4) Hence, if $\frac{du}{dx} = \frac{1}{x\sqrt{a^2-x^2}}$, $u = \frac{1}{a}\cdot\log\left(\frac{x}{\sqrt{a^2-x^2}+a}\right)$.

32. Integrate $\frac{du}{dx} = \frac{1}{\sqrt{a+bx+cx^2}}$,

$$\frac{1}{\sqrt{a+bx+cx^2}} = \frac{1}{\sqrt{c}}\cdot\frac{1}{\sqrt{\frac{a}{c}+\frac{b}{c}x+x^2}}$$

$$= \frac{1}{\sqrt{c}}\cdot\frac{1}{\sqrt{\left(x+\frac{b}{2c}\right)^2+\frac{a}{c}-\frac{b^2}{4c^2}}}$$

$$= \frac{1}{\sqrt{c}}\cdot\frac{1}{\sqrt{\left(x+\frac{b}{2c}\right)^2+\frac{4ac-b^2}{4c^2}}},$$

which being of the form $\frac{1}{\sqrt{x^2 \pm a^2}}$;

$$\therefore\ u = \frac{1}{\sqrt{c}}\cdot\log\left\{\left(x+\frac{b}{2c}+\sqrt{\frac{a}{c}+\frac{bx}{c}+x^2}\right)\frac{2c}{\sqrt{4ac-b^2}}\right\}$$

$$= \frac{1}{\sqrt{c}}\cdot\log\left(\frac{2cx+b+2\sqrt{c}\sqrt{a+bx+cx^2}}{\sqrt{4ac-b^2}}\right).$$

33. Also, $\frac{1}{\sqrt{a+bx-cx^2}} = \frac{1}{\sqrt{c}}\cdot\frac{1}{\sqrt{\frac{a}{c}+\frac{bx}{c}-x^2}}$

$$= \frac{1}{\sqrt{c}} \cdot \frac{1}{\sqrt{\frac{a}{c} + \frac{b^2}{4c^2} - \left(x - \frac{b}{2c}\right)^2}} = \frac{1}{\sqrt{c}} \cdot \frac{1}{\sqrt{\frac{4ac+b^2}{4c^2} - \left(x - \frac{b}{2c}\right)^2}},$$

which is of the form $\frac{1}{\sqrt{a^2 - z^2}} = \frac{d \cdot \sin^{-1}\left(\frac{z}{a}\right)}{dz}$;

$$\therefore \int_x \frac{1}{\sqrt{a + bx - cx^2}} = \frac{1}{\sqrt{c}} \cdot \sin^{-1} \frac{x - \frac{b}{2c}}{\sqrt{\frac{4ac + b^2}{4c^2}}}$$

$$= \frac{1}{\sqrt{c}} \sin^{-1}\left(\frac{2cx - b}{\sqrt{4ac + b^2}}\right).$$

34. Integrate $\frac{du}{dx} = \frac{1}{x\sqrt{a + bx + cx^2}}$.

Let $x = \frac{1}{z}$; $\therefore \frac{dx}{dz} = -\frac{1}{z^2}$;

$$\therefore \frac{du}{dz} = \frac{du}{dx} \cdot \frac{dx}{dz} = \frac{1}{\frac{1}{z}\sqrt{a + \frac{b}{z} + \frac{c}{z^2}}} \times -\frac{1}{z^2} = \frac{-1}{\sqrt{az^2 + bz + c}};$$

or $u = -\int_z \frac{1}{\sqrt{az^2 + bz + c}}$. (Art. 31.)

35. Integrate $\frac{du}{dx} = \frac{1}{x^2\sqrt{a + bx + cx^2}}$.

Let $x = \frac{1}{z}$; $\therefore \frac{du}{dz} = \frac{du}{dx}\frac{dx}{dz} = \frac{z^2}{\sqrt{az^2 + bz + c}} \times -\frac{1}{z^2}$;

$$\therefore u = -\int_z \frac{z}{\sqrt{az^2 + bz + c}} = -\frac{1}{a} \cdot \int_z \frac{az + \frac{b}{2} - \frac{b}{2}}{\sqrt{az^2 + bz + c}}$$

$$= -\frac{1}{a}\int_z \left\{ \frac{az + \frac{b}{2}}{\sqrt{az^2 + bz + c}} - \frac{b}{2} \cdot \frac{1}{\sqrt{az^2 + bz + c}} \right\}$$

$$= -\frac{1}{a}\sqrt{az^2 + bz + c} + \frac{b}{2a} \cdot \int_z \frac{1}{\sqrt{az^2 + bz + c}};$$

the integral of which depends upon a preceding example.

36. Integrate $\frac{du}{dx} = \frac{1}{(a+bx)\sqrt{c+ex}}$;

Let $z = \sqrt{c+ex}$; $\therefore x = \frac{z^2-c}{e}$; $a+bx = \frac{ae-bc+bz^2}{e}$;

$$\therefore \frac{du}{dz} = \frac{e}{(ae-bc+bz^2)z} \cdot \frac{2z}{e} = \frac{2}{b\left(z^2 + \frac{ae-bc}{b}\right)}.$$

(1) Let $ae > bc$;

$$\therefore u = \frac{2}{b} \times \sqrt{\frac{b}{ae-bc}} \cdot \tan^{-1} \frac{z\sqrt{b}}{\sqrt{ae-bc}}$$

$$= \frac{2}{\sqrt{b}\sqrt{ae-bc}} \tan^{-1}\left(\frac{\sqrt{b}}{\sqrt{ae-bc}}\sqrt{c+ex}\right).$$

(2) Let $ae < bc$, and let $\frac{bc-ae}{b} = \beta^2$;

$$\therefore u = \frac{2}{b} \cdot \int_z \frac{1}{z^2-\beta^2} = \frac{1}{\beta b} \cdot \int_z \left\{\frac{1}{z-\beta} - \frac{1}{z+\beta}\right\}$$

$$= \frac{1}{\beta b} \log \frac{z-\beta}{z+\beta} = \frac{1}{\beta b} \log \frac{\sqrt{c+ex}-\beta}{\sqrt{c+ex}+\beta}.$$

37. Integrate $\frac{du}{dx} = \frac{1}{(a+bx)\sqrt{c+ex^2}}$.

Let $a+bx = \frac{1}{z}$; $x = \frac{1}{b}\left(\frac{1}{z} - a\right)$; $\frac{dx}{dz} = -\frac{1}{bz^2}$.

$$c + ex^2 = c + \frac{ea^2}{b^2} - \frac{2ae}{bz} + \frac{e}{b^2z^2}$$

$$= \frac{b^2c+ea^2}{b^2z^2} \cdot (\beta^2 - 2az + z^2) \text{ by substitution};$$

$$\therefore \frac{du}{dz} = \frac{bz^2}{\sqrt{b^2c+ea^2}\sqrt{\beta^2-2az+z^2}} \cdot \frac{-1}{bz^2}$$

$$= \frac{-1}{\sqrt{b^2c+ea^2}\sqrt{\beta^2-2az+z^2}};$$

$$\therefore u = \frac{-1}{\sqrt{cb^2+ea^2}} \cdot \int_z \frac{1}{\sqrt{\beta^2-2az+z^2}} \text{ (Art. 31).}$$

38. Integrate $\frac{du}{dx} = \frac{1}{(a+bx^2)\sqrt{c+ex^2}}$.

Let $\sqrt{c+ex^2} = xz$; $\therefore x^2 = \frac{c}{z^2-e}$;

$$\therefore \frac{dx}{dz} = -\frac{z}{x}\cdot\frac{c}{(z^2-e)^2},$$

$$a+bx^2 = \frac{az^2-ae+bc}{z^2-e};$$

$$\therefore \frac{du}{dz} = \frac{z^2-e}{dz^2-ae+bc}\cdot\frac{1}{xz}\cdot\frac{dx}{dz}$$

$$= -\frac{z^2-e}{az^2-ae+bc}\cdot\frac{1}{x^2}\cdot\frac{c}{(z^2-e)^2}$$

$$= -\frac{1}{az^2-ae+bc} = -\frac{1}{a}\cdot\frac{1}{z^2 \pm \beta^2},$$

which will be an angle or logarithm, according as the positive or negative sign is taken.

39. Integrate $\frac{du}{dx} = \frac{1}{(a+bx)\sqrt{cx^2+ex+f}}$.

Let $z = \frac{1}{a+bx}$; $\therefore x = \frac{1}{b}\left(\frac{1}{z}-a\right)$; $\frac{dx}{dz} = -\frac{1}{b}\frac{1}{z^2}$;

$$\therefore cx^2+ex+f = \frac{a^2}{b^2z^2}\cdot(z^2-2\beta z+\gamma);$$

$$\therefore \frac{du}{dz} = -\frac{1}{a\sqrt{z^2-2\beta z+\gamma}}. \quad \text{(Art. 37).}$$

40. Integrate $\frac{du}{dx} = X(a+bx)^{\frac{p}{q}}$, X being a rational function of x.

Let $a+bx = z^q$; $\therefore x = \frac{z^q-a}{b}$, and $\frac{dx}{dz} = \frac{q}{b}z^{q-1}$

$$\therefore \frac{du}{dz} = Z\cdot z^p\cdot\frac{q}{b}z^{q-1} = \frac{q}{b}Z\cdot z^{p+q-1};$$

where Z is the value of X, when $\frac{z^q-a}{b}$ is put for x.

41. Integrate $\frac{du}{dx} = X.(x + \sqrt{1+x^2})^q$, where X is either a rational function of x, or of x and $\sqrt{1+x^2}$.

Let $x + \sqrt{1+x^2} = z^q$; $\therefore\ 1 + x^2 = z^{2q} - 2xz^q + x^2$;

$\therefore\ x = \frac{1}{2}\{z^q - z^{-q}\}$; $\therefore\ \frac{dx}{dz} = \frac{q}{2}\{z^{q-1} + z^{-q-1}\} = \frac{q}{2}\frac{z^{2q}+1}{z^{q+1}}$,

and $\sqrt{1+x^2} = z^q + x = \frac{1}{2}(z^q + z^{-q})$;

$$\therefore\ \frac{du}{dz} = \int_z Z.z^q.\frac{q}{2}\frac{z^{2q}+1}{z^{q+1}} = \frac{q}{2}\int_z Z.\frac{z^{p+2q}+z^p}{z^{q+1}},$$

Z being the value of X, when $\frac{1}{2}(z^q - z^{-q})$ is put for x.

42. Integrate $\frac{du}{dx}$ when it is either

$$\frac{1}{(1-x^m)\sqrt[2m]{2x^m-1}} \text{ or } \frac{x^{m-1}}{(1-x^m)\sqrt[2m]{2x^m-1}}.$$

In the former, make $2x^m - 1 = z^{2m}x^{2m}$;

$$\therefore\ x^{2m} - 2x^m + 1 = x^{2m}(1 - z^{2m});$$

$$\text{or } \left(\frac{1-x^m}{x^m}\right)^2 = 1 - z^{2m} \ldots\ldots\ldots(1),$$

$$\text{or } \frac{1-x^m}{x^{2m+1}}.\frac{dx}{dz} = z^{2m-1} \ldots\ldots\ldots(2);$$

therefore by dividing (2) by (1),

$$\frac{1}{(1-x^m)x}\frac{dx}{dz} = \frac{z^{2m-2}}{1-z^{2m}};$$

$$\text{i. e. } \frac{du}{dx}.\frac{dx}{dz} \text{ or } \frac{du}{dz} = \frac{z^{2m-2}}{1-z^{2m}}.$$

In the latter, let $2x^m - 1 = z^{2m}$; $x^{m-1}\frac{dx}{dz} = z^{2m-1}$,

and $1 - x^m = 1 - \frac{1}{2}(z^{2m}+1) = \frac{1}{2}(1-z^{2m})$;

$$\therefore\ \frac{x^{m-1}}{(1-x^m)z}\frac{dx}{dz} = \frac{du}{dz} = \frac{2z^{2m-2}}{1-z^{2m}}.$$

These formulas were rationalized by Lexell.

BINOMIAL DIFFERENTIAL COEFFICIENTS.

43. To integrate $\frac{du}{dx} = x^{m-1}(a + bx^n)^{\frac{p}{q}}$.

It may be rationalized when $\frac{m}{n}$ or $\frac{m}{n} + \frac{p}{q}$ is an integer.

(1) Let $a + bx^n = z^q$; $\therefore x^n = \frac{z^q - a}{b}$, $x^m = \left(\frac{z^q - a}{b}\right)^{\frac{m}{n}}$;

$$\therefore x^{m-1}\frac{dx}{dz} = \frac{q}{nb} z^{q-1}\left(\frac{z^q - a}{b}\right)^{\frac{m}{n}-1};$$

$$\therefore \frac{du}{dz} = \frac{du}{dx}\cdot\frac{dx}{dz} = z^p . x^{m-1}\frac{dx}{dz} = \frac{q}{nb} z^{p+q-1}\left(\frac{z^q - a}{b}\right)^{\frac{m}{n}-1},$$

which is rational if $\frac{m}{n}$ be an integer, and easily integrable, by expanding the binomial.

(2) If $\frac{m}{n}$ be a fraction. Let $a + bx^n = x^n z^q$;

$$\therefore x^n = \frac{a}{z^q - b}, \quad x^m = \frac{a^{\frac{m}{n}}}{(z^q - b)^{\frac{m}{n}}};$$

$$\therefore x^{m-1}\frac{dx}{dz} = -\frac{q a^{\frac{m}{n}}}{n}\cdot\frac{z^{q-1}}{(z^q - b)^{\frac{m}{n}+1}}; \quad (a + bx^n)^{\frac{p}{q}} = \frac{a^{\frac{p}{q}} z^p}{(z^q - b)^{\frac{p}{q}}};$$

$$\therefore \frac{du}{dz} = \frac{a^{\frac{p}{q}} z^p}{(z^q - b)^{\frac{p}{q}}}\cdot x^{m-1}\cdot\frac{dx}{dz} = -\frac{q a^{\frac{m}{n}+\frac{p}{q}}}{n}\cdot\frac{z^{p+q-1}}{(z^q - b)^{\frac{m}{n}+\frac{p}{q}+1}},$$

which is rational when $\frac{m}{n} + \frac{p}{q}$ is an integer, and easily integrable if $\frac{m}{n} + \frac{p}{q}$ be a *negative* integer.

Cor. We have assumed that m and n are integers, but if they be fractions as $\frac{r}{s}$ and $\frac{s_1}{r_1}$. Make $v^{ss_1} = x$; $\therefore x^{\frac{r}{s}} = v^{rs_1}$, and $x^{\frac{r_1}{s_1}} = v^{r_1 s}$. Also n is assumed positive, for if not, let

$$x = \frac{1}{u}; \quad \therefore x^{-n} = u^n.$$

Ex. 1. Let $\frac{du}{dx} = x^3\sqrt{1+x^2}$.

Here $m-1=3$, and $n=2$; $\therefore \frac{m}{n} = \frac{4}{2} = 2$.

Let $1+x^2 = z^2$; $\therefore x^2 = z^2-1$, $x^4 = (z^2-1)^2$;

$$\therefore x^3\frac{dx}{dz} = (z^2-1)z;$$

$$\therefore \frac{du}{dz} = x^3 z\frac{dx}{dz} = z^2(z^2-1) = z^4 - z^2;$$

$$\therefore u = \frac{z^5}{5} - \frac{z^3}{3} + C = \frac{(1+x^2)^{\frac{5}{2}}}{5} - \frac{(x^2+1)^{\frac{3}{2}}}{3} + C.$$

Ex. 2. Let $\frac{du}{dx} = \frac{1}{x^4\sqrt{1+x^2}}$.

Here $\frac{m}{n} = -\frac{3}{2}$, and $\frac{p}{q} = -\frac{1}{2}$; $\therefore \frac{m}{n} + \frac{p}{q} = -2$.

And $\frac{1}{x^4\sqrt{1+x^2}} = \frac{1}{x^5\sqrt{x^{-2}+1}} = \frac{x^{-5}}{\sqrt{x^{-2}+1}}$.

Let $x^{-2}+1 = z^2$; $\therefore x^{-3} = -z.\frac{dz}{dx}$, $x^{-2} = z^2-1$;

$$\therefore \frac{du}{dz} = \frac{z^2-1}{z}x^{-3}\frac{dx}{dz} = -(z^2-1);$$

$$\therefore u = -\left(\frac{z^3}{3} - z\right) = -z.\left(\frac{z^2}{3} - 1\right)$$

$$= -\frac{\sqrt{1+x^2}}{x}\left\{\frac{1+x^2}{3x^2} - 1\right\} = \frac{(2x^2-1)\sqrt{1+x^2}}{3x^3}.$$

Ex. 3. Let $\frac{du}{dx} = \frac{1}{\sqrt[n]{1-x^n}} = (1-x^n)^{-\frac{1}{n}}$.

Here $\frac{m}{n} + \frac{p}{q} = \frac{1}{n} - \frac{1}{n} = 0$.

Let $\therefore 1-x^n = x^n z^n$; $\therefore x^n = \frac{1}{1+z^n}$;

$$\therefore n\log x = -\log(1+z^n); \quad \therefore \frac{1}{x}\frac{dx}{dz} = -\frac{z^{n-1}}{1+z^n};$$

$$\therefore \frac{du}{dz} = \frac{1}{xz} \cdot \frac{dx}{dz} = -\frac{x^{n-2}}{1+x^n},$$

which may be integrated by partial fractions.

44. This method of substitution is seldom adopted, the integration by parts being more generally useful, we shall henceforth confine our attention chiefly to it.

45. Let $\frac{du}{dx} = x^{m-1} \cdot (a+bx^n)^r$; r being a fraction and m and n positive integers.

Now we must diminish either m or r; first to diminish the index m;

$$x^{m-1}(a+bx^n)^r = x^{m-n} \cdot x^{n-1}(a+bx^n)^r;$$

$$\therefore p = x^{m-n}; \quad \frac{dq}{dx} = x^{n-1}(a+bx^n)^r; \quad \therefore q = \frac{(a+bx^n)^{r+1}}{nb(r+1)};$$

$$\therefore u = \frac{x^{m-n} \cdot (a+bx^n)^{r+1}}{nb(r+1)} - \frac{m-n}{nb(r+1)} \cdot \int_x x^{m-n-1}(a+bx^n)^{r+1}$$

$$= \frac{x^{m-n} \cdot (a+bx^n)^{r+1}}{nb(r+1)} - \frac{m-n}{nb(r+1)} \cdot \{a\int_x x^{m-n-1}(a+bx^n)^r + bu\};$$

$$\because (a+bx^n)^{r+1} = (a+bx^n)^r \cdot (a+bx^n);$$

$$\therefore \left\{1 + \frac{m-n}{n(r+1)}\right\} u = \frac{x^{m-n} \cdot (a+bx^n)^{r+1}}{nb(r+1)}$$

$$- \frac{(m-n) \cdot a}{nb(r+1)} \cdot \int_x x^{m-n-1}(a+bx^n)^r;$$

$$\therefore u = \frac{x^{m-n} \cdot (a+bx^n)^{r+1}}{b(nr+m)} - \frac{a(m-n)}{b \cdot (nr+m)} \int_x x^{m-n-1} \cdot (a+bx^n)^r \quad (A),$$

in which the integral depends upon one in which the index of x without the parenthesis is less by n than in the original function; after s reductions we shall come to $\int_x x^{m-sn-1} \cdot (a+bx^n)^r$, the coefficient of which is $-\frac{a(m-sn)}{b(rn+m)-(s-1)n}$; which vanishes when m is a multiple of n.

46. Next to diminish the index r;

$$\int_x x^{m-1}(a+bx^n)^r = a\int_x x^{m-1}(a+bx^n)^{r-1} + b\int_x x^{m+n-1}(a+bx^n)^{r-1};$$

but from (A), by putting $m+n$ for m, and $r-1$ for r,

$$\int x^{m+n-1}(a+bx^n)^{r-1} = \frac{x^m(a+bx^n)^r - am\int x^{m-1}(a+bx^n)^{r-1}}{b(rn+m)};$$

substituting this in the preceding equation, we have

$$u = \frac{x^m(a+bx^n)^r}{rn+m} + \frac{rna}{rn+m}\int x^{m-1}(a+bx^n)^{r-1} \quad \text{(B)}.$$

By this formula successive unities may be taken from r, and thus $\int x^{m-1}(a+bx^n)^r$ may be made to depend upon $\int x^{m-sn-1}(a+bx^n)^{r-t}$; sn being the greatest multiple of n contained in $m-1$, and t the greatest whole number in r.

47. When m and r are negative the formulas (A) and (B) will not apply, for the exponent of x, as also that of $a+bx^n$ would be increased by them. Suitable formulas may however be deduced from them. For from (A) we have

$$\int x^{m-n-1}(a+bx^n)^r$$
$$= \frac{x^{m-n}(a+bx^n)^{r+1} - b(m+nr)\int x^{m-1}(a+bx^n)^r}{a(m-n)}.$$

For m put $m+n$; $\therefore m-n-1 = m-1$;

$$\therefore u = \frac{x^m(a+bx^n)^{r+1} - b(m+n+nr)\int x^{m+n-1}(a+bx^n)^r}{ma} \quad \text{(C)};$$

a formula which decreases m, if it be negative.

Also from (B),

$$\int x^{m-1}(a+bx^n)^{r-1} = -\frac{x^m(a+bx^n)^r - (m+nr)\int x^{m-1}(a+bx^n)^r}{r.na};$$

$\therefore$ writing $r+1$ for r, we have

$$u = -\frac{x^m(a+bx^n)^{r+1} - (m+n+nr)\int x^{m-1}(a+bx^n)^{r+1}}{(r+1)na} \quad \text{(D)};$$

which diminishes (r), (r) being negative.

48. Ex. $\frac{du}{dx} = \frac{x^m}{\sqrt{1-x^2}}$; here formula (A) applies:

$$a=1,\ b=-1,\ r=-\tfrac{1}{2},\ n=2,\ m-1=m;$$

$$\therefore u = -\frac{x^{m-1}\sqrt{1-x^2}}{m} + \frac{m-1}{m}\cdot\int\frac{x^{m-2}}{\sqrt{1-x^2}}.$$

But in practice we proceed thus:

$$\because \frac{x^m}{\sqrt{1-x^2}} = x^{m-1}\frac{x}{\sqrt{1-x^2}}; \quad \therefore p = x^{m-1}, \frac{dq}{dx} = \frac{x}{\sqrt{1-x^2}};$$

$$\therefore \frac{dp}{dx} = (m-1)x^{m-2}; \quad q = -\sqrt{1-x^2};$$

$$\therefore u = -x^{m-1}\sqrt{1-x^2} + (m-1)\int x^{m-2}\sqrt{1-x^2}$$

$$= -x^{m-1}\sqrt{1-x^2} + (m-1)\int \frac{x^{m-2}}{\sqrt{1-x^2}} - (m-1)u^*;$$

$$\therefore m\int \frac{x^m}{\sqrt{1-x^2}} = -x^{m-1}\sqrt{1-x^2} + (m-1)\int \frac{x^{m-2}}{\sqrt{1-x^2}};$$

$$\therefore \int \frac{x^m}{\sqrt{1-x^2}} = -\frac{x^{m-1}\sqrt{1-x^2}}{m} + \frac{m-1}{m}\int \frac{x^{m-2}}{\sqrt{1-x^2}};$$

and by putting $m-2$, $m-4$, &c. for m, the integral may be reduced either to

$$\int \frac{x}{\sqrt{1-x^2}}, \text{ or } \int \frac{1}{\sqrt{1-x^2}};$$

or, to $-\sqrt{1-x^2}$, or $\sin^{-1}x$, according as m is odd or even.

Ex. Let $\int \frac{x^4}{\sqrt{1-x^2}}$ be required. Here $m = 4$.

$$\int \frac{x^4}{\sqrt{1-x^2}} = -\frac{x^3\sqrt{1-x^2}}{4} + \tfrac{3}{4}\int \frac{x^2}{\sqrt{1-x^2}};$$

$$\int \frac{x^2}{\sqrt{1-x^2}} = -\frac{x\sqrt{1-x^2}}{2} + \tfrac{1}{2}\int \frac{1}{\sqrt{1-x^2}}$$

$$= -\frac{x\sqrt{1-x^2}}{2} + \tfrac{1}{2}\sin^{-1}x + C;$$

$$\therefore \int \frac{x^4}{\sqrt{1-x^2}} = -\sqrt{1-x^2}\left\{\frac{x^3}{4} + \frac{3x}{2.4}\right\} + \frac{1.3}{2.4}\sin^{-1}x + C.$$

49. To find the general value of the integral.

(1) Let m be even $= 2n$,

* Since $x^{m-2}\sqrt{1-x^2} = \frac{x^{m-2}}{\sqrt{1-x^2}} - \frac{x^m}{\sqrt{1-x^2}}$.

let $P_{2n} = \int_s \frac{x^{2n}}{\sqrt{1-x^2}}$, and $Q_{2n-1} = x^{2n-1}\sqrt{1-x^2}$;

$$\therefore P_{2n} = -\frac{1}{2n} Q_{2n-1} + \frac{2n-1}{2n} P_{2n-2},$$

$$P_{2n-2} = -\frac{1}{2n-2} Q_{2n-3} + \frac{2n-3}{2n-2} P_{2n-4},$$

$$P_{2n-4} = -\frac{1}{2n-4} Q_{2n-5} + \frac{2n-5}{2n-4} P_{2n-6},$$

$$\vdots$$

$$P_2 = -\tfrac{1}{2} Q_1 + \tfrac{1}{2} P_0, \text{ where } P_0 = \sin^{-1} x;$$

$$\therefore P_{2n} = -\left\{\frac{Q_{2n-1}}{2n} + \frac{2n-1}{2n.(2n-2)} Q_{2n-3}\right.$$

$$\left. + \frac{(2n-1).(2n-3)}{2n.(2n-2)(2n-4)} Q_{2n-5} + \&c.\right\}$$

$$+ \frac{(2n-1).(2n-3)(2n-5).....3.1}{2n.(2n-2)(2n-4).....4.2} \sin^{-1} x + C.$$

If the integral $= 0$, when $x = 0$. Then $C = 0$, for Q_{2n-1}, Q_{2n-3}, &c. each $= 0$.

If $x = 1$, Q_{2n-1}, Q_{2n-3}, &c. each $= 0$, and $\sin^{-1} x = \frac{\pi}{2}$;

$$\therefore \int_s \frac{x^{2n}}{\sqrt{1-x^2}}, \begin{matrix}\text{from } x=0\\ \text{to } x=1\end{matrix} = \frac{(2n-1).(2n-3).(2n-5)...3.1}{2n.(2n-2).(2n-4)...4.2} \cdot \frac{\pi}{2}.$$

(2) Let m be odd and $= 2n + 1$;

$$\therefore P_{2n+1} = -\frac{1}{2n+1} Q_{2n} + \frac{2n}{2n+1} P_{2n-1},$$

$$P_{2n-1} = -\frac{1}{2n-1} Q_{2n-2} + \frac{2n-2}{2n-1} P_{2n-3},$$

$$\vdots$$

$$P_3 = -\tfrac{1}{3} Q_2 + \tfrac{2}{3} P_1,$$

and $P_1 = \int_s \frac{x}{\sqrt{1-x^2}} = -\sqrt{1-x^2}$;

$$\therefore P_{2n+1} = -\left\{\frac{1}{2n+1} Q_{2n} + \frac{2n}{(2n+1)(2n-1)} Q_{2n-2}\right.$$

$$\left. + \frac{2n.(2n-2)}{(2n+1)(2n-1)(2n-3)} Q_{2n-4} + \&c.\right\}$$

$$- \frac{2n.(2n-2)(2n-4)...4.2}{(2n+1)(2n-1)(2n-3)...5.3} \sqrt{1-x^2} + C.$$

If $P_{2n+1} = 0$ when $x = 0$, since then $Q_{2n} = 0$;

$$\therefore\ 0 = -\frac{2n\,.\,(2n-2)\,(2n-4)\ldots4\,.\,2}{(2n+1)\,(2n-1)\,(2n-3)\ldots5\,.\,3} + C;$$

whence by subtraction,

$$P_{2n+1} = \frac{2n\,.\,(2n-2)\,(2n-4)\ldots4\,.\,2}{(2n+1)\,(2n-1)\,(2n-3)\ldots5\,.\,3}$$

$$-\left\{\frac{1}{2n+1}Q_{2n} + \frac{2n}{(2n+1)\,(2n-1)}Q_{2n-2} + \&c.\right\}.$$

Let $x = 1$;

$$\therefore \int_x \frac{x^{2n+1}}{\sqrt{1-x^2}}, \quad \begin{matrix}\text{from } x=0\\ \text{to } x=1\end{matrix}\Big\} = \frac{2n\,.\,(2n-2)\ldots6\,.\,4\,.\,2}{(2n+1)\,(2n-1)\ldots7\,.\,5\,.\,3}.$$

Cor. If n be infinite, we may make $P_{2n} = P_{2n+1}$,

$$\text{or } \frac{\pi}{2}\,.\,\frac{1\,.\,3\,.\,5\,.\,7,\ \&c.}{2\,.\,4\,.\,6\,.\,8,\ \&c.} = \frac{2\,.\,4\,.\,6\,.\,8,\ \&c.}{3\,.\,5\,.\,7\,.\,9,\ \&c.},$$

$$\text{or } \frac{\pi}{2} = \frac{2\,.\,2\,.\,4\,.\,4\,.\,6\,.\,6\,.\,8\,.\,8,\ \&c.}{1\,.\,3\,.\,3\,.\,5\,.\,5\,.\,7\,.\,7\,.\,9,\ \&c.},$$

which is Wallis's Theorem for the length of the circle.

50. Let $\dfrac{du}{dx} = (a^2 - x^2)^{\frac{n}{2}}$.

$$(a^2 - x^2)^{\frac{n}{2}} = a^2\,.\,(a^2 - x^2)^{\frac{n-2}{2}} - x^2\,.\,(a^2 - x^2)^{\frac{n-2}{2}};$$

$$\therefore\ u = a^2\int_x (a^2 - x^2)^{\frac{n-2}{2}} - \int_x x^2\,.\,(a^2 - x^2)^{\frac{n-2}{2}}$$

$$= a^2\int_x (a^2 - x^2)^{\frac{n-2}{2}} + \frac{x\,(a^2 - x^2)^{\frac{n}{2}}}{n} - \frac{u}{n};$$

$$\therefore\ u = \frac{x\,(a^2 - x^2)^{\frac{n}{2}}}{n+1} + \frac{n\,a^2}{n+1}\,.\int_x (a^2 - x^2)^{\frac{n-2}{2}};$$

by which u is reduced, n being odd to $\int_x (a^2 - x^2)^{\frac{1}{2}}$.

$$\text{Also } \int_x (a^2 - x^2)^{\frac{1}{2}} = \int_x \frac{a^2}{\sqrt{a^2 - x^2}} - \int_x \frac{x^2}{\sqrt{a^2 - x^2}}$$

$$= \frac{x}{2}\sqrt{a^2 - x^2} + \frac{a^2}{2}\,.\,\sin^{-1}\,.\,\frac{x}{a}.$$

If the integral be required between $x = 0$ and $x = a$;

$$\int_x (a^2 - x^2)^{\frac{1}{2}} = \frac{\pi a^2}{4} = \frac{1}{2} \cdot \frac{\pi a^2}{2};$$

$$\therefore \int_x (a^2 - x^2)^{\frac{3}{2}} = \frac{3a^2}{4} \int (a^2 - x^2)^{\frac{1}{2}} = \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{\pi a^4}{2}$$

$$\int_x (a^2 - x^2)^{\frac{5}{2}} = \frac{5a^2}{6} \cdot \int (a^2 - x^2)^{\frac{3}{2}} = \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \cdot \frac{\pi a^6}{2};$$

and $$\int_x (a^2 - x^2)^{\frac{n}{2}} = \frac{1 \cdot 3 \cdot 5 \cdot 7 \ldots (n-2) \cdot n}{2 \cdot 4 \cdot 6 \cdot 8 \ldots (n-1)(n+1)} \cdot \frac{\pi a^{n+1}}{2}.$$

51. Integrate $\dfrac{du}{dx} = \dfrac{1}{x^m \sqrt{1 + x^2}}$.

$$\int_x \frac{1}{x^m \sqrt{1 + x^2}} = \int_x \frac{x}{x^{m+1} \sqrt{1 + x^2}}.$$

Here $p = \dfrac{1}{x^{m+1}}$, and $\dfrac{dq}{dx} = \dfrac{x}{\sqrt{1 + x^2}}$;

$$\therefore \frac{dp}{dx} = -\frac{m+1}{x^{m+2}}, \text{ and } q = \sqrt{1 + x^2};$$

$$\therefore \int_x \frac{1}{x^m \sqrt{1 + x^2}} = \frac{\sqrt{1 + x^2}}{x^{m+1}} + (m+1) \cdot \int_x \frac{\sqrt{1 + x^2}}{x^{m+2}}$$

$$= \frac{\sqrt{1 + x^2}}{x^{m+1}} + (m+1) \cdot \int_x \frac{1}{x^{m+2} \sqrt{1 + x^2}} + (m+1) \cdot \int_x \frac{1}{x^m \sqrt{1 + x^2}};$$

$$\therefore \int_x \frac{1}{x^{m+2} \sqrt{1 + x^2}} = -\frac{1}{m+1} \cdot \frac{\sqrt{1 + x^2}}{x^{m+1}} - \frac{m}{m+1} \cdot \int_x \frac{1}{x^m \sqrt{1 + x^2}}.$$

For $m + 2$ put m;

$$\therefore \int_x \frac{1}{x^m \sqrt{1 + x^2}} = -\frac{1}{m-1} \cdot \frac{\sqrt{1 + x^2}}{x^{m-1}} - \frac{m-2}{m-1} \cdot \int_x \frac{1}{x^{m-2} \sqrt{1 + x^2}};$$

and therefore the integral may be reduced either to

$\int_x \dfrac{1}{x \sqrt{1 + x^2}}$, or $\int_x \dfrac{1}{x^2 \sqrt{1 + x^2}}$, according as m is odd or even,

also $$\int_x \frac{1}{x \sqrt{1 + x^2}} = \log \frac{1}{1 + \sqrt{1 + x^2}},$$

and $\int_x \frac{1}{x^2\sqrt{1+x^2}} = \int_x \frac{1}{x^3\sqrt{x^{-2}+1}} = -\sqrt{x^{-2}+1} = -\frac{\sqrt{1+x^2}}{x}$.

52. Integrate $\frac{du}{dx} = \frac{1}{x^m\sqrt{x^2-1}}$.

$$\int_x \frac{1}{x^m\sqrt{x^2-1}} = \int_x \frac{x}{x^{m+1}\sqrt{x^2-1}} = \int_x \frac{1}{x^{m+1}} \cdot \frac{d\cdot\sqrt{x^2-1}}{dx}$$

$$= \frac{\sqrt{x^2-1}}{x^{m+1}} + (m+1)\cdot\int_x \frac{\sqrt{x^2-1}}{x^{m+2}}$$

$$= \frac{\sqrt{x^2-1}}{x^{m+1}} + (m+1)\cdot\int_x \frac{1}{x^m\sqrt{x^2-1}} - (m+1)\cdot\int_x \frac{1}{x^{m+2}\sqrt{x^2-1}};$$

$$\therefore \int_x \frac{1}{x^{m+2}\sqrt{x^2-1}} = \frac{1}{m+1}\cdot\frac{\sqrt{x^2-1}}{x^{m+1}} + \frac{m}{m+1}\cdot\int_x \frac{1}{x^m\sqrt{x^2-1}};$$

therefore, writing m for $(m+2)$,

$$\int_x \frac{1}{x^m\sqrt{x^2-1}} = \frac{1}{m-1}\cdot\frac{\sqrt{x^2-1}}{x^{m-1}} + \frac{m-2}{m-1}\cdot\int_x \frac{1}{x^{m-2}\sqrt{x^2-1}},$$

and therefore u may be reduced, m odd, to

$\int \frac{1}{x\sqrt{x^2-1}} = \sec^{-1}x$, and m even, to $\int_x \frac{1}{x^2\sqrt{x^2-1}} = \frac{\sqrt{x^2-1}}{x}$.

EXAMPLE. Find $\int_x \frac{1}{x^5\sqrt{x^2-1}}$.

$$\int_x \frac{1}{x^5\sqrt{x^2-1}} = \tfrac{1}{4}\cdot\frac{\sqrt{x^2-1}}{x^4} + \tfrac{3}{4}\cdot\int_x \frac{1}{x^3\sqrt{x^2-1}},$$

$$\int_x \frac{1}{x^3\sqrt{x^2-1}} = \tfrac{1}{2}\cdot\frac{\sqrt{x^2-1}}{x^2} + \tfrac{1}{2}\cdot\sec^{-1}x;$$

$$\therefore \int_x \frac{1}{x^5\sqrt{x^2-1}} = \tfrac{1}{4}\cdot\frac{\sqrt{x^2-1}}{x^4} + \frac{3}{2.4}\cdot\frac{\sqrt{x^2-1}}{x^2} + \frac{1.3}{2.4}\sec^{-1}x.$$

53. Integrate $\frac{du}{dx} = \frac{x^m}{\sqrt{2ax-x^2}}$.

$$\int_x \frac{x^m}{\sqrt{2ax-x^2}} = \int_x \frac{-x^{m-1}.(a-x)+ax^{m-1}}{\sqrt{2ax-x^2}}$$

$$= -\int_x \frac{x^{m-1}(a-x)}{\sqrt{2ax-x^2}} + a.\int_x \frac{x^{m-1}}{\sqrt{2ax-x^2}}.$$

Now $\int_x x^{m-1}\frac{a-x}{\sqrt{2ax-x^2}} = x^{m-1}\sqrt{2ax-x^2}-(m-1).\int_x \sqrt{2ax-x^2}.x^{m-2}$

$$= x^{m-1}\sqrt{2ax-x^2}-2.(m-1)a.\int_x \frac{x^{m-1}}{\sqrt{2ax-x^2}}+(m-1).\int_x \frac{x^m}{\sqrt{2ax-x^2}};$$

therefore, substituting

$$m\int_x \frac{x^m}{\sqrt{2ax-x^2}} = -x^{m-1}\sqrt{2ax-x^2}+(2m-1)a.\int_x \frac{x^{m-1}}{\sqrt{2ax-x^2}};$$

$$\therefore \int_x \frac{x^m}{\sqrt{2ax-x^2}} = -\frac{x^{m-1}\sqrt{2ax-x^2}}{m}+\frac{2m-1}{m}.a.\int_x \frac{x^{m-1}}{\sqrt{2ax-x^2}},$$

by which u may be reduced to $\int_x \frac{1}{\sqrt{2ax-x^2}} = V\sin^{-1}\frac{x}{a}$.

The last term $= a^m.\frac{(2m-1)(2m-3)(2m-5)\ldots3.1}{m.(m-1).(m-2)\ldots2.1}V\sin^{-1}\frac{x}{a}$.

Cor. If $u=0$, when $x=0$, and its value be required when $x=2a$. Then, since all the terms vanish when $x=0$; $\therefore C=0$: and when $x=2a$, all the terms of the form $x^{m-1}\sqrt{2ax-x^2}$ vanish; but $V\sin^{-1}\frac{x}{2a}=\pi$;

$$\therefore \int_x \frac{x^m}{\sqrt{2ax-x^2}} \text{ from } x=0 \text{ to } x=2a,$$

$$= \pi.a^m.\frac{1.3.5\ldots(2m-3).(2m-1)}{1.2.3\ldots(m-1)\quad m}$$

54. Integrate $\frac{du}{dx} = \frac{1}{x^m\sqrt{2ax-x^2}}$. Let $x=\frac{1}{z}$;

$$\therefore \frac{du}{dz} = \frac{z^{m+1}}{\sqrt{2az-1}}\times -\frac{1}{z^2} = \frac{-z^{m-1}}{\sqrt{2az-1}}$$

$$= -\sqrt{\beta}.\frac{z^{m-1}}{\sqrt{z-\beta}}; \text{ if } \beta=\frac{1}{2a},$$

which may be integrated by parts, or rationalized by putting $\sqrt{z-\beta}=v$.

55. Integrate $\frac{du}{dx} = \frac{x^m}{\sqrt{a+bx+cx^2}}$.

$$\int_x \frac{x^m}{\sqrt{a+bx+cx^2}} = \int_x \frac{x^m}{\sqrt{c}\,\sqrt{\frac{a}{c}+\frac{b}{c}x+x^2}}.$$

Let $x+\frac{b}{2c}=z$; $\therefore\ x^2+\frac{b}{c}x+\frac{a}{c}=z^2+\frac{a}{c}-\frac{b^2}{4c^2}=z^2\pm\beta^2$;

$$\int_x \frac{x^m}{\sqrt{a+bx+cx^2}} = \frac{1}{\sqrt{c}}\int_z \frac{(z-a)^m}{\sqrt{z^2\pm\beta^2}};\quad \text{if } \frac{b}{2c}=a,$$

which may be made to depend upon $\int_z \frac{z^m}{\sqrt{z^2\pm\beta^2}}$.

56. Integrate $\frac{du}{dx} = \frac{1}{x^m\sqrt{a+bx+cx^2}}$.

Let $x=\frac{1}{z}$; $\therefore\ \frac{du}{dz} = -\frac{1}{z^2}\cdot\frac{du}{dx} = -\frac{z^{m+1}}{z^2\sqrt{az^2+bz+c}}$;

$$\therefore\ u = C - \int_z \frac{z^{m-1}}{\sqrt{az^2+bz+c}};$$

which may be integrated by the preceding method.

57. Integrate $\frac{du}{dx} = \frac{1}{\sqrt{c-x}\sqrt{2ax-x^2}}$.

$$\frac{1}{\sqrt{c-x}\sqrt{2ax-x^2}} = \frac{1}{\sqrt{c}}\frac{\left(1-\frac{x}{c}\right)^{-\frac{1}{2}}}{\sqrt{2ax-x^2}}$$

$$= \frac{1}{\sqrt{c}}\cdot\frac{1}{\sqrt{2ax-x^2}}\cdot\left\{1+\frac{1}{2}\cdot\frac{x}{c}+\frac{1\cdot3}{2\cdot4}\cdot\frac{x^2}{c^2}+\&c.\right\}$$

and thus u depends upon $\int_x \frac{x^m}{\sqrt{2ax-x^2}}$; this integral is met with in mechanics.

58. Lastly to prove Bernouilli's series for $\int_x u$.

Since $\int_x ux = \int_x x\frac{du}{dx}$,

$$\text{and } \int_x x\frac{du}{dx} = \frac{x^2}{2}\cdot\frac{du}{dx} - \frac{1}{2}\cdot\int_x x^2\frac{d^2u}{dx^2},$$

$$\int_x x^2 \frac{d^2u}{dx^2} = \frac{x^3}{3}\cdot\frac{d^2u}{dx^2} - \tfrac{1}{3}\cdot\int_x x^3\frac{d^3u}{dx^3},$$

$$\int_x x^3 \frac{d^3u}{dx^3} = \frac{x^4}{4}\cdot\frac{d^3u}{dx^3} - \tfrac{1}{4}\cdot\int_x x^4\frac{d^4u}{dx^4};$$

$$\&c.... = \&c.$$

$$\therefore \int_x u = ux - \frac{x^2}{1.2}\cdot\frac{du}{dx} + \frac{x^3}{2.3}\cdot\frac{d^2u}{dx^2} - \frac{x^4}{2.3.4}\cdot\frac{d^3u}{dx^3} + \&c. \pm \int_x \frac{d^nu}{dx^n}\cdot\frac{x^n}{1.2...n}.$$

Ex. Let $u = ax^3 + bx^2 + cx + e$;

$$\therefore \frac{du}{dx} = 3ax^2 + 2bx + c,$$

$$\frac{d^2u}{dx^2} = 6ax + 2b,$$

$$\frac{d^3u}{dx^3} = 6a, \text{ and } \frac{d^4u}{dx^4} = 0;$$

$$\therefore \int_x u = ax^4 + bx^3 + cx^2 + ex - \frac{3ax^4 + 2bx^3 + cx^2}{2} + ax^4 + \frac{bx^3}{3} - \frac{ax^4}{4}$$

$$= \frac{ax^4}{4} + \frac{bx^3}{3} + \frac{cx^2}{2} + ex.$$

EXAMPLES.

(1) $$\int_x x^2.\sqrt{a+bx} = \left\{\frac{(a+bx)^2}{7} - \frac{2a(a+bx)}{5} + \frac{a^2}{3}\right\}\frac{2(a+bx)^{\frac{3}{2}}}{b^3}.$$

(2) $$\int_x \frac{\sqrt{a+bx}}{x^2} = -\frac{\sqrt{a+bx}}{x} + \frac{b}{2\sqrt{a}}\log\left\{\frac{\sqrt{a+bx}-\sqrt{a}}{\sqrt{a+bx}+\sqrt{a}}\right\}.$$

(3) $$\int_x \frac{x^3}{\sqrt{a+bx}} = \left\{\frac{(a+bx)^3}{7} - \frac{3}{5}a(a+bx)^2 + a^2.(a+bx) - a^3\right\}\frac{2\sqrt{a+bx}}{b^4}.$$

(4) $$\int_x \frac{1}{x\sqrt{a+bx}} = \frac{1}{\sqrt{a}}\log\frac{\sqrt{a+bx}-\sqrt{a}}{\sqrt{a+bx}+\sqrt{a}}.$$

(5) $$\int_x \frac{1}{x\sqrt{bx-a}} = \frac{2}{\sqrt{a}}\tan^{-1}\sqrt{\frac{bx-a}{a}} = \frac{2}{\sqrt{a}}.\sin^{-1}\sqrt{\frac{bx-a}{bx}}.$$

(6) $\int_x \frac{1}{x^2\sqrt{4+3x}} = -\frac{\sqrt{4+3x}}{4x} - \frac{3}{16}\log\frac{\sqrt{4+3x}-2}{\sqrt{4+3x}+2}.$

(7) $\int_x \frac{x^2}{(1+x)^{\frac{1}{2}}} = \left\{x^2 - 4x - 8\right\}\frac{2}{3\sqrt{1+x}}.$

(8) $\int_x \frac{1}{x(a+bx)^{\frac{1}{2}}} = \frac{2}{a(a+bx)^{\frac{1}{2}}} + \frac{1}{a\sqrt{a}}.\log\left(\frac{\sqrt{a+bx}-\sqrt{a}}{\sqrt{a+bx}+\sqrt{a}}\right).$

(9) $\int_x \frac{x^2}{(1+x)^{\frac{5}{2}}} = (x^2 - 6x - 9)\frac{3}{4(1+x)^{\frac{3}{2}}}.$

(10) $\int_x \frac{1}{x(1+2x)^{\frac{5}{2}}} = \frac{2+3x}{3}\frac{4}{(1+2x)^{\frac{3}{2}}} + \log\left(\frac{\sqrt{1+2x}-1}{\sqrt{1+2x}+1}\right).$

(11) $\int_x \frac{x^3}{(2+x)^{\frac{9}{2}}} = -\left(x^3 + 4x^2 + \frac{32x}{5} + \frac{128}{35}\right)\frac{2}{(2+x)^{\frac{7}{2}}}.$

(12) $\int_x \frac{x^2}{\sqrt[3]{2+3x}} = (2+3x)^{\frac{2}{3}}.\left\{\frac{4-4x+5x^2}{40}\right\}.$

(13) $\int_x \frac{x^3}{(1+x)^{\frac{3}{2}}} = (14x^3 - 18x^2 + 27x - 81)\frac{3\sqrt[3]{1+x}}{4.5.7}.$

(14) $\int_x x^5\sqrt{1+x^2} = \left(x^4 - \frac{4x^2}{5} + \frac{8}{15}\right)\frac{(1+x^2)^{\frac{3}{2}}}{7}.$

(15) $\int_x \frac{\sqrt{1+x^2}}{x^6} = -\left\{\frac{1}{x^5} - \frac{2}{3x^3}\right\}\frac{(1+x^2)^{\frac{3}{2}}}{5}.$

(16) $\int_x x^3(1+x^2)^{\frac{3}{2}} = \frac{5x^2-2}{35}(1+x^2)^{\frac{5}{2}}.$

(17) $\int_x x^3(1+x^2)^{\frac{5}{2}} = \frac{7x^2-2}{63}(1+x^2)^{\frac{7}{2}}.$

(18) $\int_x \frac{x^5}{\sqrt{1-x^2}} = -\left\{\frac{x^4}{5} + \frac{4x^2}{15} + \frac{8}{15}\right\}\sqrt{1-x^2}.$

(19) $\int_x \frac{x^6}{\sqrt{1-x^2}} = -\left\{\frac{x^5}{6} + \frac{5x^3}{24} + \frac{5x}{16}\right\}\sqrt{1-x^2} + \frac{5}{16}\sin^{-1}x.$

(20) $\int_x \frac{1}{x^6\sqrt{1+x^2}} = -\left\{\frac{1}{5x^5} - \frac{4}{15x^3} + \frac{8}{15x}\right\}\sqrt{1+x^2}.$

(21) $\int_x \frac{1}{(a+bx^2)^{\frac{5}{2}}} = \left\{\frac{1}{3a(a+bx^2)} + \frac{2}{3a^2}\right\}\frac{x}{\sqrt{a+bx^2}}.$

(22) $\int_x \frac{x^3}{(1+x^2)^{\frac{3}{2}}} = (x^2+2)\frac{1}{\sqrt{1+x^2}}.$

(23) $\int_x \frac{x^4}{(1+x^2)^{\frac{3}{2}}} = \frac{x^3+3x}{2}\frac{1}{\sqrt{1+x^2}} - \frac{3}{2}\log(x+\sqrt{1+x^2}).$

(24) $\int_x \frac{x^3}{(1+x^2)^{\frac{5}{2}}} = -\left(x^2+\frac{2}{3}\right)\frac{1}{(1+x^2)^{\frac{3}{2}}}.$

(25) $\int_x \frac{x^5}{(1+x^2)^{\frac{5}{2}}} = \left(x^4+4x^2+\frac{8}{3}\right)\frac{1}{(1+x^2)^{\frac{5}{2}}}.$

(26) $\int_x \frac{x^2}{(1+x^2)^{\frac{9}{2}}} = \left(\frac{8x^7}{105}+\frac{4x^5}{15}+\frac{x^3}{3}\right)\frac{1}{(1+x^2)^{\frac{7}{2}}}.$

(27) $\int_x \frac{1}{(1-x)\sqrt{1-2x}} = \tan^{-1}.\left(\frac{\sqrt{1-2x}}{x}\right).$

(28) $\int_x \frac{1}{\sqrt{1+x+x^2}} = \log(2x+1+2\sqrt{1+x+x^2}).$

(29) $\int_x \frac{1}{\sqrt{1+x-x^2}} = \sin^{-1}\left(\frac{2x-1}{\sqrt{5}}\right).$

(30) $\int_x \frac{1}{(1+x)\sqrt{1-x}} = \frac{1}{\sqrt{2}}.\log\left(\frac{x-3+2\sqrt{2}\sqrt{1-x}}{1+x}\right).$

(31) $\int_x \frac{1}{x\sqrt{1+x+x^2}} = \log\left(\frac{2+x-2\sqrt{1+x+x^2}}{x}\right).$

(32) $\int_x \frac{1}{x\sqrt{x^2+x-1}} = \sin^{-1}.\left(\frac{x-2}{\sqrt{5}}\right).$

(33) $\int_x \frac{1}{(1+x)\sqrt{1+x^2}} = \frac{1}{\sqrt{2}}.\log\left(\frac{1-x-\sqrt{2}\sqrt{1+x^2}}{1+x}\right).$

(34) $\int_x \frac{1}{(1+x^2)\sqrt{1-x^2}} = \cos^{-1}.\sqrt{\frac{1-x^2}{1+x^2}}.$

(35) $\int_x \frac{1}{(1-x^2)\sqrt{1+x^2}} = \frac{1}{\sqrt{2}}\log\left(\frac{\sqrt{1+x^2}+x\sqrt{2}}{\sqrt{1-x^2}}\right).$

(36) $\int_x \frac{1}{(1+x)\sqrt{1+x-x^2}} = \tan^{-1}.\left(\frac{1+3x}{2\sqrt{1+x-x^2}}\right).$

(37) $$\int_x \frac{1}{(1+x)\sqrt{1-x-x^2}} = \log\left(\frac{x+3-2\sqrt{1-x-x^2}}{1+x}\right).$$

(38) $$\int_x \frac{1}{\sqrt{1+2x-x^2}} = \cos^{-1}.\left(\frac{1-x}{\sqrt{2}}\right).$$

(39) $$\int_x \frac{x}{\sqrt{a^4-x^4}} = \frac{1}{2}.\sin^{-1}.\left(\frac{x^2}{a^2}\right).$$

(40) $$\int_x \frac{x}{\sqrt{(x^2-a^2)(b^2-x^2)}} = \sin^{-1}.\sqrt{\frac{x^2-a^2}{b^2-a^2}}.$$

(41) $$\int_x \frac{1}{(a+bx^2)^{\frac{3}{2}}} = \frac{x}{a\sqrt{a+bx^2}}.$$

(42) $$\int_x \sqrt{\frac{1+x}{1-x}} = \sin^{-1}x - \sqrt{1-x^2}.$$

(43) $$\int_x \frac{1}{x}\sqrt{\frac{1+x}{1-x}} = \sin^{-1}x + \log\left(\frac{x}{1+\sqrt{1-x^2}}\right).$$

(44) $$\int_x x\sqrt{\frac{1+x}{1-x}} = \frac{1}{2}\sin^{-1}x - \frac{1}{2}(2+x)\sqrt{1-x^2}.$$

(45) $$\int_x \frac{\sqrt{x}}{\sqrt{a^3-x^3}} = \frac{2}{3}.\tan^{-1}.\sqrt{\frac{a^3-x^3}{x^3}}.$$

(46) $$\int_x \frac{1}{(2ax+x^2)^{\frac{3}{2}}} = -\frac{x+a}{a^2\sqrt{2ax+x^2}}.$$

(47) $$\int_x \frac{1}{(2ax+x^2)^{\frac{5}{2}}} = -\left\{\frac{1}{3(2ax+x^2)} - \frac{2}{3a^2}\right\}\frac{x+a}{a^2\sqrt{2ax+x^2}}.$$

(48) $$\int_x \frac{1}{(1+x+x^2)^{\frac{3}{2}}} = \frac{2.(2x+1)}{3\sqrt{1+x+x^2}}.$$

(49) $$\int_x \frac{1}{(1+x+x^2)^{\frac{5}{2}}} = \left\{\frac{1}{1+x+x^2} + \frac{8}{3}\right\}\frac{2(2x+1)}{9\sqrt{1+x+x^2}}.$$

(50) $$\int_x \frac{1}{(1+x)^2\sqrt{x}} = \frac{\sqrt{x}}{(1+x)} + \tan^{-1}(\sqrt{x}).$$

(51) $$\int_x \frac{x\sqrt{x}}{1+x} = \left(\frac{x}{3}-1\right)2\sqrt{x} + 2\tan^{-1}\sqrt{x}.$$

(52) $$\int_x \frac{x\sqrt{x}}{1+x^2} = 2\sqrt{x} + \frac{1}{\sqrt{2}}\left\{\log\left(\frac{x+1+\sqrt{2x}}{\sqrt{1+x^2}}\right) - \tan^{-1}\frac{\sqrt{2x}}{1-x}\right\}.$$

(53) $$\int_x \frac{1}{x^3\sqrt{2ax-x^2}} = -\frac{\sqrt{2ax-x^2}}{3a^2x^2}(a+x).$$

(54) $$\int_x \frac{x^2}{\sqrt{a+bx+cx^2}} = \sqrt{a+bx+cx^2}\left\{\frac{2cx-3b}{4c^2}\right\}$$
$$+\frac{4ac-3b^2}{8c^2\sqrt{c}}\log\left\{\frac{2cx+b+2\sqrt{c}\sqrt{a+bx+cx^2}}{2c}\right\}.$$

(55) Rationalize the integrals

(1) $$\int_x \frac{x^3+2x^{\frac{3}{2}}+x^{\frac{1}{2}}}{x+3x^{\frac{1}{4}}}, \qquad (2)\ \int_x \frac{x^4}{x^5+\sqrt[4]{(1+x)^3}},$$

$$(3)\ \int_x x^3\left(\frac{a+x}{a+x}\right)^{\frac{2}{3}}.$$

in (1) make $x = z^{12}$, and in (2) make $(1+x) = z^4$.

(56) $$\int_x x^3\sqrt{\frac{1+x^2}{1-x^2}} = \frac{1}{2}\tan^{-1}\sqrt[3]{\frac{1+x^2}{1-x^2}}$$
$$-\frac{1}{4}(2+x^2)\sqrt{1-x^4}.$$

(57) $$\int_x \frac{1}{x}\sqrt{\frac{a^2-c^2x^2}{a^2-x^2}} = \log\sqrt{\left(\frac{y-1}{y+1}\right)\left(\frac{y+c}{y-c}\right)^c}$$
$$\text{where } y = \sqrt{\frac{a^2-c^2x^2}{a^2-x^2}}.$$

(58) $$\int_x \frac{x^2+1}{x^2-1}\,\frac{1}{\sqrt{1-ax^2+x^4}} = \frac{1}{\sqrt{a-2}}\cos^{-1}\cdot\left(\frac{x\sqrt{a-2}}{x^2-1}\right)$$
$$\text{make } x-\frac{1}{x} = \frac{1}{z}.$$

CHAPTER IV.

INTEGRALS OF LOGARITHMIC AND EXPONENTIAL FUNCTIONS.

59. These functions are of the form $X\ (\log x)^n$, $X.a^x$, where X is a function of x.

60. Integrate $\int_x X.(\log x)^n$.

$$\text{Let } \int_x X = P, \quad \int_x P.\frac{1}{x} = Q, \quad \text{and } \int_x Q.\frac{1}{x} = R.$$

$$\text{Then } \int_x X(\log x)^n = P(\log x)^n - n.\int_x P.(\log x)^{n-1}.\frac{1}{x},$$

$$\text{and } \int_x \frac{P}{x}.(\log x)^{n-1} = Q(\log x)^{n-1} - (n-1).\int_x Q.(\log x)^{n-2}.\frac{1}{x},$$

$$\int_x \frac{Q}{x}.(\log x)^{n-2} = R(\log x)^{n-2} - (n-2).\int_x R.(\log x)^{n-3}.\frac{1}{x};$$

$$\therefore \int_x X(\log x)^n = P(\log x)^n - n.Q(\log x)^{n-1} + n.(n-1).R.(\log x)^{n-2} - \&c.$$

Ex. 1. $\int_x x^m(\log x)^n$.

$$\int_x x^m(\log x)^n = \frac{x^{m+1}(\log x)^n}{m+1} - \frac{n}{m+1}.\int_x x^{m+1}.(\log x)^{n-1}.\frac{1}{x}$$

$$= \frac{x^{m+1}(\log x)^n}{m+1} - \frac{n}{m+1}.\int_x x^m.(\log x)^{n-1},$$

$$\int_x x^m(\log x)^{n-1} = \frac{x^{m+1}(\log x)^{n-1}}{m+1} - \frac{n-1}{m+1}.\int_x x^m.(\log x)^{n-2},$$

and in this manner may the integral be reduced to

$$\int_x x^m = \frac{x^{m+1}}{m+1}, \text{ if } n \text{ be a whole number,}$$

$$\text{and } \therefore \int_x x^m.(\log x)^n = \frac{x^{m+1}}{m+1}\left\{(\log x)^n - \frac{n}{m+1}(\log x)^{n-1} + \frac{n.(n-1)}{(m+1)^2}(\log x)^{n-2} - \&c.\right\} \pm \frac{n.(n-1).(n-2)\ldots 2.1}{(m+1)^{m+1}}x^{m+1}.$$

Every term of the integral vanishes both when $x = 0$ and $x = 1$, except the last, which vanishes only when $x = 0$;

$$\therefore \int_x x^m (\log x)^n, \quad \left.\begin{matrix}\text{from } x=0\\ \text{to } x=1\end{matrix}\right\} = \pm \frac{1.2.3\ldots(n-1).n}{(m+1)^{n+1}}.$$

61. Integrate $\int_x \frac{X}{(\log x)^n}$, n a whole number.

$$\int_x \frac{X}{(\log x)^n} = \int_x \frac{X.x.\frac{d.\log x}{dx}}{(\log x)^n}. \quad \text{Since } x\frac{d.\log x}{dx} = x\frac{1}{x} = 1$$

$$= \frac{-X.x}{(n-1)(\log x)^{n-1}} + \frac{1}{n-1}.\int_x \frac{d.(Xx)}{dx} \Big/ (\log x)^{n-1}.$$

Let $\frac{d.(Xx)}{dx} = P$;

$$\therefore \int_x \frac{X}{(\log x)^n} = \frac{-Xx}{(n-1)(\log x)^{n-1}} + \frac{1}{n-1}\cdot\int_x \frac{P}{(\log x)^{n-1}};$$

and $$\int_x \frac{P}{(\log x)^{n-1}} = \frac{-P.x}{(n-2)(\log x)^{n-2}} + \frac{1}{n-2}\cdot\int_x \frac{Q}{(\log x)^{n-1}},$$

$$\text{where } Q = \frac{d.(P.x)}{dx};$$

$$\therefore \int_x \frac{X}{(\log x)^n} = \frac{-Xx}{(n-1)(\log x)^{n-1}} - \frac{Px}{(n-1)(n-2).(\log x)^{n-2}}$$

$$- \frac{Q.x}{(n-1)(n-2)(n-3)(\log x)^{n-3}} - \&c.$$

in this manner the integral may be reduced to $\int_x \frac{X_1}{(\log x)}$, which cannot be integrated except by a series.

Ex. 2. Find $\int_x \frac{x^m}{(\log x)^2}$.

$$\int_x \frac{x^m}{(\log x)^2} = \int_x \frac{x^{m+1}.\frac{1}{x}}{(\log x)^2} = \frac{-x^{m+1}}{\log x} + (m+1)\cdot\int_x \frac{x^m}{\log x}.$$

Let $\log x = z$; $\therefore x = e^z$, and $x^m = e^{mz}$;

$$\therefore \int_x \frac{x^m}{\log x} = \int_z \frac{e^{mz}}{z}\cdot\frac{dx}{dz} = \int_z \frac{e^{mz}}{z}e^z = \int_z \frac{e^{(m+1)z}}{z}$$

$$= \int_z \left\{1 + (m+1)z + \frac{(m+1)^2z^2}{1.2} + \frac{(m+1)^3z^3}{2,3} + \&c.\right\}.\frac{1}{z}$$

$$= \log x + (m+1) \cdot x + \frac{(m+1)^2 x^2}{1 \cdot 2^2} + \frac{(m+1)^3 x^3}{2 \cdot 3^2} + \&c.$$

$$= \log(\log x) + (m+1)\log x + \frac{(m+1)^2 (\log x)^2}{1 \cdot 2^2} + \frac{(m+1)^3 (\log x)^3}{2 \cdot 3^2} + \&c.$$

Cor. If $m = 0$, we have

$$\int_x \frac{1}{\log x} = \log(\log x) + (\log x) + \frac{(\log x)^2}{1 \cdot 2^2} + \&c.$$

62. Integrate $\int_x a^x \cdot X$, X being a function of x.

Since $\frac{d \cdot a^x}{dx} = a^x \log a = A a^x$; $\therefore \int_x a^x = \frac{a^x}{A}$;

$$\therefore \int_x a^x \cdot X = \frac{X a^x}{A} - \frac{1}{A} \cdot \int \frac{dX}{dx} \cdot a^x.$$

Let $\frac{dX}{dx} = P$, $\frac{dP}{dx} = Q$, &c.;

$$\therefore \int_x a^x \cdot X = \frac{X a^x}{A} - \frac{1}{A} \cdot \int_x P a^x,$$

$$\int_x a^x \cdot P = \frac{P a^x}{A} - \frac{1}{A} \cdot \int Q a^x;$$

$$\therefore \int_x a^x \cdot X = a^x \cdot \left\{ \frac{X}{A} - \frac{P}{A^2} + \frac{Q}{A^3} - \&c. \right\}.$$

Ex. 3. Let $\int_x x^m \cdot a^x$ be required.

$$\int_x x^m \cdot a^x = \frac{x^m \cdot a^x}{A} - \frac{m}{A} \cdot \int_x x^{m-1} \cdot a^x,$$

$$\int_x x^{m-1} \cdot a^x = \frac{x^{m-1} \cdot a^x}{A} - \frac{m-1}{A} \cdot \int_x x^{m-2} \cdot a^x;$$

$$\therefore \int_x x^m \cdot a^x = a^x \cdot \left\{ \frac{x^m}{A} - \frac{m x^{m-1}}{A^2} + \frac{m \cdot (m-1)\, x^{m-2}}{A^3} - \&c. \right\}.$$

Ex. 4. $$\int_x \frac{a^x}{x^n} = \frac{-a^x}{(n-1)\, x^{n-1}} + \frac{A}{n-1} \cdot \int_x \frac{a^x}{x^{n-1}};$$

$$\therefore \int_x \frac{a^x}{x^{n-1}} = \frac{-a^x}{(n-2) x^{n-2}} + \frac{A}{n-2} \cdot \int_x \frac{a^x}{x^{n-2}},$$

$$\therefore \int_x \frac{a^x}{x^n} = \frac{-a^x}{(n-1) \cdot x^{n-1}} - \frac{A a^x}{(n-1) \cdot (n-2) \cdot x^{n-2}}$$

$$- \frac{A^2 a^x}{(n-1).(n-2).(n-3).x^{n-3}} - \&c.$$

$$+ \frac{A^{n-1}}{(n-1).(n-2)\ldots 1} \cdot \int_x \frac{a^x}{x},$$

$$\text{Also } \int_x \frac{a^x}{x} = \int_x \left(\frac{1 + Ax + \frac{A^2x^2}{1.2} + \frac{A^3x^3}{2.3} + \&c.}{x} \right)$$

$$= \int_x \left(\frac{1}{x} + A + \frac{A^2 x}{1.2} + \frac{A^3 x^2}{2.3} + \&c. \right)$$

$$= \log x + Ax + \frac{A^2x^2}{1.2^2} + \frac{A^3x^3}{2.3^2} + \frac{A^4x^4}{2.3.4^2} + \&c.$$

Ex. 5. Find $\int_x \frac{1}{x} . \log(a + bx)$.

$$\log(a + bx) = \log a \left(1 + \frac{b}{a} x\right) = \log a + \log\left(1 + \frac{b}{a} x\right)$$

$$= \log a + \left(\frac{b}{a} x - \frac{b^2x^2}{2a^2} + \frac{b^3x^3}{3a^3} - \frac{b^4x^4}{4a^4} + \&c. \right);$$

$$\therefore \int_x \frac{1}{x} \log(a + bx) = \log x . \log a$$

$$+ \left(\frac{b}{a} x - \frac{b^2x^2}{2^2a^2} + \frac{b^3x^3}{3^2a^3} - \frac{b^4x^4}{4^2x^4} + \&c. \right).$$

Ex. 6. Find $\int_x x^{nx}$.

$$x^{nx} = 1 + nx \log x + \frac{(nx \log x)^2}{1.2} + \frac{(nx \log x)^3}{1.2.3} + \&c.;$$

$$\therefore \int_x x^{nx} = x + n . \int_x x \log x + \frac{n^2}{1.2} \cdot \int_x x^2 (\log x)^2$$

$$+ \frac{n^3}{2.3} \cdot \int_x x^3 (\log x)^3 + \&c.$$

Hence, the integration depends upon $\int_x x^m (\log x)^n$, Art. 60;

$$\therefore \int_x x \log x = \frac{x^2}{2} . (\log x - \tfrac{1}{2}),$$

$$\int_x x^2 (\log x)^2 = \frac{x^3}{3} \left\{ (\log x)^2 - \tfrac{2}{3} \log x + \frac{2.1}{3^2} \right\},$$

$$\int_x (x \log x)^3 = \frac{x^4}{4} \left\{ (\log x)^3 - \tfrac{3}{4} (\log x)^2 + \frac{3.2}{4^2} (\log x) - \frac{3.2}{4^3} \right\},$$

&c.

and arranging the terms according to the powers of $\log x$,

$$\int_x x^{nx} = x - \frac{nx^2}{2^2} + \frac{n^2x^3}{3^3} - \frac{n^3x^4}{4^4} + \frac{n^4x^5}{5^5} - \&c.$$
$$+ \left(\frac{x^2}{2} - \frac{nx^3}{3^2} + \frac{n^2x^4}{4^3} - \&c.\right) n \log x$$
$$+ \left(\frac{x^3}{3} - \frac{nx^4}{4^2} + \frac{n^2x^5}{5^3} - \&c.\right) \frac{n^2 (\log x)^2}{1.2}$$
$$+ \left(\frac{x^4}{4} - \frac{nx^5}{5^2} + \frac{n^2x^6}{6^3} - \&c.\right) \frac{n^3 (\log x)^3}{1.2.3}$$
$$+ \&c.$$

Cor. Since $x^n (\log x)^n = 0$; both when $x = 0$, and $x = 1$;

$$\therefore \int_x x^{nx}, \begin{matrix} \text{from } x = 0 \\ \text{to } x = 1 \end{matrix} \Big\} = 1 - \frac{n}{2^2} + \frac{n^2}{3^3} - \frac{n^3}{4^4} + \frac{n^4}{5^5} - \&c.,$$

and if $n = 1$, $\int_x x^x$ between the same values

$$= \frac{1}{1} - \frac{1}{2^2} + \frac{1}{3^3} - \frac{1}{4^4} + \frac{1}{5^5} - \&c.$$

This last integral gives the area of a curve, defined by $y = x^x$, included between two ordinates, each = 1, one drawn through the origin, and the other at a distance = 1 from it.

Ex. 7. $\int_x e^x \cdot \frac{x}{(1+x)^2}$,

$$e^x \cdot \frac{x}{(1+x)^2} = e^x \cdot \left\{\frac{1}{1+x} - \frac{1}{(1+x)^2}\right\} = \frac{d}{dx} \cdot \left(\frac{e^x}{1+x}\right);$$
$$\therefore \int_x e^x \cdot \frac{(1+x)^2}{x} = \frac{e^x}{1+x}.$$

63. Find $\int_t e^{-t^2}$ between $t = -\infty$ and $t = \infty$.

Now by Art. 49, between $x = 0$ and $x = 1$, we have

$$\left(\int_x \frac{x^{2n}}{\sqrt{1-x^2}}\right)\left(\int_x \frac{x^{2n+1}}{\sqrt{1-x^2}}\right) = \frac{1}{2n+1} \cdot \frac{\pi}{2}.$$

Let $x = e^{-qt^2}$; $\therefore \frac{dx}{dt} = -2qt \cdot e^{-qt^2}$;

$$\therefore 4q^2 \left\{\int_t \frac{te^{-q(2n+1)t^2}}{\sqrt{1-e^{-2qt^2}}} \times \int_t \frac{te^{-qt^2} \cdot e^{-qt^2(2n+1)}}{\sqrt{1-e^{-2qt^2}}}\right\} = \frac{1}{2n+1} \cdot \frac{\pi}{2},$$

the limits of t being $-\infty$ and 0; now let $q = \dfrac{1}{2n+1}$;

$$\therefore\ 2\left\{\int_t \frac{te^{-t^2}}{\sqrt{\dfrac{1-e^{-2qt^2}}{2q}}}\right\}\left\{\int_t \frac{te^{-t^2(1+q)}}{\sqrt{\dfrac{1-e^{-2qt^2}}{2q}}}\right\} = \frac{\pi}{2}.$$

Now making $q = 0$, in which case $\sqrt{\dfrac{1-e^{-2qt^2}}{2q}} = t$;

$$\therefore\ 2\left(\int_t e^{-t^2}\right)^2 = \frac{\pi}{2};\quad \therefore\ \int_t e^{-t^2} = \frac{1}{2}\sqrt{\pi};$$

but as the value of the integral is the same for t positive as t negative; $\therefore$ the integral, from $t = 0$ to $t = \infty$, $= \dfrac{1}{2}\sqrt{\pi}$;

$$\therefore \text{ from } t = -\infty \text{ to } t = +\infty;\quad \int_t e^{-t^2} = \sqrt{\pi}.$$

EXAMPLES.

(1) $\displaystyle\int_x x^4 (\log x) = \frac{x^5}{5}\left(\log x - \frac{1}{5}\right).$

(2) $\displaystyle\int_x x^2 (\log x)^2 = \frac{x^3}{3}\cdot\left\{(\log x)^2 - \frac{2}{3}\log x + \frac{2}{9}\right\}.$

(3) $\displaystyle\int_x x^3 (\log x)^3 = \frac{x^4}{4}\left\{(\log x)^3 - \frac{3}{4}(\log x)^2 + \frac{3.2}{4^2}\log x - \frac{3.2}{4^3}\right\}.$

(4) $\displaystyle\int_x \frac{1}{x}\log x = \frac{1}{2}(\log x)^2;\quad \int_x \frac{1}{x\log x} = \log.\log x.$

(5) $\displaystyle\int_x \frac{x^4}{(\log x)^3} = -\frac{x^5}{2(\log x)^2} - \frac{5x^5}{2.\log x} + \frac{25}{2}\cdot\int_x \frac{x^4}{\log x}.$

(6) $\displaystyle\int_x \frac{x^3}{\sqrt{\log x}} = \frac{x^4}{4\sqrt{\log x}}\cdot\left\{1 + \frac{1}{8\log x} + \frac{1.3}{(8.\log x)^2} + \frac{1.3.5}{(8.\log x)^3} + \&c.\right\}.$

(7) $\displaystyle\int_x a^x . x^3 = a^x\left\{\frac{x^3}{A} - \frac{3x^2}{A^2} + \frac{6x}{A^3} - \frac{6}{A^4}\right\}.$

(8) $\displaystyle\int_x e^x x^4 = e^x\left\{x^4 - 4x^3 + 12x^2 - 24x + 24\right\}.$

(9) $\displaystyle\int_x e^{-x} x^3 = -e^{-x}\left\{x^3 + 3x^2 + 6x + 6\right\}.$

(10) $$\int_x \frac{a^x}{x^4} = -a^x\left\{\frac{1}{3x^3} + \frac{A}{2.3x^2} + \frac{A^2}{2.3.x}\right\} + \frac{A^3}{2.3}\cdot\int_x \frac{a^x}{x}.$$

(11) $$\int_x \frac{a^x}{\sqrt{x}} = \frac{a^x}{A\sqrt{x}}\left\{1 + \frac{1}{2xA} + \frac{3}{(2xA)^2} + \frac{3.5}{(2xA)^3} + \&c.\right\},$$

also $$= \frac{Aa^x}{\sqrt{x}}\cdot\left\{\frac{2xA}{1} - \frac{(2xA)^2}{1.3} + \frac{(2xA)^3}{3.5} - \frac{(2xA)^4}{3.5.7} + \&c.\right\}.$$

(12) $$\int_x \frac{\log x}{(1+x)^2} = \frac{x}{x+1}\log x - \log(1+x).$$

(13) $$\int_x \frac{x}{\sqrt{1+x^2}}\log x = \sqrt{1+x^2}.\log\left(\frac{x}{e}\right) - \log\left(\frac{x}{1+\sqrt{1+x^2}}\right).$$

(14) $$\int_x \frac{1}{e^x+1} = \log\left(\frac{e^x}{e^x+1}\right).$$

(15) $$\int_x e^x\cdot\frac{x^2+1}{(x+1)^2} = e^x\cdot\left(\frac{x-1}{x+1}\right).$$

(16) $$\int_x \frac{e^x.(2-x^2)}{(1-x)\sqrt{1-x^2}} = e^x\cdot\sqrt{\frac{1+x}{1-x}}.$$

(17) $$\int_x \frac{xe^x}{(e^x-1)^3} = \log\frac{e^x}{e^x-1} - \frac{1}{e^x-1}\left\{1 + \frac{x}{2(e^x-1)}\right\}.$$

(18) $$\int x^{nx}.x^m \left\{\begin{matrix}\text{from } x=0\\ \text{to } x=1\end{matrix}\right\} = \frac{1}{m+1} - \frac{n}{(m+2)^2} + \frac{n^2}{(m+3)^3} - \frac{n^3}{(m+4)^4} + \&c.$$

CHAPTER V.

CIRCULAR FUNCTIONS.

64. THESE are of the form $\sin^n\theta$, $\cos^n\theta$, $(\sin\theta)^m(\cos\theta)^n$, $\frac{1}{(\sin\theta)^n}$, and $X.\sin^{-1}x$, where X is a function of x; these may be integrated by parts, and be reduced either to known or more simple functions such as,

$$\sin\theta,\ \cos\theta,\ \frac{1}{\cos^2\theta},\ \tan\theta,\ \cot\theta,\ \frac{1}{\sin\theta},\ \frac{1}{\cos\theta},\ \text{and}\ \frac{1}{\cos\theta\sin\theta}.$$

Also (1) $\int_\theta \sin\theta = -\cos\theta.$

(2) $\int_\theta \cos\theta = \sin\theta.$

(3) $\int_\theta \frac{1}{\cos^2\theta} = \tan\theta.$

(4) $\int_\theta \tan\theta = \int_\theta \frac{\sin\theta}{\cos\theta} = -\log\cos\theta.$

(5) $\int_\theta \cot\theta = \int_\theta \frac{\cos\theta}{\sin\theta} = \log\sin\theta.$

65. Integrate $\frac{1}{\sin\theta}$, $\frac{1}{\cos\theta}$, and $\frac{1}{\sin\theta\cos\theta}$.

(1) $$\int_\theta \frac{1}{\sin\theta} = \int_\theta \frac{\sin\theta}{1-\cos^2\theta} = \tfrac{1}{2}\int_\theta \left(\frac{\sin\theta}{1-\cos\theta} + \frac{\sin\theta}{1+\cos\theta}\right)$$

$$= \tfrac{1}{2}\log\left(\frac{1-\cos\theta}{1+\cos\theta}\right) = \tfrac{1}{2}\log\frac{\sin^2\frac{\theta}{2}}{\cos^2\frac{\theta}{2}} = \log\left(\tan\frac{\theta}{2}\right).$$

(2) $$\int_\theta \frac{1}{\cos\theta} = \int_\theta \frac{\cos\theta}{1-\sin^2\theta} = \tfrac{1}{2}\int_\theta \left\{\frac{\cos\theta}{1+\sin\theta} + \frac{\cos\theta}{1-\sin\theta}\right\}$$

$$= \tfrac{1}{2}.\log\left(\frac{1+\sin\theta}{1-\sin\theta}\right) = \log\tan\left(\frac{\pi}{4}+\frac{\theta}{2}\right).$$

(3) $$\int_\theta \frac{1}{\sin\theta\cos\theta} = \int_\theta \frac{\sin^2\theta+\cos^2\theta}{\sin\theta.\cos\theta} = \int_\theta \frac{\sin\theta}{\cos\theta} + \int_\theta \frac{\cos\theta}{\sin\theta}$$

$$= -\log\cos\theta + \log\sin\theta = \log(\tan\theta).$$

66. Find $\int_x X \cdot \sin^{-1}x$, where X is a function of x. Make $\int_x X = P$, and then integrating by parts,

$$\int_x X \cdot \sin^{-1}x = P\sin^{-1}x - \int_x \frac{P}{\sqrt{1-x^2}},$$

and $\int_x \frac{P}{\sqrt{1-x^2}}$ has been integrated.

Ex. $\int_x \frac{x}{\sqrt{1-x^2}} \sin^{-1}x$. Here $P = -\sqrt{1-x^2}$;

$$\therefore \int_x \frac{x\sin^{-1}x}{\sqrt{1-x^2}} = -\sqrt{1-x^2}\cdot\sin^{-1}x + \int_x \sqrt{1-x^2}\frac{1}{\sqrt{1-x^2}}$$

$$= -\sqrt{1-x^2}\cdot\sin^{-1}x + x.$$

Similarly may $\int_x X\cos^{-1}x$ be integrated.

67. To integrate $\frac{du}{dx} = X\tan^{-1}x$.

Let $\int_x X = P$;

$$\therefore \int_x X\tan^{-1}x = P\tan^{-1}x - \int_x \frac{P}{1+x^2}.$$

68. Integrate $\frac{du}{d\theta} = \sin^n\theta$.

Integrating by parts, since $\sin^n\theta = \sin^{n-1}\theta \cdot \sin\theta$;

$$\therefore \int_\theta \sin^n\theta = \int \sin^{n-1}\theta \cdot \sin\theta$$

$$= -\sin^{n-1}\theta\cdot\cos\theta + (n-1)\cdot\int_\theta \sin^{n-2}\theta\cos^2\theta;$$

and putting $1-\sin^2\theta$ for $\cos^2\theta$

$$= -\sin^{n-1}\theta\cos\theta + (n-1)\cdot\int_\theta \sin^{n-2}\theta - (n-1)\int_\theta \sin^n\theta;$$

$$\therefore \int_\theta \cdot \sin^n\theta = -\frac{\sin^{n-1}\theta\cdot\cos\theta}{n} + \frac{n-1}{n}\int_\theta \sin^{n-2}\theta,$$

a formula by which $\int_\theta \sin^n\theta$ may be reduced to $-\cos\theta$, or θ, according as n is odd or even.

Suppose n to be even or $= 2m$, to find the value of $\int_\theta (\sin\theta)^{2m}$ between $\theta = 0$, and $\theta = \frac{\pi}{2}$.

Let $\int_\theta (\sin\theta)^{2m} = P_{2m}$; $\sin^{2m-1}\theta\cos\theta = Q_{2m-1}$;

$$\therefore P_{2m} = -\frac{1}{2m}\cdot Q_{2m-1} + \frac{2m-1}{2m}P_{2m-2}.$$

But $Q_{2m-1} = 0$ both for $\theta = 0$ and $\theta = \frac{\pi}{2}$;

$$\therefore P_{2m} = \frac{2m-1}{2m} P_{2m-2}; \quad \therefore P_{2m-2} = \frac{2m-3}{2m-2} \cdot P_{2m-4};$$

$$\therefore P_{2m} = \frac{(2m-1)\,.\,(2m-3)}{2m\,.\,(2m-2)} \cdot P_{2m-4}; \text{ hence } \therefore$$

$$= \frac{(2m-1)\,.\,(2m-3)\ldots 3\,.\,1}{2m\,.\,(2m-2)\,.\,4\,.\,2} \cdot P_0$$

$$= \frac{1\,.\,3\,.\,5\ldots(2m-1)}{2\,.\,4\,.\,6\ldots 2m} \cdot \frac{\pi}{2}.$$

69. Integrate $\frac{du}{d\theta} = \cos^n\theta$.

$$\int_\theta \cos^n\theta = \int_\theta \cos^{n-1}\theta\,.\,\cos\theta$$

$$= +\cos^{n-1}\sin\theta + (n-1)\,.\int_\theta \cos^{n-2}\theta\sin^2\theta$$

$$= \cos^{n-1}\theta\sin\theta + (n-1)\int_\theta \cos^{n-2}\theta - (n-1)\int_\theta \cos^n\theta$$

$$= \frac{\cos^{n-1}\theta\sin\theta}{n} + \frac{n-1}{n}\int_\theta \cos^{n-2}\theta,$$

a formula by which $\int_\theta \cos^n\theta$ may be reduced to $\sin\theta$ or θ, according as n is odd or even.

70. Let $\frac{du}{d\theta} = \frac{1}{(\sin\theta)^n}$. Since $\sin^2\theta + \cos^2\theta = 1$;

$$\therefore u = \int_\theta \frac{\sin^2\theta + \cos^2\theta}{(\sin\theta)^n} = \int \frac{1}{(\sin\theta)^{n-2}} + \int_\theta \frac{\cos^2\theta}{(\sin\theta)^n},$$

and $\int_\theta \frac{\cos^2\theta}{(\sin\theta)^n} = -\frac{\cos\theta}{(n-1)\,(\sin\theta)^{n-1}} - \frac{1}{n-1}\int_\theta \frac{\sin\theta}{(\sin\theta)^{n-1}}$;

$$\therefore u = -\frac{\cos\theta}{(n-1)\,(\sin\theta)^{n-1}} + \left(1 - \frac{1}{n-1}\right)\int_\theta \frac{1}{(\sin\theta)^{n-2}}$$

$$= -\frac{\cos\theta}{(n-1)\,(\sin\theta)^{n-1}} + \frac{n-2}{n-1}\cdot\int_\theta \frac{1}{(\sin\theta)^{n-2}},$$

a formula by which n may be diminished.

71. If $\frac{du}{d\theta} = \frac{1}{(\cos\theta)^n}$, then, as in last article,

$$u = \int_\theta \frac{1}{(\cos\theta)^{n-2}} + \int_\theta \frac{\sin^2\theta}{(\cos\theta)^n},$$

and $\int_\theta \frac{\sin^2\theta}{(\cos\theta)^n} = \frac{\sin\theta}{(n-1)\,.\,(\cos\theta)^{n-1}} - \frac{1}{n-1}\int_\theta \frac{\cos\theta}{(\cos\theta)^{n-1}}$;

$$\therefore\ u = \frac{\sin\theta}{(n-1)\,(\cos\theta)^{n-1}} + \frac{n-2}{n-1}\int_\theta \frac{1}{(\cos\theta)^{n-2}}\,.$$

72. Let $\frac{du}{d\theta} = (\sin\theta)^m\,(\cos\theta)^n$, m and n both integers,

$$(\sin\theta)^m\,(\cos\theta)^n = (\sin\theta)^m \cos\theta\,(\cos\theta)^{n-1};$$

$$\therefore \int_\theta (\sin\theta)^m(\cos\theta)^n = \frac{(\sin\theta)^{m+1}(\cos\theta)^{n-1}}{m+1} + \frac{n-1}{m+1}\int_\theta (\sin\theta)^{m+2}(\cos\theta)^{n-2}$$

$$= \frac{(\sin\theta)^{m+1}(\cos\theta)^{n-1}}{m+1} + \frac{n-1}{m+1}\left\{\int_\theta (\sin\theta)^m(\cos\theta)^{n-2} - \int_\theta (\sin\theta)^m(\cos\theta)^n\right\};$$

$$\therefore\ \left(1 + \frac{n-1}{m+1}\right)u = \frac{m+n}{m+1}\,u = \frac{(\sin\theta)^{m+1}(\cos\theta)^{n-1}}{m+1}$$

$$+ \frac{n-1}{m+1}\cdot\int_\theta (\sin\theta)^m(\cos\theta)^{n-2};$$

$$\therefore\ u = \frac{(\sin\theta)^{m+1}(\cos\theta)^{n-1}}{m+n} + \frac{n-1}{m+n}\cdot\int_\theta (\sin\theta)^m(\cos\theta)^{n-2},$$

a formula by which the integral may be reduced to

$$\int_\theta (\sin\theta)^m, \text{ or } \int_\theta (\sin\theta)^m\cos\theta.$$

73. Let $\frac{du}{d\theta} = \frac{\sin^m\theta}{\cos^n\theta}$,

$$u = \int_\theta \frac{\sin^{m-1}\theta\,\sin\theta}{(\cos\theta)^n} = \frac{(\sin\theta)^{m-1}}{(n-1)\,.\,(\cos\theta)^{n-1}} - \frac{m-1}{n-1}\int_\theta \frac{(\sin\theta)^{m-2}}{(\cos\theta)^{n-2}},$$

a formula by which the integral is reducible to a known form.

74. Let $\frac{du}{d\theta} = \theta^n\,.\,\sin\theta$.

$$\int_\theta \theta^n \sin\theta = -\theta^n\cos\theta + n\,.\int\theta^{n-1}\cos\theta,$$

$$\int_\theta \theta^{n-1}\cos\theta = +\theta^{n-1}\sin\theta - (n-1)\int_\theta \theta^{n-2}\sin\theta,$$

$$\int_\theta \theta^{n-2}\sin\theta = -\theta^{n-2}\cos\theta + (n-2)\int_\theta \theta^{n-3}\cos\theta,$$

$$\&c. \quad = \quad \&c. \quad \&c.$$

$$\int_\theta \theta^n\sin\theta = -\theta^n\cos\theta + n\theta^{n-1}\sin\theta + n\,(n-1)\,\theta^{n-2}\cos\theta$$

$$- n\,(n-1)\,(n-2)\,\theta^{n-3}\sin\theta - \&c.$$

Cor. Similarly may $\int_\theta \theta^n \cos\theta$ be found and shewn to be

$$= \theta^n \sin\theta + n\theta^{n-1}\cos\theta - n(n-1)\theta^{n-2}\sin\theta$$
$$- n(n-1)(n-2)\theta^{n-3}\cos\theta + \&c.$$

75. Let $\frac{du}{d\theta} = \theta^{-n}\sin\theta = \frac{\sin\theta}{\theta^n}$;

$$\therefore \int_\theta \frac{\sin\theta}{\theta^n} = -\frac{\sin\theta}{(n-1)\theta^{n-1}} + \frac{1}{n-1}\int_\theta \frac{\cos\theta}{\theta^{n-1}},$$

and $$\int_\theta \frac{\cos\theta}{\theta^{n-1}} = -\frac{\cos\theta}{(n-2\theta)^{n-2}} - \frac{1}{n-2}\cdot\int_\theta \frac{\sin\theta}{\theta^{n-2}};$$

$$\therefore \int_\theta \frac{\sin\theta}{\theta^n} = -\frac{\sin\theta}{(n-1)\theta^{n-1}} - \frac{\cos\theta}{(n-1)(n-2)\theta^{n-2}} - \frac{1}{(n-1)(-2)}\cdot\int\frac{\sin\theta}{\theta^{n-2}}$$

$$= -\frac{\sin\theta}{(n-1)\theta^{n-1}} - \frac{\cos\theta}{(n-1)(n-2)\theta^{n-2}} + \frac{\sin\theta}{(n-1)(n-2)(n-3)\theta^{n-3}} + \&c.$$

by which the integral may be reduced to $\int_\theta \frac{\sin\theta}{\theta}$, (if n be an integer) $= \int_\theta \left\{1 - \frac{\theta^2}{2.3} + \frac{\theta^4}{2.3.4.5} - \&c.\right\} = \theta - \frac{\theta^3}{2.3^2} + \frac{\theta^5}{2.3.4.5} - \&c.$

a similar method applies to $\int_\theta \frac{\cos\theta}{\theta^n}$.

76. Integrate $\sin m\theta \cdot \cos n\theta$, $\sin m\theta \cdot \sin n\theta$, and $\cos m\theta \cdot \cos n\theta$.

Since $\sin A \cdot \cos B = \frac{1}{2}\cdot\{\sin(A+B) + \sin(A-B)\}$;

$$\therefore \sin m\theta \cdot \cos n\theta = \tfrac{1}{2}\cdot\{\sin(m+n)\theta + \sin(m-n)\theta\};$$

$$\therefore \int_\theta (\sin m\theta \cdot \cos n\theta) = -\tfrac{1}{2}\cdot\left\{\frac{\cos(m+n)\theta}{m+n} + \frac{\cos(m-n)\theta}{m-n}\right\}.$$

Also since $\cos m\theta \cdot \cos n\theta = \frac{1}{2}\cdot\{\cos(m+n)\theta + \cos(m-n)\theta\}$,

and $\sin m\theta \cdot \sin n\theta = \frac{1}{2}\cdot\{\cos(m-n)\theta - \cos(m+n)\theta\}$;

$$\therefore \int_\theta (\cos m\theta \cdot \cos n\theta) = \tfrac{1}{2}\cdot\left\{\frac{\sin(m+n)\theta}{m+n} + \frac{\sin(m-n)\theta}{m-n}\right\},$$

and $$\int_\theta (\sin m\theta \sin n\theta) = -\tfrac{1}{2}\cdot\left\{\frac{\sin(m+n)\theta}{m+n} + \frac{\sin(m-n)\theta}{m-n}\right\}.$$

Cor. Similarly if $\frac{du}{d\theta} = \sin(a+m\theta)\cdot\cos(b+n\theta)$,

put for $\sin(a+m\theta)\cdot\cos(b+n\theta)$ its equivalent expression $\frac{1}{2}[\sin\{a+b+(m+n)\theta\} + \sin\{a-b+(m-n)\theta\}]$.

77. Integrate $(\tan\theta)^m$, and $(\tan\theta)^{-m}$.

$$(\tan\theta)^m = (\tan\theta)^{m-2}\{1+\tan^2\theta-1\}$$

$$= (\tan\theta)^{m-2}\frac{d.\tan\theta}{d\theta} - (\tan\theta)^{m-2};$$

$$\therefore \int_\theta (\tan\theta)^m = \frac{(\tan\theta)^{m-1}}{m-1} - \int_\theta (\tan\theta)^{m-2},$$

$$\int_\theta (\tan\theta)^{m-2} = \frac{(\tan\theta)^{m-3}}{m-3} - \int_\theta (\tan\theta)^{m-4},$$

&c. &c.

$$\therefore \int_\theta (\tan\theta)^m = \frac{(\tan\theta)^{m-1}}{m-1} - \frac{(\tan\theta)^{m-3}}{m-3} + \frac{(\tan\theta)^{m-5}}{m-5} - \text{&c.}$$

by which u is reduced either to θ, or $\int_\theta \tan\theta = -\log\cos\theta$.

78. $$u = \int_\theta \frac{1}{(\tan\theta)^m} = \int_\theta \frac{1+\tan^2\theta-\tan^2\theta}{(\tan\theta)^m}$$

$$= \int_\theta \frac{\frac{d.(\tan\theta)}{d\theta}}{(\tan\theta)^m} - \int_\theta \frac{1}{(\tan\theta)^{m-2}} = -\frac{1}{(m-1)(\tan\theta)^{m-1}} - \int_\theta \frac{1}{(\tan\theta)^{m-2}}$$

and $$\int_\theta \frac{1}{(\tan\theta)^{m-2}} = -\frac{1}{(m-3)(\tan\theta)^{m-3}} - \int_\theta \frac{1}{(\tan\theta)^{m-4}};$$

$$\therefore \int_\theta \frac{1}{(\tan\theta)^m} = -\frac{1}{(m-1)(\tan\theta)^{m-1}}$$

$$+ \frac{1}{(m-3)(\tan\theta)^{m-3}} - \frac{1}{m-5\,(\tan\theta)^{m-5}} + \text{&c.}$$

and thus u may be reduced to θ, or $\int_\theta \frac{1}{\tan\theta} = \log(\sin\theta)$.

79. $\int_x e^{ax}\sin kx$.

$$\int_x e^{ax}\sin kx = \frac{e^{ax}\sin kx}{a} - \frac{k}{a}\cdot\int_x e^{ax}\cos kx \ldots\ldots\ldots (1),$$

and $$\int_x e^{ax}\cos kx = \frac{e^{ax}\cos kx}{a} + \frac{k}{a}\cdot\int_x e^{ax}\sin kx.$$

Multiplying by $\frac{k}{a}$, and transposing,

$$\frac{k^2}{a^2}\cdot\int_x e^{ax}\sin kx = -\frac{ke^{ax}\cos kx}{a^2} + \frac{k}{a}\cdot\int_x e^{ax}\cos kx \ldots\ldots (2).$$

Adding (1) and (2),

$$\left(1+\frac{k^2}{a^2}\right)\int e^{ax}\sin kx = \frac{(a\,.\,\sin kx - k\cos kx)e^{ax}}{a^2};$$

$$\therefore \int e^{ax}\sin kx = \frac{(a\,.\,\sin kx - k\cos kx)e^{ax}}{k^2+a^2}.$$

80. To integrate $\dfrac{du}{dx} = \dfrac{1}{a+b\,.\,\cos x}$.

Since $\cos x = \dfrac{1-\tan^2\frac{x}{2}}{1+\tan^2\frac{x}{2}}$. Let $z = \tan\dfrac{x}{2}$;

$$\therefore \cos x = \frac{1-z^2}{1+z^2}; \quad \therefore \sin x\frac{dx}{dz} = \frac{4z}{(1+z^2)^2}.$$

But $\sin x = \sqrt{1-\left(\dfrac{1-z^2}{1+z^2}\right)^2} = \dfrac{2z}{1+z^2}$; $\therefore \dfrac{dx}{dz} = \dfrac{2}{1+z^2}$.

$$\therefore \frac{du}{dz} = \frac{1}{a+b\dfrac{1-z^2}{1+z^2}}\cdot\frac{2}{1+z^2} = \frac{2}{a(1+z^2)+b(1-z^2)}.$$

(1) Let $a>b$; $\therefore \dfrac{du}{dz} = \dfrac{2}{a+b+(a-b)z^2} = \dfrac{2}{a+b}\cdot\dfrac{1}{1+\dfrac{a-b}{a+b}\cdot z^2}$;

$$\therefore u = \frac{2}{a+b}\cdot\sqrt{\frac{a+b}{a-b}}\cdot\tan^{-1}\left(z\sqrt{\frac{a-b}{a+b}}\right)$$

$$= \frac{2}{\sqrt{a^2-b^2}}\tan^{-1}\left\{\sqrt{\frac{a-b}{a+b}}\cdot\tan\frac{x}{2}\right\}.$$

(2) Let $a<b$; $\quad \therefore \dfrac{du}{dz} = \dfrac{2}{(b-a)}\cdot\dfrac{1}{\dfrac{b+a}{b-a}-z^2}$;

$$\therefore u = \frac{1}{b-a}\cdot\sqrt{\frac{b-a}{b+a}}\cdot\log\frac{\sqrt{\dfrac{b+a}{b-a}}+z}{\sqrt{\dfrac{b+a}{b-a}}-z}$$

$$= \frac{1}{\sqrt{b^2-a^2}}\cdot\log\frac{\sqrt{b+a}+\sqrt{b-a}\,.\tan\dfrac{x}{2}}{\sqrt{b+a}-\sqrt{b-a}\,.\tan\dfrac{x}{2}}.$$

81. Similarly may $\int_x \frac{1}{a + b \sin x}$ be found.

Also $\int_x \frac{\sin x}{a + b \cos x} = \frac{1}{b} \int_x \frac{-\overline{d(b \cos x)}}{a + b \cos x} = -\frac{1}{b} . \log (a + b \cos x)$,

and $\int_x \frac{\cos x}{a + b \cos x} = \int_x \frac{a + b \cos x - a}{b (a + b \cos x)}$

$= \int_x \left\{ \frac{1}{b} - \frac{a}{b} . \frac{1}{a + b \cos x} \right\} = \frac{x}{b} - \frac{a}{b} . \int_x \frac{1}{a + b \cos x}$.

82. Integrate $\frac{du}{dx} = \frac{1}{a + b (\cos x)^2}$.

$$\int_x \frac{1}{a + b (\cos x)^2} = \int_x \frac{\sec^2 x}{a \sec^2 x + b}$$

$$= \int_x \frac{\sec^2 x}{a + b + a \tan^2 x} = \frac{1}{a + b} \int_x \frac{\frac{d}{dx} \tan x}{1 + \frac{a}{a + b} \tan^2 x}$$

$$= \frac{1}{\sqrt{a^2 + ab}} . \tan^{-1} \left(\sqrt{\frac{a}{a + b}} . \tan x \right).$$

83. If $\frac{du}{dx} = \frac{1}{a + b \tan x}$. Let $\tan x = z$; $\therefore \frac{dx}{dz} = \frac{1}{1 + z^2}$;

$$\therefore \int_x \frac{1}{a + b \tan x} = \int_z \frac{1}{(1 + z^2)(a + bz)},$$

which may be integrated by partial fractions.

84. If $\frac{du}{dx} = \frac{a' + b' \cos x}{(a + b \cos x)}$.

Let $u = \frac{A \sin x}{(a + b \cos x)^{m-1}} + \int_x \frac{B + C \cos x}{(a + b \cos x)^{m-1}}$;

$\therefore$ differentiating and omitting the denominators,

$$a' + b' \cos x = A \cos x (a + b \cos x) + (m - 1) Ab \sin^2 x$$
$$+ (B + C \cos x)(a + b \cos x);$$
$$\therefore (m - 1) Ab + Ba - a' + (Aa + Bb + Ca - b') \cos x$$
$$- \{(m - 2) A - C\} \cos^2 x = 0,$$

whence by putting the coefficients of like terms = 0,

$$A = \frac{ab' - ba'}{(m-1)(a^2-b^2)}; \quad B = \frac{aa' - bb'}{a^2-b^2}; \quad C = (m-2)A;$$

$$\therefore u = \frac{(ab' - ba')\sin x}{(m-1)(a^2-b^2)(a+b\cos x)^{m-1}}$$

$$+ \int_x \frac{(m-1)(aa' - bb') + (m-2)(ab' - a'b)\cos x}{(m-1)(a^2-b^2)(a+b\cos x)^{m-1}}.$$

Cor. 1. If $b' = 0$ and $a' = 1$, $\int_x \frac{1}{(a+b\cos x)^m}$

$$= \frac{1}{(m-1)(a^2-b^2)}\left\{\frac{-b\sin x}{(a+b\cos x)^{m-1}}\right.$$

$$\left. + \int_x \frac{(m-1)a - (m-2)b\cos x}{(a+b\cos x)^{m-1}}\right\}.$$

Cor. 2. If $a' = 0$ and $b' = 1$; $\int_x \frac{\cos x}{(a+b\cos x)^m}$

$$= \frac{1}{(m-1)(a^2-b^2)}\left\{\frac{a\sin x}{(a+b\cos x)^{m-1}} - \int_x \frac{(m-1)b - (m-2)a\cos x}{(a+b\cos x)^{m-1}}\right\}.$$

85. Integrate $\int_x (a+b\cos x)^m$ by means of multiple arcs.

$$(a+b\cos x)^m = a^m\left(1+\frac{b}{a}\cos x\right)^m = a^m(1+n\cos x)^m; \quad n = \frac{b}{a};$$

$$\text{but } (1+n\cos x)^m = \left\{1 + \frac{n}{2}(e^{x\sqrt{-1}} + e^{-x\sqrt{-1}})\right\}^m$$

$$= 1 + \frac{mn}{2}(e^{x\sqrt{-1}} + e^{-x\sqrt{-1}}) + \frac{m(m-1)}{1.2}\frac{n^2}{4}(e^{x\sqrt{-1}} + e^{-x\sqrt{-1}})^2$$

$$+ \frac{m(m-1).(m-2)}{1.2.3}\frac{n^3}{8}.(e^{x\sqrt{-1}} + e^{-x\sqrt{-1}})^3 + \&c.$$

$$= 1 + mn\cos x + \frac{m(m-1)}{1.2}.\frac{n^2}{4}(2\cos 2x + 2)$$

$$+ \frac{m(m-1).(m-2)}{1.2.3}\frac{n^3}{8}(2\cos 3x + 6\cos x)$$

$$+ \frac{m(m-1).(m-2).(m-3)}{1.2.3.4}\frac{n^4}{16}(2\cos 4x + 8\cos 2x + 6) + \&c.$$

$$= A_0 + A_1\cos x + A_2\cos 2x + A_3\cos 3x + A_4\cos 4x + \&c.$$

where $A_0 = 1 + \frac{m(m-1)}{2.2}n^2 + \frac{m(m-1).(m-2).(m-3).n^4}{2.2.4.4} + \&c.$

$$A_1 = 2n\left\{\frac{m}{2} + \frac{m(m-1).(m-2)n^3}{2.2.4} + \frac{m\ldots(m-4).n^4}{2.2.4.4.6} + \&c.\right\};$$

to find A_2, A_3, &c., take the logarithms and differentiate;

$$\therefore \frac{mn\sin x}{1+n\cos x} = \frac{A_1\sin x + 2A_2\sin 2x + 3A_3\sin 3x + \&c.}{A_0 + A_1\cos x + A_2\cos 2x + \&c.}.$$

Then $\because \sin x\cos\alpha x = \frac{1}{2}\{\sin(\alpha+1)x - \sin(\alpha-1)x\}$

$$\cos x\sin\beta x = \tfrac{1}{2}\{\sin(\beta+1)x + \sin(\beta-1)x\};$$

therefore multiplying out and arranging the terms according to the sines of the multiple arc;

$$0 = (A_1 + \frac{2A_2}{2}n - mA_0n + \frac{m}{2}A_2n)\sin x$$

$$+ (2A_2 + \frac{1}{2}A_1n + \frac{3A_3}{2}n - \frac{m}{2}A_1n + \frac{m}{2}A_3n)\sin 2x$$

$$+ (3A_3 + \frac{2}{2}A_2n + \frac{4A_4}{2}n - \frac{m}{2}A_2n + \frac{m}{2}A_4n)\sin 3x$$

$$+ \&c.;$$

$$\therefore A_2 = \frac{2mnA_0 - 2A_1}{(m+2)n};\quad A_3 = \frac{(m-1)A_1n - 4A_2}{(m+3)n};$$

$$A_4 = \frac{(m-2)A_2n - 6A_3}{(m+4)n},$$

hence if A_0, A_1 are known, the other coefficients are also known.

86. When $m = -1$, or $u = \int_x \frac{1}{1+n\cos x}$.

$$A_0 = 1 + \frac{1}{2}n^2 + \frac{1.3}{2.4}n^4 + \&c. = \frac{1}{\sqrt{1-n^2}};$$

$$A_1 = 2n\left(-\frac{1}{2} - \frac{1.3}{2.4}n^2 - \&c.\right) = \frac{2}{n}.\left(1 - \frac{1}{\sqrt{1-n^2}}\right) = \frac{2}{n}(1-A_0),$$

$$A_2 = -\frac{2}{n}.(nA_0 + A_1);\quad A_3 = -\frac{1}{n}.(2A_1n + 2A_2);$$

$$A_4 = -\frac{1}{n}(A_2n + 2A_2);$$

similarly we may find the coefficients when $m = -\frac{1}{2}$, $m = -\frac{3}{2}$; the latter case is useful in Physical Astronomy.

87. Let $\frac{du}{dx} = \log(1 + n\cos x)$.

$$\log(1+n\cos x) = n\cos x - \tfrac{1}{2}n^2\cos^2 x + \tfrac{1}{3}n^3\cos^3 x - \tfrac{1}{4}n^4\cos^4 x + \&c.$$

$$= -\frac{1}{2}\frac{n^2}{2} - \frac{1.3}{2.4}\frac{n^4}{4} - \frac{1.3.5}{2.4.6}\frac{n^6}{6} - \&c.$$

$$+ \left(n + \frac{3}{4}\frac{n^3}{3} + \frac{3.5}{4.6}\frac{n^5}{5} + \&c.\right)\cos x + \&c.$$

$$= -A_0 + A_1\cos x - A_2\cos 2x + A_3\cos 3x + \&c. \quad (1);$$

where $A_0 = \frac{1}{2}\frac{n^2}{2} + \frac{1.3}{2.4}\frac{n^4}{4} + \frac{1.3.5}{2.4.6}\cdot\frac{n^6}{6} + \&c.$;

$$\therefore\ n\frac{dA_0}{dn} = \frac{1}{2}.n^2 + \frac{1.3}{2.4}.n^4 + \frac{1.3.5}{2.4.6}.n^6 + \&c. = \frac{1}{\sqrt{1-n^2}} - 1;$$

$$\therefore\ \frac{dA_0}{dn} = \frac{1}{n\sqrt{1-n^2}} - \frac{1}{n};\quad \therefore\ A_0 = \log\frac{1-\sqrt{1-n^2}}{n} + \log\frac{1}{n} + C,$$

and $C = \log 2$, for $A_0 = 0$, when $n = 0$;

$$\therefore\ A_0 = \log\left(\frac{2-2\sqrt{1-n^2}}{n^2}\right),$$

and $A_1 = n + \frac{1.3}{2.4}\cdot\frac{n^3}{3} + \frac{1.3.5}{4.6}\cdot\frac{n^5}{5} + \&c.$;

$$\therefore\ \frac{dA_1}{dn} = \frac{2}{n^2}\left\{\frac{n^2}{2} + \frac{1.3}{4}\cdot\frac{n^4}{2} + \frac{1.3.5}{4.6}\cdot\frac{n^6}{2} + \&c.\right\}$$

$$= \frac{2}{n^2\sqrt{1-n^2}} - \frac{2}{n^2};$$

$$\therefore\ A_1 = 2\left(\frac{1-\sqrt{1-n^2}}{n}\right) + C = 2\left(\frac{1-\sqrt{1-n^2}}{n}\right);\ \because\ C = 0;$$

$$\therefore\ \text{if } \frac{1-\sqrt{1-n^2}}{n} = N;\quad A_0 = \log\left(\frac{2N}{n}\right);\quad A_1 = 2N;$$

and to find A_2, A_3, &c. differentiate (1);

$$\therefore\ \frac{n\sin x}{1+n\cos x} = A_1\sin x - 2A_2\sin 2x + 3A_3\sin 3x - 4A_4\cos 4x + \&c.$$

$$\therefore\ A_2 = \frac{A_1 - n}{n};\quad A_3 = \frac{4A_2 - A_1 n}{3n};\quad A_4 = \frac{6A_2 - 2A_2 n}{4n};$$

$$\therefore\ A_2 = \frac{2}{2}N^2,\quad A_3 = \frac{2}{3}N^3,\quad A_4 = \frac{2}{4}N^4,\quad A_5 = \frac{2}{5}N^5,\ \&c.;$$

$$\therefore \int_x \log(1 + n\cos x) = \int_x \left\{\log\left(\frac{n}{2N}\right) + 2N\cos x - \frac{2}{2}N^2\cos 2x + \&c.\right\}$$

$$= x\log\frac{n}{2N} + 2N\sin x - \frac{2}{2.2}N^2.\sin 2x + \frac{2}{3.3}N^3.\sin 3x - \&c.$$

Cor. If $n = 1$; $\therefore N = 1$, and $\log(1 + \cos x) = \log\left(2\cos^2\frac{x}{2}\right)$;

$$\therefore\ = \log 2 + 2\log\cos\frac{x}{2};\ \text{hence putting } 2x \text{ for } x,$$

$$\log\cos x = \log\tfrac{1}{2} + \cos 2x - \tfrac{1}{2}\cos 4x + \tfrac{1}{3}\cos 6x - \&c.$$

And $n = -1$; $\therefore N = -1$, and $\log(1 - \cos x) = \log 2 - 2\log\sin\frac{x}{2}$

$$= -\log\tfrac{1}{2} - 2\log\sin\frac{x}{2};\quad \therefore \text{ putting } 2x \text{ for } x,$$

$$\log\sin x = \log\tfrac{1}{2} - \cos 2x - \tfrac{1}{2}\cos 4x - \tfrac{1}{3}\cos 6x - \&c.;$$

$$\therefore \int_x \log\cos x = x\log\tfrac{1}{2} + \tfrac{1}{2}\sin 2x - \frac{1}{2.4}\sin 4x + \frac{1}{3.6}\sin 6x - \&c.$$

$$\int_x \log\sin x = x\log\tfrac{1}{2} - \tfrac{1}{2}\sin 2x - \frac{1}{2.4}\sin 4x - \frac{1}{3.6}\sin 6x - \&c.$$

and $\int_x \log\tan x = -\sin 2x - \frac{1}{9}\sin 6x - \&c.$

EXAMPLES.

(1) $\int_\theta (\sin\theta)^3 = -\frac{1}{3}.(\sin\theta)^2.\cos\theta - \frac{2}{3}\cos\theta.$

(2) $\int_\theta (\cos\theta)^3 = \frac{1}{3}.(\cos\theta)^2.\sin\theta + \frac{2}{3}.\sin\theta.$

(3) $\int_\theta (\sin\theta)^5 = -\cos\theta\left\{\frac{(\sin\theta)^4}{5} + \frac{4(\sin\theta)^2}{15} + \frac{8}{15}\right\}.$

(4) $\int_\theta (\cos\theta)^6 = \sin\theta\left\{\frac{(\cos\theta)^5}{6} + \frac{5(\cos\theta)^3}{24} + \frac{5\cos\theta}{16}\right\} + \frac{5\theta}{16}.$

(5) $\int_\theta (\sin\theta)^3.(\cos\theta)^2 = \frac{\cos\theta}{5}\left\{(\sin\theta)^4 - \frac{1}{3}(\sin\theta)^2 - \frac{2}{3}\right\}.$

(6) $\int_\theta (\sin\theta)^2 (\cos\theta)^4 = \frac{(\sin\theta \cdot \cos\theta)^3}{6} + \frac{1}{8}\sin^3\theta\cos\theta - \frac{1}{16}\sin\theta\cos\theta + \frac{\theta}{16}.$

(7) $\int_\theta (\sin\theta)^6 (\cos\theta)^3 = \left\{\frac{(\cos\theta)^2}{9} + \frac{2}{63}\right\}(\sin\theta)^7$

(8) $\int_\theta \frac{1}{(\sin\theta)^3} = -\frac{1}{2}\cdot\frac{\cos\theta}{(\sin\theta)^2} + \frac{1}{2}\log\left(\tan\frac{\theta}{2}\right).$

(9) $\int_\theta \frac{1}{(\sin\theta)^5} = -\cos\theta\left\{\frac{1}{4(\sin\theta)^4} + \frac{3}{8(\sin\theta)^2}\right\} + \frac{3}{8}\log\tan\frac{\theta}{2}.$

(10) $\int_\theta \frac{1}{(\cos\theta)^6} = \sin\theta\left\{\frac{1}{5(\cos\theta)^5} + \frac{4}{15(\cos\theta)^3} + \frac{8}{15\cdot\cos\theta}\right\}.$

(11) $\int_\theta \frac{(\sin\theta)^3}{(\cos\theta)^4} = \frac{1}{(\cos\theta)^3}\left\{(\sin\theta)^2 - \frac{2}{3}\right\}.$

(12) $\int_\theta \frac{(\sin\theta)^5}{(\cos\theta)^2} = -\frac{1}{\cos\theta}\left\{\frac{(\sin\theta)^4}{3} + \frac{4(\sin\theta)^2}{3} - \frac{8}{3}\right\}.$

(13) $\int_\theta \frac{(\cos\theta)^4}{(\sin\theta)^3} = \left\{(\cos\theta)^3 - \frac{3\cos\theta}{2}\right\}\frac{1}{(\sin\theta)^2} - \frac{3}{2}\cdot\log\tan\frac{\theta}{2}.$

(14) $\int_\theta \frac{1}{(\sin\theta)^2\cdot(\cos\theta)^3} = \left\{\frac{1}{2(\cos\theta)^2} - \frac{3}{2}\right\}\frac{1}{\sin\theta} + \frac{3}{2}\cdot\log\tan\left(\frac{\pi}{4} + \frac{\theta}{2}\right).$

(15) $\int_\theta \frac{1}{(\sin\theta)^4(\cos\theta)^2} = -\frac{1}{3\cos\theta(\sin\theta)^3} - \frac{8}{3}\cot 2\theta.$

(16) $\int_\theta (\tan\theta)^4 = \frac{1}{3}\tan^3\theta - \tan\theta + \theta.$

(17) $\int_\theta \frac{1}{(\tan\theta)^5} = \frac{-1}{4(\tan\theta)^4} + \frac{1}{2(\tan\theta)^2} + \log(\sin\theta).$

(18) $\int_\theta \theta^3\cdot\cos\theta = \theta^3\sin\theta + 3\theta^2\cos\theta - 6\theta\sin\theta - 6\cos\theta.$

(19) $\int_x \frac{x^2}{\sqrt{1-x^2}}\sin^{-1}x = \frac{(\sin^{-1}x)^2}{4} - \frac{x\sqrt{1-x^2}}{2}\sin^{-1}x + \frac{x^2}{4}.$

(20) $\int_x \frac{x}{(1-x^2)^{\frac{3}{2}}}\sin^{-1}x = \frac{\sin^{-1}x}{\sqrt{1-x^2}} + \log\sqrt{\frac{1-x}{1+x}}.$

(21) $\int_x \frac{x^2}{1+x^2} \tan^{-1}x = x\tan^{-1}x - \frac{1}{2}(\tan^{-1}x)^2 - \log\sqrt{1+x^2}.$

(22) $\int_x x^2 \tan^{-1}\sqrt{\frac{x}{a}} = \frac{x^3+a^3}{3}\tan^{-1}\sqrt{\frac{x}{a}} - \frac{\sqrt{ax}}{3}\left(\frac{x^2}{5} - \frac{ax}{3} + a^2\right).$

(23) $\int_x e^{ax}.\cos kx = \frac{e^{ax}(a\cos kx + k\sin kx)}{k^2+a^2}.$

(24) $\int_x e^{-ax}.\sin kx = -\frac{e^{-ax}.(a\sin kx + k\cos kx)}{k^2+a^2}.$

(25) $\int_x e^{ax}.(\sin x)^2 = \frac{e^{ax}.\sin x(a\sin x - 2\cos x)}{a^2+4} + \frac{2.e^{ax}}{a(a^2+4)}.$

(26) $\int_\theta \frac{1}{a\cos^2\theta + b\sin^2\theta} = \frac{1}{\sqrt{ab}}.\tan^{-1}\left(\sqrt{\frac{b}{a}}\tan\theta\right).$

(27) $\int_\theta \frac{\cos\theta}{(1-e^2\cos^2\theta)^{\frac{3}{2}}} = \frac{1}{1-e^2}.\frac{\sin\theta}{\sqrt{1-e^2\cos^2\theta}}.$

(28) $\int_x \frac{1}{(a+b\cos x)^2} = \frac{1}{a^2-b^2}\left\{\frac{-b\sin x}{a+b\cos x} + a\int_x \frac{1}{a+b\cos x}\right\}.$

(29) $\int_\theta (\cos 2\theta)^{\frac{3}{2}}\cos\theta = \frac{1}{8}\sin\theta(3+2\cos 2\theta)\sqrt{\cos 2\theta} + \frac{3}{8\sqrt{2}}\sin^{-1}(\sqrt{2}\sin\theta).$

(30) $\int_\theta \frac{\sin\theta}{\sqrt{\sin^2\alpha - \sin^2\theta}} = \log.\left(\frac{\cos\alpha}{\cos\theta + \sqrt{\sin^2\alpha - \sin^2\theta}}\right)$ from $\theta=\theta$ to $\theta=\alpha$.

(31) $\int_x e^{2A\cos x}$, (where $A = \log a$), between $x=0$, and $x=\pi$

$= 1 + A^2 + \left(\frac{A^2}{2}\right)^2 + \left(\frac{A^3}{2.3}\right)^3 + \&c.$

CHAPTER VI.

APPLICATION OF THE INTEGRAL CALCULUS TO DETERMINE THE AREAS AND LENGTHS OF PLANE CURVES, AND THE VOLUMES AND SURFACES OF SOLIDS OF REVOLUTION.

88. We have seen in the Differential Calculus, that if $y = f(x)$ be the equation to a curve, and A the area of a portion ANP, that $\frac{dA}{dx} = y = f(x)$. Hence, when the equation to a curve is given, its area may be found by finding the value of $\int_x f(x)$, and this integral may in general be found by means of the rules given in the preceding Chapters. If the equation to the curve be between polar co-ordinates, then

$$\frac{dA}{d\theta} = \frac{r^2}{2}; \quad \therefore A = \int_\theta \frac{r^2}{2}.$$

It is sometimes convenient to substitute z for $\phi(x)$; but then, since $y = f(z)$,

$$\frac{dA}{dz} = \frac{dA}{dx} \cdot \frac{dx}{dz} = y \frac{dx}{dz};$$

$$\therefore A = \int_z y \cdot \frac{dx}{dz} = \int_z f(z) \cdot \frac{dx}{dz}.$$

89. Again, if s represents the length of a curve, of which the equation is $y = f(x)$,

$$\text{since } \frac{ds}{dx} = \sqrt{1 + \frac{dy^2}{dx^2}}; \quad \therefore s = \int_x \sqrt{1 + \frac{dy^2}{dx^2}},$$

where $\frac{dy}{dx}$ may be found from $y = f(x)$.

90. Also, if V and S respectively represent the volume and surface of a solid of revolution, since

$$\frac{dV}{dx} = \pi y^2, \quad \text{and } \frac{dS}{dx} = 2\pi y \sqrt{1 + \frac{dy^2}{dx^2}};$$

$$\therefore V = \pi \int_x y^2, \quad \text{and } S = 2\pi \cdot \int_x y \sqrt{1 + \frac{dy^2}{dx^2}}.$$

91. A constant must be added to each of these integrals, the determination of which depends upon the nature of the particular problem.

As an illustration, let the area ABD be required, the nature of the curve ANP being known by the equation $y=f(x)$, where $AN=x$, and $NP=y$.



Let $AB=a$, and $ANP=A$;

$$\therefore \frac{dA}{dx}=y=f(x);$$

$$\therefore A=ANP=\int_a f(x)=\phi(x)+C.....(1).$$

To find C, we observe that if $x=0$ the area $=0$; if therefore at the same time $\phi(x)=0$; $\therefore C=0$,

and $ANP=\phi(x)$, and $ABD=\phi(a)$;

the same result as would have been obtained had we successively put $x=0$ and $x=a$ in equation (1), and subtracted the former result from the latter. This process is called integrating between the limits of $x=0$ and $x=a$, and is commonly represented by the symbol $\int_0^a f(x)$; the first limit being placed below, the second above the sign of integration.

To take a second instance, let the area $DBCE$ be required where $AC=b$; putting a for x in equation (1),

$$\text{area } ABD=\phi(a)+C,$$

$$\text{and area } ACE=\phi(b)+C;$$

$$\therefore \text{area } BDEC=\phi(b)-\phi(a).$$

Hence, if the value of an integral $u=\phi(x)$ be required between two values a and b of x, omit the constant, and having put a and b successively for x in $\phi(x)$, subtract $\phi(a)$ from $\phi(b)$. This is called integrating between the limits or values a and b of x, and the integral so found is called a *definite integral*, and is expressed by $\int_a^b f(x)$.

We have already found definite integrals in the preceding pages, and if we use the symbol mentioned above,

$$\int_0^1 \frac{x^{2n}}{\sqrt{1-x^2}}=\frac{1.3.5...(2n-3).(2n-1)}{2.4.6...2n-2\ .\ 2n}.\frac{\pi}{2}. \quad \text{Art. (49);}$$

$$\int_0^{2a} \frac{x^m}{\sqrt{2ax-x^2}}=\frac{1.3.5...(2m-3)\,(2m-1)}{2.4.6...(m-1)\ .\ m}.\pi a^m. \quad \text{Art. (53);}$$

$$\int_{-\infty}^{\infty} e^{-t^2}=\sqrt{\pi}. \quad \text{Art. (63), and some others}$$

When C is not determined, the value, $\phi(x)+C$, is called the *general* integral.

AREAS OF CURVES.

92. To find the areas of curves, or to integrate

$$\frac{dA}{dx} = y, \quad \text{or} \quad \frac{dA}{d\theta} = \frac{r^2}{2}.$$

Ex. 1. To find the area of the circle.



$$\left.\begin{matrix} CN = x \\ NP = y \\ CA = a \end{matrix}\right\}; \quad \therefore y = \sqrt{a^2 - x^2};$$

$$\therefore A = \int_x y = \int_x \sqrt{a^2 - x^2};$$

$$\therefore \text{area } CBPN = \int_x \sqrt{a^2 - x^2}.$$

But $$\int_x \sqrt{a^2 - x^2} = \int_x \frac{a^2 - x^2}{\sqrt{a^2 - x^2}}$$

$$= \frac{a^2}{2} \sin^{-1} \frac{x}{a} + \frac{x}{2} \sqrt{a^2 - x^2} + C.$$

$C = 0$, since area $= 0$; if $x = 0$;

$\sin^{-1}\frac{x}{a}$ can only be approximated to, by means of an infinite series, but if $x = a$, it $= \frac{\pi}{2}$, and,

$$\text{quadrant } ACB = \int_0^a \sqrt{a^2 - x^2} = \frac{a^2}{2} \cdot \frac{\pi}{2} = \frac{\pi a^2}{4};$$

$$\therefore \text{area of the circle} = \pi a^2.$$

Cor. 1. If $AN = x$, $y = \sqrt{2ax - x^2}$,

$$\therefore ANP = \int_0^x \sqrt{2ax - x^2},$$

and when $x = a$, ANP becomes a quadrant;

$$\therefore \int_x^a \sqrt{2ax - x^2} = \frac{\pi a^2}{4}.$$

The two definite integrals $\int_0^a \sqrt{a^2 - x^2}$, and $\int_0^a \sqrt{2ax - x^2}$, should be carefully remembered.

Also $\int_x \sqrt{a^2 - x^2}$ being $= CBPN$, where CN is sometimes called the cosine to radius CA; $\therefore \int_x \sqrt{a^2 - x^2}$ is called

a circular area of which cosine $= x$ and radius $= a$; and $\int_s \sqrt{2ax - x^2}$ in which $AN = x$, is called a circular area, of which ver. sine $= x$ and radius $= a$.

If AN = the diameter, the area ANP is a semicircle;

$$\therefore \int_0^{2a} \sqrt{2ax - x^2} = \frac{\pi a^2}{2}.$$

Also $\because \int_0^{-a} \sqrt{a^2 - x^2}$ = area of the second quadrant;

$$\therefore \int_{-a}^{a} \sqrt{a^2 - x^2} = \frac{\pi a^2}{2}.$$

Cor. 2. To find the area of the sector ACP.

Let A = area ACP; $\theta = \angle ACP$;

$$\therefore \frac{dA}{d\theta} = \tfrac{1}{2} r^2 = \tfrac{1}{2} a^2;$$

$$\therefore A = \frac{a^2\theta}{2} = a \times \frac{a\theta}{2} = \frac{\text{radius} \times \text{arc}}{2}.$$

Ex. 2. To find the area of an ellipse.

The centre the origin; $CN = x$; $NP = y$,

$$\therefore y = \frac{b}{a}\sqrt{a^2 - x^2};$$

$$\therefore A = \int_s y = \frac{b}{a} \cdot \int_s \sqrt{a^2 - x^2};$$

$$\text{elliptic quadrant} = \frac{b}{a}\int_0^a \sqrt{a^2 - x^2} = \frac{b}{a} \cdot \frac{\pi a^2}{4} = \frac{\pi ab}{4};$$

$$\therefore \text{area of ellipse} = \pi ab.$$

Had the vertex been the origin, and $AN = x$,

$$\text{elliptic quadrant} = \frac{b}{a}\int_0^a \sqrt{2ax - x^2} = \frac{b}{a}\frac{\pi a^2}{4} = \frac{\pi ab}{4}.$$

Ex. 3. To find the area of the common parabola.

$$y^2 = 4mx; \quad \therefore y = 2\sqrt{mx}.$$

$$A = \int_s y = 2\int_s \sqrt{mx} = 2\sqrt{m} \cdot \tfrac{2}{3} x^{\frac{3}{2}} + C;$$

$$\therefore \text{area} = \int_0^x y = \frac{4\sqrt{m}}{3} x^{\frac{3}{2}} = \tfrac{2}{3} 2\sqrt{mx} \cdot x = \tfrac{2}{3} yx$$

$$= \tfrac{2}{3} \text{ of circumscribing rectangle.}$$

Ex. 4. To find the area of the Witch.

$$y = \frac{2a}{x}\sqrt{2ax - x^2};$$

$$\therefore \text{ area} = \int_x y = 2a\int_x \frac{\sqrt{2ax - x^2}}{x} = 2a\int_x \frac{2a - x}{\sqrt{2ax - x^2}}$$

$$= 2a\left\{\int_x \frac{a - x}{\sqrt{2ax - x^2}} + a\int_x \frac{1}{\sqrt{2ax - x^2}}\right\}$$

$$= 2x\left\{\sqrt{2ax - x^2} + a \text{ ver-sin}^{-1}\frac{x}{a}\right\} + C.$$

And area = 0, if $x = 0$; $\therefore C = 0$;

$$\therefore \text{ area} = 2a\left\{\sqrt{2ax - x^2} + a \text{ ver-sin}^{-1}\frac{x}{a}\right\}.$$

Let $x = 2a$; $\therefore$ area $= 2a \times a \,.\, \pi = 2\pi a^2$.

Ex. 5. Find the area of the hyperbolic sector CAP.

Sector $CAP = \Delta CNP -$ area ANP.



Let $\left.\begin{matrix} CN = x \\ NP = y \\ CA = a \end{matrix}\right\}$; $\therefore y = \frac{b}{a}\sqrt{x^2 - a^2}$.

$$ANP = \int_x y = \frac{b}{a}\int_x \sqrt{x^2 - a^2} = \frac{b}{a}\cdot\int_x \frac{x^2 - a^2}{\sqrt{x^2 - a^2}}$$

$$= \frac{b}{a}\cdot\left\{\frac{x}{2}\sqrt{x^2 - a^2} - \frac{a^2}{2}\cdot\log\left(x + \sqrt{x^2 - a^2}\right)\right\} + C,$$

and $0 = -\frac{ba}{2}\cdot\log a + C$; $\because ANP = 0$; if $x = a$;

$$\therefore ANP = \frac{xy}{2} - \frac{ba}{2}\cdot\log\left(\frac{x + \sqrt{x^2 - a^2}}{a}\right)$$

$$= \Delta CNP - \frac{ba}{2}\cdot\log\left(\frac{x + \sqrt{x^2 - a^2}}{a}\right);$$

$$\therefore \text{ sector } CAP = \frac{ba}{2}\cdot\log\left(\frac{x}{a} + \frac{y}{b}\right).$$

Ex. 6. Find the area of the portion $PNMQ$, PQ being an arc of the rectangular hyperbola.



Here $yx = \frac{a^2}{2}$. Let $CN = \alpha$, and $CM = \beta$,

$$\int_x y = \frac{a^2}{2}\int_x \frac{1}{x} = \frac{a^2}{2}.\log x + C;$$

$$\therefore\ PQMN = \frac{a^2}{2}.(\log\beta - \log\alpha) = \frac{a^2}{2}.\log\left(\frac{\beta}{\alpha}\right).$$

Also $\because\ \frac{yx}{2} = \frac{a^2}{4};\quad \therefore\ \Delta\ CNP = \Delta\ CQM;$

$$\therefore\ \text{sector } CPQ = \text{area } PNMQ = \frac{a^2}{2}\log\left(\frac{\beta}{\alpha}\right).$$

Ex. 7. Find the area of the cissoid.

$$\text{Here } y^2 = \frac{x^3}{2a - x};\quad \therefore\ y = \frac{x^{\frac{3}{2}}}{\sqrt{2a - x}};$$

$$\therefore\ \text{area} = \int_x y = \int_x \frac{x^{\frac{3}{2}}}{\sqrt{2a - x}}$$

$$= -2\sqrt{2a - x}\,.\,x^{\frac{3}{2}} + 3\,.\int_x x^{\frac{1}{2}}\sqrt{2a - x}$$

$$= -2x\sqrt{2ax - x^2} + 3\,.\int_x \sqrt{2ax - x^2},$$

$$\text{area} = \int_0^{2a} y = 3\,.\frac{\pi a^2}{2} = \frac{3\pi}{2}a^2.$$

Ex. 8. Find the area of the cycloid.

Origin from the vertex, $\frac{dy}{dx} = \frac{\sqrt{2ax - x^2}}{x}$,

$$\text{area} = \int_x y = yx - \int_x x\frac{dy}{dx} = yx - \int_x \sqrt{2ax - x^2},$$

also $\because$ if $x = 0$, $y = 0$; if $x = 2a$, $y = \pi a$;

$$\therefore\ \int_0^{2a} y = \text{semicycloid} = 2\pi a^2 - \tfrac{1}{2}\pi a^2 = \frac{3\pi a^2}{2};$$

$$\therefore\ \text{cycloid} = 3\pi a^2 = 3\,.\ \text{area of generating circle.}$$

Ex. 9. Area of the conchoid.

$$\text{Here } xy = (a + x)\sqrt{b^2 - x^2},$$

$$\text{area} = ab \log . \left(\frac{x}{b + \sqrt{b^2 - x^2}}\right) + \left(a + \frac{x}{2}\right)\sqrt{b^2 - x^2} + \frac{b^2}{2}\sin^{-1}\left(\frac{x}{b}\right) + C.$$

Ex. 10. In the common parabola, to find the area ASP.



$SA = a$, $\angle ASP = \theta$, $SP = r$;

$$\therefore\ r = \frac{2a}{1 + \cos\theta} = \frac{a}{\cos^2\frac{\theta}{2}};$$

$$\therefore\ A = \tfrac{1}{2}\int_\theta r^2 = a^2 \cdot \int_\theta \frac{\frac{1}{2}}{\cos^4\frac{\theta}{2}} = a^2 \cdot \int_\theta \frac{\frac{1}{2}}{\cos^2\frac{\theta}{2}}\left(1 + \tan^2\frac{\theta}{2}\right)$$

$$= a^2\left\{\tan\frac{\theta}{2} + \frac{1}{3}\tan^3\frac{\theta}{2}\right\}.$$

Ex. 11. Find the area of the lemniscata.

$$\text{Here } r^2 = a^2\cos 2\theta;$$

$$\therefore\ \int_\theta \tfrac{1}{2}r^2 = \frac{a^2}{2} \cdot \int_\theta \cos 2\theta = \frac{a^2}{4}\sin 2\theta + C.$$

Area = 0, if $\theta = 0$; $\therefore\ C = 0$; and area $= \frac{a^2}{4}\sin 2\theta$.

Let $\theta = 45^0$; $\therefore$ $\frac{1}{4}$th of lemniscata $= \frac{a^2}{4}$; lemniscata $= a^2$.

Ex. 12. Find the area of the spiral where $r = a\theta^n$.

$$\text{Here } \frac{dA}{d\theta} = \tfrac{1}{2}r^2;\ \theta = \left(\frac{r}{a}\right)^{\frac{1}{n}};$$

$$\therefore\ \frac{d\theta}{dr} = \frac{1}{na^{\frac{1}{n}}}r^{\frac{1}{n}-1};\ \therefore\ \frac{dA}{dr} = \frac{1}{2na^{\frac{1}{n}}}r^{1+\frac{1}{n}};$$

$$\therefore\ A = \frac{1}{2na^{\frac{1}{n}}} \cdot \frac{n}{2n+1}r^{\frac{2n+1}{n}} + C,$$

and $C = 0$, if $A = 0$, when $r = 0$.

Cor. Let $n = 1$, or the spiral be that of Archimedes;

$$\therefore \text{ area} = \frac{r^3}{6a} = \frac{\pi r^3}{3R},$$

$$\text{if } R = r \text{ when } \theta = 2\pi;$$

$$\text{therefore area of spiral in first revolution} = \frac{\pi R^2}{3}.$$

The area after two revolutions of the radius vector is when $\theta = 4\pi$, or when $r = 2R$. But before $r = 2R$, it will have made two revolutions, and therefore have twice generated the area from $r = 0$ to $r = R$.

Consequently we must subtract the area described in the first revolution from that in the second;

$$\therefore \text{ area} = \frac{\pi \cdot (2R)^3}{3R} - \frac{\pi R^2}{3} = \frac{7\pi R^2}{3}.$$

And the space between the arcs of the first and second areas

$$= \frac{7\pi R^2}{3} - \frac{\pi R^2}{3} = 2\pi R^2.$$

$$\text{At the } n^{\text{th}} \text{ revolution } r = nR,$$

$$\ldots\ldots(n-1)^{\text{th}} \ldots\ldots r = (n-1)R;$$

$$\therefore \text{ area after } n \text{ revolutions} = \frac{\pi}{3} \cdot \frac{(nR)^3 - (n-1)^3 R^3}{R}$$

$$= \frac{\pi R^2}{3}\{n^3 - (n-1)^3\}.$$

$$\text{Area after } (n+1) \text{ revolutions} = \frac{\pi R^2}{3}\{(n+1)^3 - n^3\};$$

$\therefore$ space between the arcs after $n+1$ and n revolutions

$$= \frac{\pi R^2}{3} \cdot \{(n+1)^3 + (n-1)^3 - 2n^3\} = \frac{\pi R^2}{3} \cdot 6n = 2n\pi R^2$$

$= n$ times the space between the first and second.

Ex. 13. Find the area of the involute of the circle, where $r^2 - p^2 = a^2$.

$$\frac{dA}{d\theta} = \tfrac{1}{2} r^2 \frac{d\theta}{dr} = \frac{p}{r\sqrt{r^2 - p^2}};$$

$$\therefore \frac{dA}{dr} = \frac{pr}{2\sqrt{r^2 - p^2}} = \frac{r\sqrt{r^2 - a^2}}{2a};$$

$$\therefore A = \frac{(r^2 - a^2)^{\frac{3}{2}}}{6a} + C; \text{ and } C = 0.$$

If $p = 2\pi a$; $A = \frac{4}{3}\pi^3 a^2$, or subtracting πa^2, the area of the involute exterior of the circle after one unwrapping of the string $= \pi a^2 \left(\frac{4\pi^2}{3} - 1\right)$.

Ex. 14. Find the area of the curve of which the equation is

$$y^3 - 3axy + x^3 = 0.$$

If the curve be traced there will be found a nodus as $APMQ$, to which the axes Ay and Ax are tangents.



Let $y = xz$; $\therefore z = \frac{y}{x} = \tan PAN$;

$$\therefore x^3 z^3 - 3ax^2 z + x^3 = 0;$$

$$\therefore x = \frac{3az}{1 + z^3}, \text{ and } y = \frac{3az^2}{1 + z^3};$$

and since x is $= 0$, for each of the branches APM and AQM, this will happen if $z = \infty$ or $= 0$.

$$\text{Now } \frac{dA}{dz} = \frac{dA}{dx} \cdot \frac{dx}{dz} = y \cdot \frac{dx}{dz},$$

$$\text{and } \frac{dx}{dz} = \frac{3a \cdot \{1 + z^3 - 3z^3\}}{(1 + z^3)^2} = 3a \frac{1 - 2z^3}{(1 + z^3)^2};$$

$$\therefore A = \int_z y \cdot \frac{dx}{dz} = 9a^2 \cdot \int_z \frac{z^2 (1 - 2z^3)}{(1 + z^3)^3}$$

$$= 9a^2 \int_z \left\{ \frac{z^2 [1 - 2(z^3 + 1) + 2]}{(1 + z^3)^3} \right\}$$

$$= 9a^2 \int_x \left\{ \frac{3x^2}{(1+x^3)^2} - \int_x \frac{2x^2}{(1+x^3)^3} \right\}$$

$$= 9a^2 \left\{ -\tfrac{1}{2} \cdot \frac{1}{(1+x^3)^2} + \frac{2}{3} \cdot \frac{1}{1+x^3} \right\} + C.$$

Let $x = 0$; $\therefore C = -\dfrac{3a^2}{2}$, and let $x_1 = \dfrac{y}{x}$ at M;

$$\therefore \text{ area } AQMm = -\frac{3a^2}{2} + 9a^2 \left\{ -\tfrac{1}{2} \cdot \frac{1}{(1+x_1^3)^2} + \frac{2}{3} \frac{1}{1+x_1^3} \right\}.$$

Integrating $x = \infty$ and $x = x_1$ for the branch APM,

$$\text{area } APMm = 9a^2 \left\{ -\tfrac{1}{2} \frac{1}{(1+x_1^3)^2} + \frac{2}{3} \cdot \frac{1}{1+x_1^3} \right\};$$

$$\therefore \text{ the nodus } APMQ = \text{area } APMm - \text{area } AQMm = \frac{3a^2}{2}.$$

If the area of the nodus only be required, the following method is often useful;

Since $\tan\theta = \dfrac{y}{x} = x$; $\therefore \dfrac{d\theta}{dx} = \cos^2\theta$;

$$\therefore \frac{dA}{dx} = \tfrac{1}{2} r^2 \frac{d\theta}{dx} = \tfrac{1}{2} r^2 \cos^2\theta = \frac{x^2}{2}.$$

and $\therefore A = \tfrac{1}{2} \int_x x^2$.

Here $x^2 = \dfrac{9a^2x^2}{(1+x^3)^2}$;

$$\therefore A = \frac{9a^2}{2} \cdot \int_x \frac{x^2}{(1+x^3)^2} = C - \frac{3a^2}{2} \cdot \frac{1}{1+x^3},$$

integrating from $x = 0$, to $x = \infty$: area $= \dfrac{3a^2}{2}$.

Ex. 15. To find the area of the evolute of an ellipse.

Here $\left(\dfrac{x}{\alpha}\right)^{\frac{2}{3}} + \left(\dfrac{y}{\beta}\right)^{\frac{2}{3}} = 1.$

We might put $y = xx$; but better thus:

Let $\dfrac{x}{\alpha} = \cos^3\theta$; $\therefore \dfrac{y}{\beta} = \sin^3\theta$; $\dfrac{dx}{d\theta} = -3\alpha \sin\theta \cdot \cos^2\theta$;

$$\therefore \frac{dA}{d\theta} = \frac{dA}{dx} \cdot \frac{dx}{d\theta} = y \frac{dx}{d\theta} = -3\alpha\beta \sin^4\theta \cos^2\theta;$$

$\therefore A = 3a\beta . \int_0 \sin^4\theta\cos^2\theta$; the limits of which are $\theta = \frac{\pi}{2}$ and $\theta = 0$, since those of x are, 0 and a.

$$\text{Now } \int_0 \sin^4\theta\cos^2\theta = \frac{\cos\theta\sin^5\theta}{5} + \frac{1}{5}\int_0 \sin^6\theta;$$

$$\text{but } \frac{\cos\theta . \sin^5\theta}{5} = 0, \text{ both when } \theta = 0 \text{ and } \theta = \frac{\pi}{2};$$

$$\therefore \int_{\frac{\pi}{2}}^{0} \sin^4\theta\cos^2\theta = -1\int_{\frac{\pi}{2}}^{0} \sin^6\theta = -\frac{1}{5}\times\frac{5.3.1}{6.4.2}.\frac{\pi}{2};$$

$$\therefore A = 3a\beta . \frac{3\pi}{6\times 4\times 4} = \frac{3a\beta}{8}.\frac{\pi}{4};$$

$$\therefore 4A = \text{whole area} = \frac{3\pi a\beta}{8} = \frac{3}{8}\pi\frac{(a^2-b^2)^2}{ab}.$$

Cor. For the evolute of the hyperbola, where $\left(\frac{x}{a}\right)^{\frac{2}{3}} - \left(\frac{y}{\beta}\right)^{\frac{2}{3}} = 1$, make $\frac{x}{a} = \sec^3\theta$; $\therefore \frac{y}{\beta} = \tan^3\theta$; $\therefore \frac{dx}{d\theta} = 3a\sec^3\theta . \tan\theta$;

$$\therefore A = 3a\beta . \int_0 \sec^3\theta . \tan^4\theta = 3a\beta . \int \frac{\sin^4\theta}{\cos^7\theta};$$

the limits of θ being 0, and some finite value θ'.

Ex. 16. The same substitution applies to find the area of

$$\left(\frac{y}{\beta}\right)^{\frac{2}{2n+1}} + \left(\frac{x}{a}\right)^{\frac{2}{2n+1}} = 1.$$

For making $x = a\cos^{2n+1}\theta$; $\therefore y = \beta\sin^{2n+1}\theta$,

$$\frac{dx}{d\theta} = -(2n+1)a\cos^{2n}\theta\sin\theta;$$

$$\therefore A = -(2n+1)a\beta\int_{\frac{\pi}{2}}^{0} \sin^{2n+2}\theta\cos^{2n}\theta$$

$$= -(2n+1)a\beta\int_{\frac{\pi}{2}}^{0} \{\sin^{2n+2}\theta(1-\sin^2\theta)^n\};$$

whence expanding, $\because \int_{\frac{\pi}{2}}^{0} \sin^{2n}\theta = -\frac{(2n-1).(2n-3)\ldots 3.1}{2n.(2n-2)\ldots 4.2}.\frac{\pi}{2}$;

$$\therefore \text{area} = 4A = (4n+2)\pi a\beta.\left\{\frac{1.3.5\ldots(2n+1)}{2.4.6\ldots(2n+2)} - n.\frac{1.3\ldots(2n+3)}{2.4\ldots(2n+4)}\right.$$

$$\left. +n.\frac{(n-1)}{2}.\frac{1.3\ldots(2n+5)}{2.4\ldots(2n+6)} - n\frac{(n-1)(n-2)}{2.3}.\frac{1.3\ldots(2n+7)}{2.4\ldots(2n+8)} + \&c.\right\}.$$

If $n = 1$, $\left(\frac{y}{\beta}\right)^{\frac{1}{3}} + \left(\frac{x}{\alpha}\right)^{\frac{1}{3}} = 1$, and area $= 6\pi\alpha\beta\left\{\frac{3}{8} - \frac{3 \cdot 5}{8 \cdot 6}\right\}$

$= \frac{3}{8}\pi\alpha\beta$, the result obtained in the preceding example.

THE LENGTHS OF CURVES.

93. To find the lengths of curves, or to integrate

$$\frac{ds}{dx} = \sqrt{1 + \frac{dy^2}{dx^2}}, \text{ when } y = f(x).$$

Ex. 17. Find the length of an arc of the parabola.

$$y^2 = 4mx; \quad \therefore \frac{dy}{dx} = \frac{2m}{y}; \quad \therefore \frac{dy^2}{dx^2} = \frac{4m^2}{y^2} = \frac{m}{x};$$

$$s = \int_x \sqrt{1 + \frac{dy^2}{dx^2}} = \int_x \sqrt{1 + \frac{m}{x}} = \int_x \frac{\sqrt{x+m}}{\sqrt{x}}$$

$$= \int_x \frac{x+m}{\sqrt{x^2+mx}} = \int_x \frac{x + \frac{m}{2}}{\sqrt{x^2+mx}} + \int_x \frac{\frac{m}{2}}{\sqrt{x^2+mx}}$$

$$= \sqrt{x^2+mx} + \frac{m}{2}\log\left(x + \frac{m}{2} + \sqrt{x^2+mx}\right) + C.$$

And $s = 0$, if $x = 0$; $\quad \therefore 0 = \frac{m}{2}\log\left(\frac{m}{2}\right) + C$;

$$\therefore s = \sqrt{x^2+mx} + \frac{m}{2}\log\left(\frac{2x + m + 2\sqrt{x^2+mx}}{m}\right).$$

Ex. 18. Find when curves included under the general equation $y = ax^{\frac{m}{n}}$ are rectifiable.

$$\frac{dy}{dx} = \frac{m}{n}ax^{\frac{m-n}{n}}; \quad \therefore s = \int_x \sqrt{1 + \frac{m^2a^2}{n^2} \cdot x^{\frac{2m-2n}{n}}};$$

which is integrable,

(1) When $\frac{n}{2m-2n}$ is an integer $= r$,

$$\text{or } \frac{m}{n} - 1 = \frac{1}{2r}, \text{ or } \frac{m}{n} = \frac{1}{2r} + 1 = \frac{2r+1}{2r}.$$

(2) When $\frac{n}{2m-2n}+\frac{1}{2}=$ an integer $=q$,

$$\text{or } \frac{m}{n}-1=\frac{1}{2q-1}, \text{ or } \frac{m}{n}=\frac{2q}{2q-1}.$$

Let $r=1, 2, 3,$ &c. $q=1, 2, 3,$ &c.;

$$\therefore \frac{m}{n}=\frac{3}{2}, \frac{5}{4}, \frac{7}{6}, \text{ &c. and } \frac{m}{n}=\frac{2}{1}, \frac{4}{3}, \frac{6}{5}, \text{ &c.}$$

Ex. 19. Let $\frac{m}{n}=\frac{3}{2}$, or the curve be the semi-cubical parabola;

$$\therefore y=ax^{\frac{3}{2}}; \frac{dy}{dx}=\frac{3a}{2}x^{\frac{1}{2}}=\frac{\sqrt{x}}{\sqrt{c}}, \text{ if } \sqrt{c}=\frac{2}{3a};$$

$$\therefore s=\int_x\sqrt{1+\frac{x}{c}}=\frac{1}{\sqrt{c}}\cdot\int_x\sqrt{x+c}=\frac{1}{\sqrt{c}}\frac{2}{3}(x+c)^{\frac{3}{2}}+C.$$

$$\text{If } s=0, x=0; \therefore C=-\frac{1}{\sqrt{c}}\frac{2}{3}c^{\frac{3}{2}};$$

$$\therefore s=\frac{1}{\sqrt{c}}\cdot\frac{2}{3}\cdot\left\{(x+c)^{\frac{3}{2}}-c^{\frac{3}{2}}\right\}.$$

Ex. 20. Find the length of the cycloid.

$$\frac{dy}{dx}=\sqrt{\frac{2a-x}{x}}; \quad \therefore 1+\frac{dy^2}{dx^2}=1+\frac{2a-x}{x}=\frac{2a}{x};$$

$$\therefore s=\int_x\sqrt{\frac{2a}{x}}=\sqrt{2a}\cdot\int_x\frac{1}{\sqrt{x}}=\sqrt{2ax}+C,$$

and $C=0$, since $s=0$ when $x=0$;

therefore $s=2\sqrt{2ax}=$ twice the chord of the arc of the generating circle, corresponding to the arc of the cycloid.

Hence the cycloid is rectifiable.

If $x=2a$, $s=2\sqrt{4a^2}=4a$, or the length of the semi-cycloid = twice the diameter of the circle.

Ex. 21. Find the length of the arc of an ellipse.

$$y=\frac{b}{a}\sqrt{a^2-x^2}; \frac{dy}{dx}=-\frac{b}{a}\cdot\frac{x}{\sqrt{a^2-x^2}};$$

$$\therefore 1+\frac{dy^2}{dx^2}=1+\frac{b^2x^2}{a^2(a^2-x^2)}=\frac{a^4-(a^2-b^2)x^2}{a^2(a^2-x^2)}=\frac{a^2-e^2x^2}{a^2-x^2};$$

$$\therefore\ s = \int_x \frac{\sqrt{a^2 - e^2x^2}}{\sqrt{a^2 - x^2}} = a\int_x \frac{\sqrt{1 - e^2x^2}}{\sqrt{1 - x^2}},\ \text{(if } x = ax\text{)}$$

$$= a\int_x \frac{1}{\sqrt{1 - x^2}} \cdot \left\{1 - \tfrac{1}{2}e^2x^2 - \frac{1.1}{2.4}e^4x^4 - \frac{1.1.3}{2.4.6}\cdot e^6x^6 - \&c.\right\},$$

the integration of which depends on $\int_x \frac{x^{2n}}{\sqrt{1 - x^2}}$.

If the quadrant be required, we must integrate from $x = 0$ to $x = a$, or from $x = 0$ to $x = 1$, but then

$$\int_0^1 \frac{x^{2n}}{\sqrt{1 - x^2}} = \frac{\pi}{2}\cdot\frac{1.3.5.....(2n-1)}{2.4.6.....2n};$$

$$\therefore \int_0^1 \frac{x^2}{\sqrt{1 - x^2}} = \frac{\pi}{2}\cdot\frac{1}{2};\quad \int_0^1 \frac{x^4}{\sqrt{1 - x^2}} = \frac{\pi}{2}\cdot\frac{1.3}{2.4};$$

$$\int_0^1 \frac{x^6}{\sqrt{1 - x^2}} = \frac{\pi}{2}\cdot\frac{1.3.5}{2.4.6},\ \&c.;\quad \int_0^1 \frac{1}{\sqrt{1 - x^2}} = \frac{\pi}{2};$$

therefore elliptic quadrant

$$= \frac{\pi a}{2}\cdot\left\{1 - \frac{1}{2^2}e^2 - \frac{1.3}{2^2.4^2}e^4 - \frac{1.3^2.5}{2^2.4^2.6^2}e^6 - \&c.\right\},$$

a series rapidly convergent when e is a small fraction.

(3) To find the same by circular functions.

Let $x = a\cos\theta$; $\therefore\ y = b\sin\theta$, $\because\ x$ is $< a$;

$$\therefore \frac{ds}{d\theta} = \sqrt{a^2\sin^2\theta + b^2\cos^2\theta}$$

$$= \sqrt{a^2 - (a^2 - b^2)\cos^2\theta} = a\sqrt{1 - e^2\cos^2\theta}$$

$$= a\left\{1 - \tfrac{1}{2}e^2\cos^2\theta - \frac{1.1}{2.4}e^4\cos^4\theta - \frac{1.1.3}{2.4.6}e^6\cos^6\theta - \&c.\right\}.$$

Now $\int_\theta \cos^{2n}\theta = \sin\theta\,.\,\cos^{2n-1}\theta + (2n-1)\,.\,\int_\theta \cos^{2n-2}\theta\,.\,\sin^2\theta$

$$= \frac{\sin\theta\cos^{2n-1}\theta}{2n} + \frac{2n-1}{2n}\cdot\int_\theta \cos^{2n-2}\theta,$$

and $\sin\theta\cos^{2n-1}\theta = 0$, when $\theta = 0$, and $\theta = \frac{\pi}{2}$;

$\therefore$ calling $\int_0^{\frac{\pi}{2}} \cos^{2n}\theta = P_{2n}$,

$$P_{2n} = \frac{2n-1}{2n} \cdot P_{2n-2} = \frac{(2n-1)(2n-3)}{2n(2n-2)} P_{2n-4}$$

$$= \frac{(2n-1) \cdot (2n-3) \ldots\ldots 3 \cdot 1}{2n \cdot (2n-2) \ldots\ldots 4 \cdot 2} \cdot \frac{\pi}{2}; \quad \because P_0 = \frac{\pi}{2}.$$

$$\therefore P_2 = \tfrac{1}{2} \cdot \frac{\pi}{2}; \qquad P_4 = \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{\pi}{2}, \quad P_6 = \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \cdot \frac{\pi}{2} \text{ &c.};$$

$$\therefore s = \frac{\pi a}{2} \cdot \left\{1 - \frac{1}{2^2}e^2 - \frac{1 \cdot 3}{2^2 \cdot 4^2}e^4 - \frac{1 \cdot 3^2 \cdot 5}{2^2 \cdot 4^2 \cdot 6^2} \cdot e^6 - \frac{1 \cdot 3^2 \cdot 5^2 \cdot 7}{2^2 \cdot 4^2 \cdot 6^2 \cdot 8^2}e^8 - \text{&c.}\right\}.$$

Ex. 22. Find the length of a hyperbolic arc.

$$y = \frac{b}{a}\sqrt{x^2 - a^2}, \quad \frac{dy}{dx} = \frac{b}{a} \cdot \frac{x}{\sqrt{x^2 - a^2}};$$

$$\therefore \frac{ds}{dx} = \sqrt{1 + \frac{dy^2}{dx^2}} = \sqrt{\frac{(b^2 + a^2)x^2 - a^4}{a^2(x^2 - a^2)}} = \sqrt{\frac{e^2x^2 - a^2}{x^2 - a^2}};$$

$$\therefore s = \int_x \sqrt{\frac{e^2x^2 - a^2}{x^2 - a^2}} = a \cdot \int_z \sqrt{\frac{e^2z^2 - 1}{z^2 - 1}}, \text{ (if } x = az\text{)},$$

$$= ae \int_z z \frac{\sqrt{1 - \frac{1}{e^2z^2}}}{\sqrt{z^2 - 1}}$$

$$= a \int_z \frac{ez}{\sqrt{z^2 - 1}} \cdot \left\{1 - \tfrac{1}{2}\frac{1}{(ez)^2} - \frac{1 \cdot 1}{2 \cdot 4} \cdot \frac{1}{(ez)^4} - \frac{1 \cdot 1 \cdot 3}{2 \cdot 4 \cdot 6} \cdot \frac{1}{(ez)^6}\right.$$

$$\left. - \frac{1 \cdot 1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6 \cdot 8} \cdot \frac{1}{(ez)^8} - \text{&c.}\right\},$$

the limits being $x = a$; $x = \infty$, or $z = 1$; $z = \infty$;

now every term, except the first, depends upon

$$\int_z \frac{1}{z^m\sqrt{z^2 - 1}}, \quad m \text{ odd},$$

and $\displaystyle\int \frac{1}{z^m\sqrt{z^2 - 1}} = \frac{1}{m-1} \cdot \frac{\sqrt{z^2 - 1}}{z^{m-1}} + \frac{m-2}{m-1} \cdot \int_z \frac{1}{z^{m-2}\sqrt{z^2 - 1}}$,

and $\dfrac{\sqrt{z^2 - 1}}{z^{m-1}}$ vanishes both when $z = 1$, and $z = \infty$;

$$\therefore \int_1^\infty \frac{1}{x^m\sqrt{x^2-1}} = \frac{m-2}{m-1}\int_1^\infty \frac{1}{x^{m-2}\sqrt{x^2-1}}.$$

But $\int_1^\infty \frac{1}{x\sqrt{x^2-1}} = \sec^{-1}x = \frac{\pi}{2}$;

$$\therefore \int_1^\infty \frac{1}{x^3\sqrt{x^2-1}} = \tfrac{1}{2}\int_1^\infty \frac{1}{x\sqrt{x^2-1}} = \tfrac{1}{2}\cdot\frac{\pi}{2};$$

$$\int_1^\infty \frac{1}{x^5\sqrt{x^2-1}} = \tfrac{3}{4}\cdot\int_1^\infty \frac{1}{x^3\sqrt{x^2-1}} = \frac{1.3}{2.4}\cdot\frac{\pi}{2};$$

$$\int_1^\infty \frac{1}{x^7\sqrt{x^2-1}} = \tfrac{5}{6}\cdot\int_1^\infty \frac{1}{x^5\sqrt{x^2-1}} = \frac{1.3.5}{2.4.6}\cdot\frac{\pi}{2};$$

$$\therefore s = ae\cdot\int_z \frac{x}{\sqrt{x^2-1}} - \frac{\pi a}{2}\cdot\left\{\tfrac{1}{2}\cdot\frac{1}{e} + \frac{1.1}{2^2.4}\cdot\frac{1}{e^3}\right.$$

$$\left. + \frac{1.1.3^2}{2^2.4^2.6}\cdot\frac{1}{e^5} + \frac{1.1.3^2.5^2}{2^2.4^2.6^2.8}\cdot\frac{1}{e^7} + \&c.\right\}.$$

Now the equation to the asymptote is $y = \frac{bx}{a}$;

$$\therefore \text{length of asymptote} = \sqrt{x^2 + \frac{b^2x^2}{a^2}} = x\sqrt{\frac{a^2+b^2}{a^2}} = ex = aez.$$

But $ae\int_z \frac{x}{\sqrt{x^2-1}} = ae\sqrt{x^2-1} = aez$ from $x=1$ to $x=\infty$.

If therefore l be the length of the asymptote,

$$l - s = \frac{\pi a}{2}\left\{\tfrac{1}{2}\cdot\frac{1}{e} + \frac{1.1}{2^2.4}\cdot\frac{1}{e^3} + \frac{1.1.3^2}{2^2.4^2.6}\cdot\frac{1}{e^5}\right.$$

$$\left. + \frac{1.1.3^2.5^2}{2^2.4^2.6^2.8}\cdot\frac{1}{e^7} + \&c.\right\}.$$

Ex. 23. Find the length of an arc of the logarithmic curve.

Here $y = a^x$, and $\frac{dy}{dx} = Aa^x = A.y$,

and $\frac{ds}{dy} = \frac{ds}{dx}\cdot\frac{dx}{dy} = \sqrt{1+A^2y^2}\cdot\frac{1}{Ay}$;

$$\therefore s = \int_y \frac{\sqrt{1 + A^2y^2}}{Ay} = \int_y \frac{Ay}{\sqrt{1 + A^2y^2}} + \int_y \frac{1}{Ay\sqrt{1 + A^2y^2}}$$

$$= \frac{\sqrt{1 + A^2y^2}}{A} + \frac{1}{A} \log \frac{Ay}{1 + \sqrt{1 + A^2y^2}} + C.$$

Ex. 24. Find the length of an arc of the Lemniscata.

$$r^2 = a^2 \cos 2\theta, \quad \text{and} \quad s = \int_r \sqrt{1 + r^2 \frac{d\theta^2}{dr^2}};$$

$$\therefore s = \int_r \frac{a^2}{\sqrt{a^4 - r^4}} = a \cdot \int_x \frac{1}{\sqrt{1 - x^4}}, \quad \text{if } r = ax$$

$$= a \int_x \cdot \left(\frac{1}{\sqrt{1 - x^2}} \cdot \frac{1}{\sqrt{1 + x^2}} \right)$$

$$= a \int_x \cdot \frac{1}{\sqrt{1 - x^2}} \cdot \left\{ 1 - \tfrac{1}{2}x^2 + \frac{1.3}{2.4}x^4 - \frac{1.3.5}{2.4.6}x^6 + \&c. \right\}.$$

Integrate from $\theta = 45^0$ to $\theta = 0$; or from $x = 0$ to $x = 1$;

$$\therefore \int_0^1 \frac{1}{\sqrt{1 - x^2}} = \frac{\pi}{2}, \quad \int_0^1 \frac{x^2}{\sqrt{1 - x^2}} = \tfrac{1}{2} \cdot \frac{\pi}{2};$$

$$\int_0^1 \frac{x^4}{\sqrt{1 - x^2}} = \frac{1.3}{2.4} \cdot \frac{\pi}{2}, \quad \text{and} \int_0^1 \frac{x^6}{\sqrt{1 - x^2}} = \frac{1.3.5}{2.4.6} \cdot \frac{\pi}{2};$$

$$\therefore s = \frac{\pi a}{2} \cdot \left\{ 1 - \frac{1}{2^2} + \frac{1^2.3^2}{2^2.4^2} - \frac{1^2.3^2.5^2}{2^2.4^2.6^2} + \frac{1^2.3^2.5^2.7^2}{2^2.4^2.6^2.8^2} - \&c. \right\}.$$

The whole length of the lemniscata $= 4s$.

Ex. 25. To find the length of the involute of a circle.

Here $r^2 - p^2 = a^2$; $\quad \frac{ds}{dr} = \frac{r}{\sqrt{r^2 - p^2}} = \frac{r}{a}$;

$$\therefore s = \frac{r^2}{2a} + C = \frac{r^2 - a^2}{2a} = \frac{p^2}{2a} = \frac{a}{2} \cdot \theta^2.$$

If $\theta = 2\pi$, or the string be unwound once, $s = 2\pi^2 a$.

If $\theta = 2n\pi$, or the string be unwound n times length $= \frac{a}{2}(2n\pi)^2$.

Ex. 26. If the radius of the circle be unity, and θ be a

circular arc, θ_1 the length of its involute, θ_2 the length of the involute of θ_1, &c., prove that

$$\theta + \theta_1 + \theta_2 + \&c. = e^\theta - 1.$$



$$AQ = \theta, \qquad QP = \rho,$$
$$AP = \theta_1, \qquad PP_1 = \rho_1,$$
$$AP_1 = \theta_2, \qquad P_1P_2 = \rho_2,$$
$$AP_3 = \theta_3.$$

Then $\angle PQp = \angle P_1Pp_1$; $\therefore \dfrac{Pp}{QP} = \dfrac{P_1p_1}{PP_1}$;

$$\therefore \frac{d\theta_1}{QP} \text{ or } \frac{d\theta_1}{\theta} = \frac{d\theta_2}{\theta_1}; \text{ but } \theta_1 = \tfrac{1}{2}\theta^2;$$

$$\therefore d\theta_2 = \frac{\theta_2}{\theta} d\theta_1 = \tfrac{1}{2}\theta^2 d\theta; \quad \therefore \theta_2 = \frac{1}{2.3}\theta^3.$$

$$\text{And } d\theta_3 = \frac{\theta_2}{\theta_1} d\theta_2 = \theta_2 . \frac{d\theta_1}{\theta} = \frac{1}{2.3} . \theta^3 . d\theta;$$

$$\therefore \theta_3 = \frac{1}{2.3.4}\theta^4;$$

$$\therefore \theta + \theta_1 + \theta_2 + \&c. = \theta + \tfrac{1}{2}\theta^2 + \frac{1}{2.3}\theta^3 + \&c. = e^\theta - 1.$$

THE VOLUMES AND SURFACES OF SOLIDS OF REVOLUTION.

94. To find the volumes and surfaces of solids, or to integrate the functions

$$\frac{dV}{dx} = \pi y^2, \text{ and } \frac{dS}{dx} = 2\pi y \sqrt{1 + \frac{dy^2}{dx^2}}.$$

Ex. 27. To find the volume of a cone with a circular base.

Let a = altitude, b = radius of base.

Then if the vertex be the origin and the altitude the axis of x, $y = \frac{b}{a}x$;

$$\therefore\ V = \pi \int_x y^2 = \frac{\pi b^2}{a^2} \int_x x^2 = \frac{\pi b^2}{a^2} \cdot \frac{x^3}{3} + C.$$

And $V = 0$ if $x = 0$; $\therefore\ C = 0$; $\therefore\ V = \frac{\pi b^2}{a^2} \cdot \frac{x^3}{3}$.

Let $x = a$; $\therefore$ whole cone $= \frac{\pi b^2 a}{3} = \frac{1}{3}$ of a cylinder of the same altitude and on the same base.

Ex. 28. Find the volume of the paraboloid.

$y^2 = 4mx$ is the equation to the generating curve;

$\therefore\ V = \pi \int_x y^2 = \pi \int_x 4mx = 2\pi m \cdot x^2 + C$, and $C = 0$;

$$\therefore\ V = 2\pi m x^2 = \frac{\pi 4mx \cdot x}{2} = \frac{\pi y^2 x}{2}.$$

But $\pi y^2 x$ = volume of a cylinder, base $= \pi y^2$ and altitude $= x$;

$\therefore$ paraboloid $= \frac{1}{2}$ circumscribing cylinder.

Ex. 29. Find the volume of a sphere.

Here $y^2 = 2ax - x^2$;

$$\therefore\ V = \pi \int_x \cdot (2ax - x^2) = \pi \left(ax^2 - \frac{x^3}{3} \right) + C,$$

and $V = 0$ if $x = 0$; $\therefore\ C = 0$; $\therefore\ V = \pi x^2 \left\{ a - \frac{x}{3} \right\}$.

Let $x = 2a$; $\therefore$ sphere $= 4\pi a^2 \left(a - \frac{2}{3}a\right) = \frac{4}{3}\pi a^3$.

Since circumscribing cylinder $= 2a \cdot \pi a^2 = 2\pi a^3$;

$\therefore$ sphere $= \frac{2}{3}$ of circumscribing cylinder.

Ex. 30. Find the volume of the prolate spheroid formed by the revolution of an ellipse round its major axis.

$$y^2 = \frac{b^2}{a^2}(a^2 - x^2) \ldots\ldots (1);$$

$$\therefore\ V = \pi \int_{\frac{b^2}{a^2}} (a^2 - x^2) = \pi \frac{b^2}{a^2}\left(a^2 x - \frac{x^3}{3}\right) + C$$

$$= \frac{4}{3}\pi b^2 a;\quad \text{from } x = -a,\ \text{to } x = +a.$$

If the solid content of the oblate spheroid, which is formed by revolution round the minor axis be required; take the minor axis for the axis of x, and the major for that of y.

Then in equation (1) put y for x and x for y, we have

$$x^2 = \frac{b^2}{a^2}(a^2 - y^2);\quad \therefore\ y^2 = \frac{a^2}{b^2}(b^2 - x^2);$$

$$\therefore\ \text{solid} = \pi \cdot \int_{\frac{a^2}{b^2}} (b^2 - x^2) = \frac{\pi a^2}{b^2}\left(b^2 x - \frac{x^3}{3}\right)$$

$$= \frac{4}{3}\pi a^2 b;\quad \text{from } x = -b,\ \text{to } x = +b;$$

$\therefore$ prolate spheroid : oblate spheroid :: b : a.

Cor. Sphere on major axis : prolate spheroid :: $a^2 : b^2$,

sphere on minor axis : oblate spheroid :: $b^2 : a^2$.

Ex. 31. Find the solid generated by the conchoid round the asymptote.

Here $xy = (a + x)\sqrt{b^2 - x^2}$;

and since the curve revolves round the axis of y;

$$\therefore\ dV = \pi x^2 dy = -\pi \frac{ab^2 + x^3}{\sqrt{b^2 - x^2}} \cdot dx;$$

$$\therefore\ V = -\pi \int_{\frac{ab^2}{\sqrt{b^2 - x^2}}} \left\{ + \frac{x^3}{\sqrt{b^2 - x^2}} \right\}$$

$$= C - \pi \cdot \left\{ ab^2 \sin^{-1}\frac{x}{b} - y^2\sqrt{b^2 - x^2} - \tfrac{2}{3}(b^2 - x^2)^{\frac{3}{2}} \right\},$$

and $x = b$; $V = 0$; $\therefore\ 0 = C - \pi\left\{ab^2 \frac{\pi}{2}\right\}$; $\therefore\ C = \frac{\pi^2 ab^2}{2}$;

$$\therefore\ V = \frac{\pi^2 ab^2}{2} - \pi\left\{ab^2 \sin^{-1}\frac{x}{b} - \frac{\sqrt{b^2 - x^2}}{3}(x^2 + 2b^2)\right\}.$$

Let $x = 0$; $\therefore$ whole volume $= \pi b^3 \left\{\frac{\pi a}{2} + \frac{2b}{3}\right\}$.

Ex. 32. Find the volume generated by the revolution of the cissoid round its asymptote.



$AB = 2a$, $BM = x$, $MQ = y$.

Now $NQ^2 = \dfrac{AN^3}{BN}$,

or $x^2 = \dfrac{(2a-y)^3}{y}$;

$$\therefore\ V = \pi \int_x y^2 = \pi y^2 x - 2\pi \int_y yx.$$

But $x^2y^2 = y\,(2a-y)^3$; $\therefore\ xy = \sqrt{y}\,.(2a-y)^{\frac{3}{2}}$;

$$\therefore \int_y xy = \int_y (2a-y)\sqrt{2ay-y^2}$$

$$= \int_y (a-y)\sqrt{2ay-y^2} + a\int_y \sqrt{2ay-y^2}$$

$$= \frac{(2ay-y^2)^{\frac{3}{2}}}{3} + a\int_y \sqrt{2ay-y^2};$$

$$\therefore\ V = \pi\left\{(2ay-y^2)^{\frac{3}{2}} - \tfrac{2}{3}(2ay-y^2)^{\frac{3}{2}} - 2a\int_y \sqrt{2ay-y^2}\right\};$$

$$\therefore \text{ whole solid} = \int_{2a}^{0} y^2 = \pi\,.\,2a\,.\,\frac{\pi a^2}{2} = \pi^2 a^3.$$

Ex. 33. Find the solid generated by the revolution of the semi-cycloid round its base.

Make the base the axis of x; $\therefore\ \dfrac{dy}{dx} = \dfrac{\sqrt{2ay-y^2}}{y}$;

$$\therefore\ V = \int_y \pi y^2 \cdot \frac{dx}{dy} = \pi \cdot \int_y \frac{y^3}{\sqrt{2ay-y^2}}\,.$$

But $\displaystyle\int \frac{y^3}{\sqrt{2ay-y^2}} = -\frac{y^2\sqrt{2ay-y^2}}{3} - \frac{5}{2\,.\,3}\,ay\sqrt{2ay-y^2}$

$$-\frac{3\,.\,5}{2\,.\,3}\,a^2\sqrt{2ay-y^2} + \frac{3\,.\,5}{2\,.\,3}\,a^3 \text{ ver-sin}^{-1}\frac{y}{a};$$

$$\therefore\ V = \pi\int_0^{2a} \frac{y^3}{\sqrt{2ay-y^2}} = \frac{5\pi^2 a^3}{2}\,.$$

Ex. 34. Find the solid generated by the revolution of the cycloid round its axis.

$$V = \pi \int_x y^2 = \pi \int_\theta y^2 \frac{dx}{d\theta};$$

and $y = a\,(\theta + \sin\theta)$; $x = a\,(1 - \cos\theta)$;

$$\therefore\ V = \pi a^3 \int_0 (\theta + \sin\theta)^2 \sin\theta$$
$$= \pi a^3 \int_0 \{\theta^2 \sin\theta + 2\theta \sin^2\theta + \sin^3\theta\},$$

whence integrating from $\theta = 0$ to $\theta = \pi$;

$$\therefore\ V = \pi a^3 \left\{\frac{3\pi^2}{2} - \frac{8}{3}\right\}.$$

Ex. 35. To find the volume of a conical figure, the base of which is bounded by a given curve.

From A draw AD perpendicular to the base, and $= a$. In AD take $AN = x$, N being a point in a section bc, parallel and similar to the base BC.



Let A = area of the base,

S = area of section bc;

$$\therefore \frac{S}{A} = \frac{bN^2}{BD^2} = \frac{AN^2}{AD^2} = \frac{x^2}{a^2};$$

$$\therefore\ S = A\frac{x^2}{a^2},\ \text{and}\ \frac{dV}{dx} = S = A.\frac{x^2}{a^2};$$

$$\therefore\ V = \frac{A}{a^2}\int_x x^2 = \frac{Ax^3}{3a^2} + C,\ \text{and}\ C = 0;$$

$$\therefore\ ABC = \frac{Aa^3}{3a^2} = \frac{A.a}{3} = \text{base} \times \tfrac{1}{3}\ \text{of the altitude.}$$

Cor. This proposition is manifestly true for a pyramid of any base.

Ex. 36. To find the volume of a Groin; a solid of which in this instance, the sections parallel to the base are squares, and those perpendicular, bounded by a given curve.

Let the given curve AD be a quadrant

$AN = x$, $NP = y$; $AB = BD = a$;

therefore generating area $= (2y)^2 = 4y^2$;



$$\therefore \frac{dV}{dx} = 4y^2 = 4(2ax - x^2);$$

$$\therefore\ V = 4\left(ax^2 - \frac{x^3}{3}\right) = \frac{8}{3}a^3,\ \text{if}\ x = a.$$

Again, $\because$ generating surface = perimeter of square $= 8y$;

$$\therefore \frac{dS}{dx} = 8y\frac{ds}{dx} = 8\sqrt{a^2 - x^2}\,.\,\frac{a}{\sqrt{a^2 - x^2}} = 8a;$$

$$\therefore S = 8ax = 8a^2.$$

And similarly may the volume and surface be found, whatever be the curve APD. Also, if the base be any other figure, of which the area is a function of y, as a circle, a parabola, a triangle, &c. and APB be a curve of which the equation is $y = f(x)$, the surface and volume may be found.

Ex. 37. Find the solid generated by a parabolic area round its ordinate.



$AM = x$, $BN = x_1$, $AB = a$,

$MP = x$, $NP = y_1$, $BC = b$;

$$\frac{dV}{dx_1} = \pi y_1^2 = \pi(a - x)^2 = \pi\left(a - \frac{y^2}{4m}\right)^2$$

$$= \pi\left(\frac{4ma - y^2}{4m}\right)^2 = \frac{\pi}{(4m)^2}(b^2 - y^2)^2,$$

and $\dfrac{dV}{dx_1} = \dfrac{dV}{dy}$; $\therefore = \dfrac{\pi}{(4m)^2}\{b^4 - 2b^2y^2 + y^4\}$;

$$\therefore V = \frac{\pi}{(4m)^2}\left\{b^4y - \frac{2b^2y^3}{3} + \frac{y^5}{5}\right\};$$

$$\therefore \text{volume} = \frac{\pi b^5}{(4m)^2}\left\{1 - \tfrac{2}{3} + \tfrac{1}{5}\right\} = \frac{\pi b^5}{(4m)^2}\,.\,\frac{8}{15} = \frac{8}{15}\pi a^2 b.$$

Ex. 38. Find the volume and surface of the solid generated by the circle BQP round an axis ANx, in its own plane.



Let $AO = b$, $OB = a$,

$MQ = y$, $OM = x$.

Then surface generated by QP

$$= \pi(NP^2 - NQ^2) = \pi\{(b + y)^2 - (b - y)^2\} = 4\pi by;$$

$$\therefore \frac{dV}{dx} = 4\pi by; \quad \therefore V = 4\pi b\int_x y = 4\pi b\frac{\pi a^2}{2} = 2\pi^2 a^2 b,$$

$$S = 2\pi\,.\int_x (NP + NQ)\,.\frac{ds}{dx} = 4\pi b\,.\int_x \frac{ds}{dx} = 4\pi b\,.\,\pi a = 4\pi^2 ba.$$

Ex. 39. The surface of a sphere.

$$y = \sqrt{2ax - x^2}, \text{ and } \frac{dy}{dx} = \frac{a - x}{\sqrt{2ax - x^2}},$$

$$1 + \frac{dy^2}{dx^2} = 1 + \frac{(a-x)^2}{2ax - x^2} = \frac{a^2}{2ax - x^2} = \frac{a^2}{y^2},$$

$$S = 2\pi \int y \sqrt{1 + \frac{dy^2}{dx^2}} = 2\pi \int y \cdot \frac{a}{y} = 2\pi \int a = 2\pi ax + C,$$

$S = 0$, if $x = 0$; $\therefore C = 0$; $\therefore$ surface of a segment $= 2\pi ax$;

$\therefore$ surface of sphere $= 2\pi a \cdot 2a = 4\pi a^2$.

Ex. 40. Convex surface of a paraboloid.

$$y^2 = 4mx, \quad \frac{dy}{dx} = \frac{2m}{y};$$

$$\therefore 1 + \frac{dy^2}{dx^2} = 1 + \frac{4m^2}{y^2} = 1 + \frac{m}{x} = \frac{x+m}{x};$$

$$\therefore S = \int 2\pi y \sqrt{1 + \frac{dy^2}{dx^2}} = 4\pi \sqrt{m} \cdot \int \sqrt{x} \sqrt{\frac{x+m}{x}}$$

$$= 4\pi \sqrt{m} \int \sqrt{x+m} = 4\pi \sqrt{m} \tfrac{2}{3} (x+m)^{\frac{3}{2}} + C,$$

and if $x = 0$, $S = 0$; $\therefore C = -\frac{8}{3}\pi \sqrt{m} \cdot m^{\frac{3}{2}}$;

$$\therefore \text{surface} = \frac{8\pi\sqrt{m}}{3} \cdot \{(x+m)^{\frac{3}{2}} - m^{\frac{3}{2}}\}.$$

Ex. 41. Surface generated by a semi-cycloid round its base.

Here $\frac{dy}{dx} = \frac{\sqrt{2ay - y^2}}{y}$; $\therefore \sqrt{1 + \frac{dy^2}{dx^2}} = \sqrt{\frac{2a}{y}}$;

$$\therefore \frac{dS}{dy} = \frac{dS}{dx} \cdot \frac{dx}{dy} = 2\pi \sqrt{2ay} \cdot \frac{y}{\sqrt{2ay - y^2}};$$

$$\therefore S = 2\pi \sqrt{2a} \cdot \int_y \frac{y}{\sqrt{2a - y}}$$

$$= 2\pi \sqrt{2a} \{-2y \sqrt{2a - y} - \tfrac{4}{3}(2a - y)^{\frac{3}{2}}\};$$

$\therefore$ surface by semi-cycloid $= 2\pi \cdot \frac{4}{3}(2a)^2 = \frac{32}{3}\pi a^2$.

Ex. 42. Find the same when round the axis.

The vertex the origin, $\frac{dy}{dx} = \sqrt{\frac{2a-x}{x}}$.

$$\text{Surface} = 2\pi \int_x y \frac{ds}{dx} = 2\pi \left\{ ys - \int_x s \frac{dy}{dx} \right\}; \quad s = 2\sqrt{2ax}$$

$$= 2\pi \left\{ 2y\sqrt{2ax} - 2\sqrt{2a} \int_x \sqrt{x} \sqrt{\frac{2a-x}{x}} \right\}$$

$$= 4\pi \left\{ y\sqrt{2ax} - \sqrt{2a} \int_x \sqrt{2a-x} \right\}$$

$$= 4\pi\sqrt{2a} \left\{ y\sqrt{x} + \frac{2}{3} \left\{ (2a-x)^{\frac{3}{2}} \right\} \right\},$$

from $x = 0$ to $x = 2a$, or $y = 0$ to $y = \pi a$,

$$S = 4\pi\sqrt{2a} \left\{ \pi a \sqrt{2a} - \frac{2}{3} . (2a)^{\frac{3}{2}} \right\}$$

$$= 8\pi a \left\{ \pi a - \frac{4}{3} a \right\} = 8\pi a^2 . \left\{ \pi - \frac{4}{3} \right\}.$$

Ex. 43. To find the surface of the prolate spheroid.

$$y = \frac{b}{a}\sqrt{a^2 - x^2}, \text{ and } 1 + \frac{dy^2}{dx^2} = \frac{a^2 - e^2x^2}{a^2 - x^2},$$

$$\frac{dS}{dx} = 2\pi y \frac{ds}{dx} = 2\pi \frac{b}{a}\sqrt{a^2 - x^2} \sqrt{\frac{a^2 - e^2x^2}{a^2 - x^2}}$$

$$= \frac{2\pi b}{a}\sqrt{a^2 - e^2x^2} = 2\pi b . \sqrt{1 - \frac{e^2x^2}{a^2}};$$

$$\therefore S = \frac{\pi b a}{e} . \left\{ \sin^{-1}\left(\frac{ex}{a}\right) + \frac{ex}{a}\sqrt{1 - \frac{e^2x^2}{a^2}} \right\};$$

$$\therefore \left\{ \begin{matrix} \text{from } x = -a \\ \text{to } \ x = +a \end{matrix} \right\} S = \frac{2\pi b a}{e} . \left\{ \sin^{-1} e + e\sqrt{1 - e^2} \right\}$$

$$= 2\pi a^2 \left\{ \sqrt{1 - e^2} . \frac{\sin^{-1} e}{e} + 1 - e^2 \right\}.$$

Let $e = 0$, or spheroid become a sphere; $\therefore \frac{\sin^{-1} e}{e} = 1$,

and surface $= 2\pi a^2 \{1 + 1\} = 4\pi a^2$.

Ex. 44. To find the surface of an oblate spheroid.

$BM = x_1$ $CN = x$,

$MP = y_1$ $NP = y$,



$$\frac{dS}{dx_1} = 2\pi y_1 \cdot \frac{ds}{dx_1};$$

or $$\frac{dS}{dy} = 2\pi x \cdot \frac{ds}{dy};$$

$$\therefore \frac{dS}{dx} = 2\pi x \frac{ds}{dx} = 2\pi x \sqrt{\frac{a^2 - e^2x^2}{a^2 - x^2}}.$$

Make $\sqrt{a^2 - x^2} = z$; and $c = \frac{b}{e}$;

$$\therefore S = -2\pi e \int_z \sqrt{c^2 + z^2}$$

$$= C - \pi e \left\{ z\sqrt{c^2 + z^2} + c^2 \log (z + \sqrt{z^2 + c^2}) \right\}.$$

From $x = 0$ to $x = a$; or from $z = a$ to $z = 0$,

$$S = \pi e \cdot \left\{ a\sqrt{c^2 + a^2} + c^2 \log \frac{a + \sqrt{a^2 + c^2}}{c} \right\};$$

$$\therefore \text{surface} = 2S = 2\pi e \cdot \left\{ a\sqrt{c^2 + a^2} + c^2 \log \frac{a + \sqrt{a^2 + c^2}}{c} \right\}$$

$$= 2\pi e \left\{ \frac{a^2}{e} + \frac{a^2(1 - e^2)}{e^2} \log \frac{a + \frac{a}{e}}{\frac{a}{e}\sqrt{1 - e^2}} \right\}$$

$$= 2\pi a^2 \cdot \left\{ 1 + \frac{(1 - e^2)}{2e} \log \left(\frac{1 + e}{1 - e} \right) \right\};$$

which $= 4\pi a^2$, when $e = 0$, or if the spheroid become a sphere.

MISCELLANEOUS EXAMPLES.

(1) How much of the Earth's surface may be seen by a person elevated the $\frac{1}{n}$th part of the Earth's radius above it.

Ans. $\frac{1}{2n + 2}$ th part.

(2) Find the length of the curve, where $x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}$.

$s = 6a$.

(3) If h = height of a parabolic frustum, a and b the radii of the ends, shew that

$$\text{Frustum} = \frac{\pi h}{2} \cdot (a^2 + b^2).$$

(4) Find the area of the catenary, $y = \frac{a}{2}(e^{\frac{x}{a}} + e^{-\frac{x}{a}})$,

$$\text{Area} = a\sqrt{y^2 - a^2}.$$

(5) The area of $(x^2 + y^2)^2 = a^2x^2 - b^2y^2$ is

$$= ab + (a^2 - b^2)\tan^{-1}\left(\frac{a}{b}\right).$$

(6) The area of a parabolic segment, cut off by any chord $= \frac{2}{3}$ of circumscribing parallelogram.

(7) If $r = a\sec\frac{\theta}{2}$, the area included by the curve, the asymptotes and tangent at vertex $= 4a^2$.

(8) Find the area of $x^4y^4 - a^4y^4 = a^8$.

$$\text{Area} = \frac{a^2}{2}\left\{\log\sqrt{\frac{xy + a^2}{xy - a^2}} - \tan^{-1}\left(\frac{a^2}{xy}\right)\right\}.$$

(9) In a parabola, find the area included between the curve, its evolute, and its radius of curvature.

$$\text{Area} = 4\sqrt{\frac{x}{a}}\left\{a^2 + \frac{2}{3}ax + \frac{1}{5}x^2\right\}.$$

(10) If the subtangent of the logarithmic curve = that of the spiral, $\theta = \frac{a}{r}$: the arc included by two radii of the spiral = arc included by two respectively equal ordinates of the curve.

(11) Find the length of the spiral of Archimedes.

$$s = \frac{r}{2a}\sqrt{r^2 + a^2} + \frac{a}{2} \cdot \log\left(\frac{r + \sqrt{r^2 + a^2}}{2a}\right).$$

(12) The length of the epiycyloid after one revolution of the generating circle $= 8\frac{b}{a}(a + b)$, and the area between the epicycloid and the circle $= \pi b^2\left(3 + \frac{2b}{a}\right)$.

(13) Find the volume generated by the revolution of the Witch, round its asymptote. Volume $= 4\pi^2a^3$.

(14) The area of the curve in which

$$r = \frac{(a^2 - b^2)\sin\theta\cos\theta}{\sqrt{a^2\sin^2\theta + b^2\cos^2\theta}}; \text{ is} = \frac{\pi}{2}(a - b)^2.$$

CHAPTER VII.

DIFFERENTIAL EQUATIONS.

95. In the integrations which have been performed in the preceding Chapters, the differential coefficient has either been a given function of one of the variables, or else has been expressed in such terms of the two, that by a very evident process it has been reduced to a function of one only. We now proceed to integrate differentials, when the differential coefficients and the variables x and y are mingled together.

96. Differential equations are divided into classes, dependent upon the *order* and *degree* of the differential coefficient.

Thus an equation involving

$$\frac{dy}{dx}, \quad \frac{d^2y}{dx^2}, \quad \frac{d^3y}{dx^3}, \text{ &c. } \frac{d^ny}{dx^n},$$

is called a differential equation of the n^{th} order and of the first degree, while one containing

$$\frac{dy}{dx}, \quad \left(\frac{dy}{dx}\right)^2, \quad \left(\frac{dy}{dx}\right)^3, \text{ &c. } \left(\frac{dy}{dx}\right)^n$$

is said to be of the first order, and of the n^{th} degree: and finally, an equation in which are to be found the n^{th} powers of the differential coefficients and the m^{th} differential coefficient, is named an equation of the m^{th} order and the n^{th} degree.

We shall begin with that class in which the first power of the first differential coefficient is alone found.

Differential Equations of the first Order and the first Degree.

97. These are included under the formula

$$M + N \cdot \frac{dy}{dx} = 0,$$

where M and N may be any functions of x and y; we shall however in the first place treat of homogeneous equations.

98. Let $M + N\frac{dy}{dx} = 0$, be a homogeneous equation, or one in which the sum of the indices of y and x together, is the same in every term.

$$\text{Make } y = vx; \quad \therefore \frac{dy}{dx} = x + v\frac{dx}{dv}.$$

Divide by N and the equation becomes,

$$\frac{M}{N} + \frac{dy}{dx} = 0, \text{ or } \frac{M}{N} + x + v\frac{dx}{dv} = 0.$$

But $\frac{M}{N}$ is of no dimensions or is a function of $\frac{y}{v}$ or x.

$$\text{Let } \therefore \frac{M}{N} = f(x); \quad \therefore v\frac{dx}{dv} = -\{x + f(x)\};$$

$$\therefore \frac{dv}{vdx} = -\frac{1}{x + f(x)}; \quad \therefore \log\left(\frac{v}{c}\right) = -\int_x \frac{1}{x + f(x)},$$

which may be integrated by the ordinary rules.

We put $x = yx$, or $y = vx$, as may be most convenient, for the solution is more easily effected, when we substitute for that differential coefficient which involves the fewest terms.

Ex. 1. Let $v + y = (v - y)\frac{dy}{dv}$.

$$\text{Here make } y = vx; \quad \therefore \frac{dy}{dv} = x + v\frac{dx}{dv};$$

$$\therefore x + v\frac{dx}{dv} = \frac{v + y}{v - y} = \frac{1 + x}{1 - x}; \quad \therefore v\frac{dx}{dv} = \frac{1 + x^2}{1 - x};$$

$$\therefore \frac{dv}{vdx} = \frac{1 - x}{1 + x^2} = \frac{1}{1 + x^2} - \frac{x}{1 + x^2};$$

$$\therefore \log\left(\frac{v}{c}\right) = \tan^{-1}x - \log\sqrt{1 + x^2};$$

$$\therefore \log\left(\frac{v}{c}\sqrt{1 + x^2}\right), \text{ or } \log\frac{\sqrt{v^2 + y^2}}{c} = \tan^{-1}\frac{y}{v}.$$

Ex. 2. Find the curve in which the subtangent is equal to the sum of the abscissa and ordinate.

$$\text{Here } y\frac{dv}{dy} = v + y; \text{ and let } v = yx;$$

$$\therefore \frac{dv}{dy} = x + y\frac{dx}{dy} = \frac{v + y}{y} = x + 1;$$

$$\therefore \frac{dy}{ydx} = 1; \quad \therefore \log\left(\frac{y}{c}\right) = x = \frac{v}{y}.$$

Ex. 3. Find the curve in which the subnormal $= y - x$.

$$y\frac{dy}{dx} = y - x; \quad \therefore \frac{dy}{dx} = 1 - \frac{x}{y}.$$

$$\text{Let } y = xz; \quad \therefore z + x\frac{dz}{dx} = 1 - \frac{1}{z} = \frac{z-1}{z};$$

$$\therefore \frac{dx}{x\,dz} = \frac{-z}{z^2 - z + 1};$$

$$\therefore \log\left(\frac{\sqrt{y^2 - yx + x^2}}{c}\right) = \frac{1}{\sqrt{3}}\cot^{-1}\left(\frac{2y - x}{x\sqrt{3}}\right).$$

Ex. 4. Find the curve in which the distance from the origin to a point in the curve equals the subtangent.

$$\text{Here } AP = NT, \text{ or } \sqrt{y^2 + x^2} = y\frac{dx}{dy}.$$

$$\text{Make } x = yz; \quad \therefore z + y.\frac{dz}{dy} = \frac{\sqrt{x^2 + y^2}}{y} = \sqrt{1 + z^2};$$

$$\therefore \frac{dy}{y\,dz} = \frac{1}{\sqrt{1 + z^2} - z} = \sqrt{1 + z^2} + z;$$

$$\text{whence } \log\left\{\frac{y^2}{c^2(x + \sqrt{x^2 + y^2})}\right\} = \frac{x}{y^2}(x + \sqrt{x^2 + y^2}).$$

Ex. 5. $(\sqrt{x} - \sqrt{y}) = \sqrt{y}.\frac{dx}{dy}$. Make $x = yz$.

Ex. 6. $x\frac{dy}{dx} - y = \sqrt{x^2 + y^2}$. Make $y = xz$,

Then $x^2 = c^2 + 2cy$.

Ex. 7. $x\frac{dy}{dx} - y = y\log\left(\frac{y}{x}\right); \quad \therefore y^c = x^c e^x$.

Ex. 8. $\int_x y = \frac{y^3}{x}; \quad \therefore (x^2 - 2y^2)^3 = cx^2$.

Ex. 9. NT and NP are the subtangent and ordinate of a curve of which the vertex is A, and $\tan TPA = m \tan APN$, find the equation to the curve.

$$(y^2 + x^2)^{m+1} = c^{2m}x^2.$$

Ex. 10. $\int_x(xy) = \frac{x}{n}\int_x y; \quad \therefore y^{n-1}.x^{n-2} = a^{2n-3}$.

Ex. 11. $\int_x(xy^2) = \frac{2}{3}x\int_x y^2$; $\therefore\ y^2 = cx$.

99. The equation $(a+bx+cy)dx+(a_1+b_1x+c_1y)dy=0$ can be rendered homogenous by making

$$v = a + bx + cy, \quad \text{and} \quad z = a_1 + b_1x + c_1y;$$

$$\therefore\ dv = bdx + cdy, \quad dz = b_1dx + c_1dy;$$

$$\therefore\ c_1dv - cdz = (bc_1 - b_1c)dx,$$

$$bdz - b_1dv = (bc_1 - b_1c)dy;$$

whence by substitution the equation becomes

$$v(c_1dv - cdz) + z(bdz - b_1dv) = 0,$$

$$\text{or } (vc_1 - b_1z)dv + (bz - cv)dz = 0,$$

which is a homogenous equation.

Cor. This method is inapplicable when $bc_1 = b_1c$; but since then $c_1 = \frac{b_1c}{b}$, the equation becomes

$$(a + bx + cy)dx + \left(a_1 + b_1x + b_1\frac{cx}{b}\right)dy = 0,$$

$$\text{i.e. } (a + bx + cy)dx + \left\{a_1 + \frac{b_1}{b}(bx + cy)\right\}dy = 0,$$

an equation in which the variables may be separated by making $bx + cy = z$; $\therefore\ dx = \frac{dz - cdy}{b}$;

$$\therefore\ (a + z)\frac{dz - cdy}{b} + \left(a_1 + \frac{b_1z}{b}\right)dy = 0;$$

$$\therefore\ (a + z)dz - (ca + cz - a_1b - b_1z)dy = 0;$$

$$\therefore\ \frac{dy}{dz} = \frac{(a + z)}{ca - a_1b + (c - b_1)z} = \frac{(a + z)}{\alpha + \beta z},$$

where $\alpha = ca - a_1b$ and $\beta = c - b_1$, the integral of which may be readily found.

100. To integrate the *linear* equation, (so called since the first power of y is alone involved),

$$\frac{dy}{dx} + Py = Q \ldots\ldots (1),$$

in which P and Q are functions of x.

$$\text{Since } \frac{d}{dx}ye^{\int.P} = \frac{dy}{dx}e^{\int.P} + e^{\int.P}.Py$$

$$= e^{\int.P}\left\{\frac{dy}{dx} + Py\right\}.$$

It is obvious that if both sides of (1) be multiplied by $e^{\int.P}$, the left hand will be a complete differential, and the right hand a function of x alone; multiply therefore by $e^{\int.P}$;

$$\therefore e^{\int.P}\left\{\frac{dy}{dx} + Py\right\} = e^{\int.P}.Q; \quad \text{or } \frac{d}{dx}(ye^{\int.P}) = e^{\int.P}.Q;$$

$$\therefore \text{ integrating } ye^{\int.P} = C + \int e^{\int.P}.Q;$$

$$\text{or } \quad y = Ce^{-\int.P} + e^{-\int.P}\int e^{\int.P}.Q.$$

Ex. 1. Let $\frac{dy}{dx} + y = ax^3$.

Here $P = 1$, $\int.P = x$; $\therefore e^{\int.P} = e^x$, $Q = ax^3$;

$$\therefore ye^x = C + a\int e^x.x^3 = C + ae^x\left\{x^3 - 3x^2 + 6x - 6\right\};$$

$$\therefore y = Ce^{-x} + a\left\{x^3 - 3x^2 + 6x - 6\right\}.$$

Ex. 2. $(1 + x^2)\frac{dy}{dx} - yx = a$; or $\frac{dy}{dx} - y\frac{x}{1+x^2} = \frac{a}{1+x^2}$.

Here $P = -\frac{x}{1+x^2}$; $\int.P = \log\frac{1}{\sqrt{1+x^2}}$; $e^{\int.P} = \frac{1}{\sqrt{1+x^2}}$;

$$\therefore y\frac{1}{\sqrt{1+x^2}} = a\int.\frac{1}{\sqrt{1+x^2}} \times \frac{1}{1+x^2}$$

$$= a\int.\frac{1}{(1+x^2)^{\frac{3}{2}}} = \frac{ax}{\sqrt{1+x^2}} + c;$$

$$\therefore y = ax + c\sqrt{1+x^2}.$$

Ex. 3. $\frac{dy}{dx} = a + bx + cy$; $\therefore Ce^{cx} = b + c(a + bx + cy)$.

Ex. 4. $\frac{dy}{dx} + \frac{ny}{x} = \frac{a}{x^n}$; $\therefore yx^n = a(x + b)$.

101. The equation $y^{m-1}\frac{dy}{dx} + Py^m = Qy^n$ may be reduced to the preceding form, in the following manner.

Divide by y^n;

$$\therefore\ y^{m-n-1}\frac{dy}{dx} + Py^{m-n} = Q.$$

Let $y^{m-n} = (m-n)z$; $\quad\therefore\ y^{m-n-1}\dfrac{dy}{dx} = \dfrac{dz}{dx}$;

$$\therefore\ \frac{dz}{dx} + (m-n)Pz = Q;$$

which is of the required form.

Ex. 1. $\quad v\dfrac{dv}{ds} - \dfrac{hv^2}{s} = -\dfrac{m}{s^2}.$

Let $v^2 = 2z$; $\quad\therefore\ v\dfrac{dv}{ds} = \dfrac{dz}{ds}$;

$$\therefore\ \frac{dz}{ds} - \frac{2hz}{s} = -\frac{m}{s^2}.$$

Here $P = -\dfrac{2h}{s}$; $\quad\therefore \int_s P = -2h\log(s) = \log\dfrac{1}{s^{2h}}$; $\quad\therefore e^{\int_s P} = \dfrac{1}{s^{2h}}$;

$$\therefore\ zs^{-2h} = -m\int_s s^{-(2h+2)} = c + \frac{ms^{-(2h+1)}}{2h+1};$$

$$\therefore\ z = \frac{v^2}{2} = cs^{2h} + \frac{m}{(2h+1)s}.$$

Ex. 2. $\quad \dfrac{dy}{dx} + y = xy^3$; $\quad\therefore\ \dfrac{1}{y^2} = x + \tfrac{1}{2} + Ce^{2x}.$

Ex. 3. $\quad \dfrac{dy}{dx} + \dfrac{xy}{1-x^2} = x\sqrt{y}$; $\quad\therefore \sqrt{y} = C\sqrt[4]{1-x^2} - \tfrac{1}{3}(1-x^2).$

Ex. 4. $\quad xy^2dy + y^3dx = \dfrac{a\,dx}{x}$;

$$\therefore\ y^3 = \frac{3a}{2x} + \frac{C}{x^3}.$$

Integration of exact Differentials. The method of finding a factor which will render a function integrable.

102. The equation $Mdx + Ndy = 0$ is not always the result of the differentiation of $f(xy) = c$; for after the differentiation, its terms may have been divided by a common factor, or the equation may arise from the elimination of an arbitrary constant between the primitive equation and its derivative.

But whenever $Mdx + Ndy = 0$ is the complete differential of a function of two variables, the condition $\frac{d^2u}{dxdy} = \frac{d^2u}{dydx}$ is fulfilled, or since $M = \frac{du}{dx}$ and $N = \frac{du}{dy}$;

$$\therefore \frac{dM}{dy} = \frac{d^2u}{dxdy} = \frac{dN}{dx}.$$

Hence, we have a method by which we may find whether any equation of the form $Mdx + Ndy = 0$, be a complete differential; and if it be, since then $\frac{du}{dx} = M$, and $\frac{du}{dy} = N$, we can by integrating these partial differential equations find u; for since $M = \frac{du}{dx}$, M is the partial differential coefficient of u, with regard to x, considering x alone to vary, and its integral will give all the terms in which x is to be found: let the integration be performed. Then

$$u = \int_x M + Y.$$

Here instead of adding a constant C, we put Y, for as y has been supposed not to vary, the constant will include those terms of the original equation which are functions of y alone. Next to determine Y: differentiate with regard to y;

$$\therefore \frac{du}{dy} = \frac{d\int_x M}{dy} + \frac{dY}{dy}.$$

But $\frac{du}{dy} = N$; $\therefore \frac{dY}{dy} = N - \frac{d\int_x M}{dy}$;

$$\therefore Y = \int_y \left(N - \frac{d\int_x M}{dy}\right) + C;$$

$$\therefore u = \int_x M + \int_y \left(N - \frac{d\int_x M}{dy}\right) + C.$$

103. Since Y ought to be a function of y only,

$\int_y \left(N - \frac{d\int_x M}{dy}\right)$ should be independent of x.

To prove this, let $y + \delta y$ be put for y in $\int_x M$;

$$\therefore \int_x \left(M + \frac{dM}{dy}\delta y + \&c.\right) = \int_x M + \delta y \int_x \frac{dM}{dy} + \&c.$$

Hence $\frac{d\int_x M}{dy} = \int \frac{dM}{dy}$; $\therefore\ Y = \int_y \left(N - \int_x \frac{dM}{dy}\right) + C$;

$\therefore\ \frac{dY}{dy} = N - \int_x \frac{dM}{dy}$; $\therefore\ \frac{d^2Y}{dx\,dy} = \frac{dN}{dx} - \frac{dM}{dy} = 0$;

or since $\frac{dY}{dy}$ differentiated with regard to x vanishes; Y is a function of y only: the same result would have been obtained by integrating N or $\frac{du}{dy}$ in the first instance.

Ex. 1. Let $du = \frac{2\,dx}{\sqrt{x^2-y^2}} - \frac{2x\,dy}{y\sqrt{x^2-y^2}}$.

Here $M = \frac{2}{\sqrt{x^2-y^2}}$; $N = \frac{-2x}{y\sqrt{x^2-y^2}}$

$\frac{dM}{dy} = \frac{2y}{(x^2-y^2)^{\frac{3}{2}}}$, $\frac{dN}{dx} = \frac{-2}{y}\left(\frac{-y^2}{(x^2-y^2)^{\frac{3}{2}}}\right) = \frac{2y}{(x^2-y^2)^{\frac{3}{2}}}$;

$\therefore\ u = \int_x M + Y = 2\log\left(x + \sqrt{x^2-y^2}\right) + Y$,

$\frac{du}{dy} = \frac{-2y}{\left(x+\sqrt{x^2-y^2}\right)\sqrt{x^2-y^2}} + \frac{dY}{dy} = \frac{-2x}{y\sqrt{x^2-y^2}}$;

$\therefore\ \frac{dY}{dy} = \frac{2}{\sqrt{x^2-y^2}}\left\{\frac{y}{x+\sqrt{x^2-y^2}} - \frac{x}{y}\right\}$

$= \frac{-2}{\sqrt{x^2-y^2}}\left\{\frac{x^2-y^2+x\sqrt{x^2-y^2}}{y\left(x+\sqrt{x^2-y^2}\right)}\right\} = \frac{-2}{y}$;

$\therefore\ Y = C - 2\log y$;

$\therefore\ u = \log\left(\frac{x+\sqrt{x^2-y^2}}{y}\right)^2 + C$.

Ex. 2. Let $du = \frac{a(x\,dx+y\,dy)}{\sqrt{x^2+y^2}} + \frac{y\,dx-x\,dy}{x^2+y^2} + 3by^2\,dy = 0$.

$M = \frac{ax}{\sqrt{x^2+y^2}} + \frac{y}{x^2+y^2}$; $N = \frac{ay}{\sqrt{x^2+y^2}} - \frac{x}{x^2+y^2} + 3by^2$.

Here $\frac{dM}{dy} = \frac{-ay}{(x^2+y^2)^{\frac{3}{2}}} + \frac{x^2-y^2}{(x^2+y^2)^2} = \frac{dN}{dx}$;

$\therefore\ u = \int_x M + Y = a\sqrt{x^2+y^2} + \tan^{-1}\frac{x}{y} + Y$,

$$\frac{du}{dy} = \frac{ay}{\sqrt{x^2+y^2}} - \frac{x}{y^2+x^2} + \frac{dY}{dy}; \quad \therefore \frac{dY}{dy} = 3by^2;$$

$$\therefore Y = by^3 + C, \text{ and } u = a\sqrt{x^2+y^2} + \tan^{-1}\frac{x}{y} + by^3 + C.$$

Ex. 3. $$\int \frac{x\,dy - y\,dx}{x^2+y^2} = \tan^{-1}\frac{x}{y} + C.$$

Ex. 4. $$\frac{dx}{x} + \frac{y^2dx}{x^3} - \frac{y\,dy}{x^2} + \frac{dy}{2y} + \frac{y\,dx - x\,dy}{x^2}\sqrt{x^2+y^2} = 0;$$

$$\therefore \log(xy) - \frac{y}{x^2}(y + \sqrt{x^2+y^2}) + \log(y\sqrt{x^2+y^2} - y^2) = C.$$

104. When the equation $M\,dx + N\,dy = 0$ does not fulfil the *criterion of integrability*, $\frac{dM}{dy} = \frac{dN}{dx}$, it is no longer a complete differential, some factor having disappeared from it. Could however the factor be restored, every equation of this class might be integrated by the same process: but there is great difficulty in finding this factor; in most cases the differential equation, by which it is to be determined, is more complicated than the original one.

Thus, suppose z to be the factor, then $Mz\,dx + Nz\,dy = 0$ is a complete differential, and therefore

$$\frac{d(Mz)}{dy} = \frac{d(Nz)}{dx};$$

$$\therefore z\frac{dM}{dy} + M\frac{dz}{dy} = z\frac{dN}{dx} + N\frac{dz}{dx};$$

whence z is to be found, a problem seldom practicable.

105. When, however, any factor of the equation

$$M\,dx + N\,dy = 0$$

is known, an infinite number of factors may be found which will render the equation integrable; for let z be a factor,

$$\therefore du = zM\,dx + Nz\,dy;$$

$$\therefore \phi(u)\,du = z\phi(u)\,M\,dz + Nz\phi(u)\,dy;$$

and since the first member of the equation is an exact differential, the second member is also; $\therefore$ z multiplied by any function of (u), will make the equation integrable.

We may sometimes find the factor when the differential equation can be divided into two parts, for each of which a factor can be found; for let the equation be

$$pdx + qdy + p_1dx + q_1dy = 0 \quad (1),$$

and suppose that $\varkappa$ and $\varkappa_1$ are the factors which will make $pdx + qdy$ and $p_1dx + q_1dy$ integrable, so that

$$\varkappa pdx + \varkappa qdy = du; \text{ and } \varkappa_1 p_1dx + \varkappa_1 q_1dy = du_1;$$

$\therefore \varkappa\phi(u)$ and $\varkappa_1\phi(u_1)$ will include all the factors which will render the two equations separately integrable; if therefore we can make $\varkappa\phi(u) = \varkappa_1\phi(u_1)$, we shall obtain a factor which will render the equation (1) integrable.

Ex. 1. Let $\dfrac{adx}{x} + \dfrac{bdy}{y} = \dfrac{cx^m dx}{y^b}$.

$$\because \frac{adx}{x} + \frac{bdy}{y} = d.\log(x^a y^b); \quad \therefore \varkappa = 1; \quad u = x^a.y^b;$$

$\dfrac{cx^m dx}{y^b}$ is integrable if $\varkappa_1 = y^b$; $\quad \therefore u_1 = \dfrac{cx^{m+1}}{m+1}$;

$$\therefore \varkappa\phi(u) = \phi(x^a y^b); \quad \varkappa_1\phi(u_1) = y^b\phi(x^{m+1}).$$

Let $\phi(x^a y^b) = x^{ka} y^{kb}$; $\quad \phi(x^{m+1}) = x^{k_1 m+1}$;

$$\therefore x^{ka} y^{kb} = y^b x^{k_1 m+1}; \quad \therefore kb = b; \quad \therefore k = 1; \quad k_1 = \frac{a-1}{m};$$

$\therefore$ the factor is $x^a y^b$, whence integrating

$$x^a y^b = \frac{cx^{m+a+1}}{m+a+1} + C.$$

Ex. 2. $xdy - 2ydx = adx$. Here

$$\varkappa = \frac{1}{xy}; \quad u = \log\left(\frac{y}{x^2}\right); \quad \varkappa_1 = 1; \quad u_1 = ax;$$

$$\therefore \frac{1}{x^3} \text{ is the factor; and } y + cx^2 = \frac{a}{2}.$$

Ex. 3. $aydx + bxdy = x^m y^n (a_1 ydx + b_1 xdy)$.

The factor is $x^{ka-1} y^{kb-1}$; $\quad k = \dfrac{a_1 n - b_1 m}{ab_1 - a_1 b}$:

the integral of this equation might also be found by making $x^a y^b = z$; $x^{a_1} y^{b_1} = v$, from which we shall obtain

$$\frac{z^k}{a_1 n - b_1 m} + \frac{v^{k_1}}{an - bm} = C; \quad \text{where } k_1 = \frac{an - bm}{ab_1 - a_1 b}.$$

106. The factor may be determined when z contains only one variable as x, for then $\frac{dz}{dy} = 0$, and therefore

$$\frac{dz}{z\,dx} = \frac{1}{N}\left(\frac{dM}{dy} - \frac{dN}{dx}\right).$$

The right-hand side must be a function of x only, which is the case in the linear equation, for $N = 1$, and M contains only the first power of y; therefore integrating,

$$\log \frac{z}{c} = X; \quad \therefore\ z = ce^{X}.$$

But to find *a priori* the multiplier which will make the equation $dy + (Py - Q)\,dx = 0$ an exact differential.

Let z be the multiplier: multiply by it;

$$\therefore\ z\,dy + z(Py - Q)\,dx = N dy + M dx;$$

$$\therefore\ \frac{dN}{dx} = \frac{dz}{dx};\quad \frac{dM}{dy} = (Py - Q)\,\frac{dz}{dy} + Pz;$$

$$\therefore\ \frac{dz}{dx} = (Py - Q)\frac{dz}{dy} + Pz;$$

$$\therefore\ \frac{dz}{dx}\,dx = (Py - Q)\,dx\,\frac{dz}{dy} + Pz\,dx$$

$$= -\frac{dz}{dy}\,dy + Pz\,dx;\ \text{ since } (Py - Q)\,dx = -dy;$$

$$\therefore\ \frac{dz}{dx}\,dx + \frac{dz}{dy}\,dy = dz = Pz\,dx;$$

$$\text{i.e. } \frac{1}{z}\frac{dz}{dx} = P;\quad \therefore\ z = e^{\int P};$$

which justifies the assumption made, in article (100).

107. The factor may also be found when the equation is homogeneous.

For let $M + N\frac{dy}{dx} = 0$ be the differential equation, supposed to be homogenous and of m dimensions, and let z be the factor, a homogeneous function of the n^{th} degree;

$$\therefore\ zM\,dx + zN\,dy = du \ldots\ldots\ldots (1).$$

Hence, since u must be of $m + n + 1$ dimensions,

$$zMx + zNy = (m + n + 1)u \ldots\ldots\ldots (2),$$

Art. (112), Diff. Calc.; therefore dividing (1) by (2),

$$\frac{Mdx + Ndy}{Mx + Ny} = \frac{1}{m+n+1} \cdot \frac{du}{u};$$

and since the right-hand side of the equation is a complete differential, the left-hand must be so also; and $\therefore \frac{1}{Mx + Ny}$ is the factor required.

Ex. Let $ydy + (x - ny)dx = 0$;

$$\therefore \text{ the factor} = \frac{1}{x^2 - nyx + y^2};$$

$$\therefore Mz = \frac{x - ny}{x^2 - nyx + y^2}, \quad Nz = \frac{y}{x^2 - nyx + y^2},$$

$$\text{and } \frac{d(Mz)}{dy} = \frac{ny^2 - 2xy}{(x^2 - nyx + y^2)^2} = \frac{d(Nz)}{dx}.$$

108. We shall now add a few problems illustrating the solution of differential equations.

Find the curve which cuts any number of curves of a given species at a given angle.

Let y and x be the co-ordinates of the curve of given species,

y_1 and x_1 those of the required curve,

m = tangent of given angle.

$$\text{Then } \tan^{-1} m = \tan^{-1} \frac{dy}{dx} - \tan^{-1} \frac{dy_1}{dx_1};$$

$$\therefore m = \frac{\frac{dy}{dx} - \frac{dy_1}{dx_1}}{1 + \frac{dy}{dx} \cdot \frac{dy_1}{dx_1}},$$

and $\frac{dy}{dx}$ may be found from the given curve, and is a function of x and y, or $\phi(xy)$, and since at the point of intersection the co-ordinates of both curves are the same, we may for x_1 and y_1 put x and y; and then the equation to the required curve is

$$m\left\{1 + \phi(xy)\frac{dy}{dx}\right\} = \phi(xy) - \frac{dy}{dx},$$

which is of the first order and degree.

Cor. If the required curve cut the given curves at right angles,

$$\text{then } m = \frac{1}{0}; \quad \therefore\ 1 + \phi(xy)\frac{dy}{dx} = 0; \quad \therefore \frac{dy}{dx} = -\frac{1}{\phi(xy)},$$

which is the equation to the *Orthogonal Trajectory*.

Ex. 1. Find the curve which will cut all the parabolas that have a common vertex and axis at right angles.

Let $y^2 = 2mx$ be the equation to one of the parabolas;

$$\therefore\ \phi(xy) = \frac{m}{y} = \frac{y}{2x};$$

$$\therefore \frac{dy}{dx} = -\frac{2x}{y}; \quad \therefore \frac{y^2}{2} = (c^2 - x^2),$$

the equation to an ellipse of which the centre is the common vertex of the parabolas, and the major axis is perpendicular to the common axis, the ratio of the axes being $\sqrt{2} : 1$; c being indeterminate shews that any ellipse of which the axes are in the given ratio will cut the parabolas at right angles.

Ex. 2. Find the curve which will cut at right angles all the ellipses that have a common centre, coincident major axes, and the ratio of their axes constant.

Let $y^2 = n(a^2 - x^2)$ be the equation to one of the ellipses in which $\sqrt{n} = \frac{b}{a}$;

$$\therefore \frac{dy}{dx} = -n\frac{x}{y} = -\frac{dx}{dy};$$

$$\therefore n\frac{dy}{y} = \frac{dx}{x}; \quad \therefore n\log\left(\frac{y}{b}\right) = \log\left(\frac{x}{c}\right); \quad \therefore y^n = \frac{b^n}{c}x,$$

the equation to a parabola, of which the vertex is in the common centre of the ellipses.

If $n = 2$, $y^2 = \frac{b^2}{c}x$, the common parabola: this case is obviously the converse of the preceding problem.

Ex. 3. Find the curve which intersects at an angle of 45^0, all the straight lines drawn from the origin to meet it.

Let $y = ax$ be one of the lines;

$$\therefore\ \phi(xy) = a = \frac{y}{x}; \quad m = 1; \quad \therefore\ 1 + \frac{y}{x}\frac{dy}{dx} = \frac{y}{x} - \frac{dy}{dx},$$

a homogeneous equation,

$$\text{whence } \log\left(\sqrt{\frac{x^2+y^2}{c}}\right) = -\tan^{-1}\left(\frac{y}{x}\right).$$

$$\text{Let } y = r\sin\theta,\ x = r\cos(\pi-\theta) = -r\cos\theta;$$

$$\therefore r = \sqrt{x^2+y^2},\ \frac{y}{x} = -\tan\theta;\ \therefore \log\left(\frac{r}{c}\right) = \theta;\ r = ce^{\theta},$$

the equation to the logarithmic spiral.

109. To integrate *Riccati's* equation, so called from its proposer,

$$\frac{dy}{dx} + by^2 = ax^m.$$

(1). If $m = 0$, then $\frac{dy}{dx} = a - by^2$, which is easily integrable.

(2). If m be not $= 0$, we must proceed as follows.

CASE 1. Let $y = \frac{1}{bx} + \frac{z}{x^2}$; $\therefore \frac{dy}{dx} = -\frac{1}{bx^2} + \frac{dz}{dx}\cdot\frac{1}{x^2} - \frac{2z}{x^3}$,

$$by^2 = \frac{1}{bx^2} + \frac{bz^2}{x^4} + \frac{2z}{x^3};$$

$$\therefore \frac{dy}{dx} + by^2 = \frac{dz}{dx}\frac{1}{x^2} + \frac{bz^2}{x^4} = ax^m;$$

$$\therefore \frac{dz}{dx} + \frac{bz^2}{x^2} = ax^{m+2},$$

which is homogeneous if $m + 2 = 0$, or $m = -2$, and the variables may be separated if $m = -4$; for then

$$\frac{dz}{dx} + \frac{bz^2}{x^2} = \frac{a}{x^2};\quad \therefore \frac{dz}{bz^2 - a} + \frac{dx}{x^2} = 0.$$

If m have any other value, make

$$z = \frac{1}{y_1};\ x^{m+3} = x_1;\quad \therefore dz = -\frac{dy_1}{y_1^2};\ x^{m+2}dx = \frac{dx_1}{m+3}.$$

$$\text{Also } \frac{1}{x} = x_1^{-\frac{1}{m+3}};\quad \therefore \frac{dx}{x^2} = \frac{dx_1}{m+3}x_1^{-\frac{m+4}{m+3}},$$

$$\text{whence } -\frac{dy_1}{y_1^2} + \frac{b}{(m+3)y_1^2}x_1^{-\frac{m+4}{m+3}}dx_1 = \frac{a}{m+3}dx_1.$$

$$\text{Let } \frac{b}{m+3} = a_1;\ \frac{a}{m+3} = b_1;\ -\frac{m+4}{m+3} = m_1.$$

Then $dy_1 + b_1 y_1^2 dx_1 = a_1 x_1^{m_1} dx_1$,

which is of the same form as the original equation, and may be made homogeneous if $m_1 = -2$, and the variables may be separated by the preceding process if $m_1 = -4$.

By continuing the same methods it is evident that we shall have a similar equation, $dy_2 + b_2 y_2^2 dx_2 = a_2 x_2^{m_2} dx_2$,

where $m_2 = -\dfrac{m_1 + 4}{m_1 + 3}$; and b_2, a_2 are derived from b_1 and a_1 as these were from b and a; which equation will be integrable if $m_2 = -4$.

And hence if among the series of indices

$$-m, \quad -\frac{m+4}{m+3}, \quad -\frac{m_1+4}{m_1+3}, \quad -\frac{m_2+4}{m_2+3}, \quad \&c.$$

any one $= -4$, the equation is integrable. And by successively putting these indices $= -4$, we find the values of m to be, -4, $-\dfrac{8}{3}$, $-\dfrac{12}{5}$, $-\dfrac{16}{7}$, &c., which are included under the form $-\dfrac{4n}{2n-1}$, n being any integer.

CASE 2. Make in the original equation $y = \dfrac{1}{y_1}$;

$$\therefore -\frac{dy_1}{y_1^2} + \frac{b}{y_1^2} dx = a x^m dx; \quad \therefore dy_1 + a y_1^2 x^m dx = b dx.$$

Let $x^{m+1} = x_1$; $\therefore x = x_1^{\frac{1}{m+1}}$; $\therefore dx = \dfrac{1}{m+1} x_1^{-\frac{m}{m+1}} dx_1$.

$$\text{And } dy_1 + \frac{a}{m+1} y_1^2 dx_1 = \frac{b}{m+1} x_1^{-\frac{m}{m+1}} dx_1;$$

or putting $\dfrac{a}{m+1} = b_1$, $\dfrac{b}{m+1} = a_1$, and $-\dfrac{m}{m+1} = m_1$;

$$dy_1 + b_1 y_1^2 dx_1 = a_1 x_1^{m_1} dx_1,$$

which may be integrated by the former method if $m_1 = \dfrac{-4n}{2n-1}$;

or if $\dfrac{m}{m+1} = \dfrac{4n}{2n-1}$, whence $m = \dfrac{-4n}{2n+1}$. Hence Riccati's equation is integrable when m is of the form $\dfrac{-4n}{2n \mp 1}$. The first case belongs to the upper, the second to the lower sign.

Ex. 1. Integrate $dy + y^2 dx = \frac{a^2 dx}{x^{\frac{4}{3}}}$.

Here $-\frac{4}{3}$ is of the form $-\frac{4n}{2n+1}$;

$\therefore$ let $y = \frac{1}{y_1}$, and let $x^{m+1} = x^{-\frac{4}{3}+1} = x^{-\frac{1}{3}} = x_1$;

$\therefore x = x_1^{-3}$; $dx = -3x_1^{-4}dx_1$; $x^{\frac{4}{3}} = x_1^{-4}$;

$$\therefore -\frac{dy_1}{y_1^2} - \frac{3}{y_1^2}x_1^{-4}dx_1 = -3a^2 dx_1,$$

$$dy_1 - 3a^2 y_1^2 dx_1 = -3x_1^{-4}dx_1.$$

Let $-3a^2 = b_1$, $-3 = a_1$; $\therefore dy_1 + b_1 y_1^2 dx_1 = a_1 x_1^{-4} dx_1$.

Now let $y_1 = \frac{1}{b_1 x_1} + \frac{z_1}{x_1^2}$.

Then $\frac{dz_1}{dx_1} + b_1 z_1^2 \frac{1}{x_1^2} = a_1 x_1^{-2}$, or $x_1^2 \frac{dz_1}{dx_1} = (a_1 - b_1 z^2)$;

$$\therefore \frac{1}{x_1^2}\frac{dx_1}{dz_1} = \frac{1}{a_1 - b_1 z_1^2} = \frac{1}{3(a^2 z_1^2 - 1)},$$

$$\frac{3a}{x_1} = \log\sqrt{\frac{az_1 + 1}{az_1 - 1}} \times \frac{1}{c} = \log\frac{1}{c}\sqrt{\frac{3a^2 x_1^2 y_1 + x_1 + 3a}{3a^2 x_1^2 y_1 + x_1 - 3a}};$$

$$\therefore \text{since } \frac{1}{x_1} = x^{\frac{1}{3}} \text{ and } y_1 = \frac{1}{y};$$

$$\therefore e^{6ax^{\frac{1}{3}}} = \frac{1}{c^2}\left\{\frac{3a^2 x^{-\frac{2}{3}} + y(3a + x^{-\frac{1}{3}})}{3a^2 x^{-\frac{2}{3}} + y(x^{-\frac{1}{3}} - 3a)}\right\}$$

$$= \frac{1}{c^2}\left\{\frac{3a^2 x^{-\frac{1}{3}} + y(1 + 3ax^{\frac{1}{3}})}{3a^2 x^{-\frac{1}{3}} + y(1 - 3ax^{\frac{1}{3}})}\right\};$$

$$\therefore c^2 = C = e^{-6ax^{\frac{1}{3}}}\left\{\frac{3a^2 x^{-\frac{1}{3}} + y(1 + 3ax^{\frac{1}{3}})}{3a^2 x^{-\frac{1}{3}} + y(1 - 3ax^{\frac{1}{3}})}\right\}.$$

Ex. 2. Let $dy + y^2 dx = \frac{a^2 dx}{x^{\frac{8}{3}}}$.

Here $-\frac{8}{3}$ is of the form $\frac{-4n}{2n-1}$;

$\therefore$ let $y = \frac{1}{x} + \frac{z}{x^2}$; and $\because b = 1$, $m = -\frac{8}{3}$;

$\therefore \frac{dz}{dx} + \frac{bz^2}{x^2} = ax^{m+2}$ becomes $\frac{dz}{dx} + z^2\frac{1}{x^2} = a^2 x^{-\frac{2}{3}}$.

Let $x = \frac{1}{y_1}$, and $\because m + 2 = -\frac{2}{3}$; $\therefore m + 1 = \frac{1}{3}$.

Let $x^{\frac{1}{3}} = x_1$; $\therefore x^{-\frac{2}{3}}dx = 3dx_1$; $\frac{1}{x} = \frac{1}{x_1^3}$; $\therefore \frac{dx}{x^2} = \frac{3dx_1}{x_1^4}$;

$\therefore \frac{-dy_1}{y_1^2} + \frac{1}{y_1^2}\frac{3dx_1}{x_1^4} = 3a^2dx_1$; $\therefore dy_1 + 3a^2y_1^2dx_1 = 3x_1^{-4}dx_1$,

which, as has been shewn, is integrable.

110. It sometimes happens that equations in which the variables are separated admit of algebraical integrals, although the integral of each part is transcendental.

Thus, since $\frac{dy}{\sqrt{1-y^2}} + \frac{dx}{\sqrt{1-x^2}} = 0$, (1);

$$\therefore \sin^{-1}x + \sin^{-1}y = C = \sin^{-1}c.$$

$$\therefore x\sqrt{1-y^2} + y\sqrt{1-x^2} = c;$$

we may however obtain the same result, thus

$$(1) \times xy; \quad \therefore \int \frac{xydx}{\sqrt{1-x^2}} + \int \frac{xydy}{\sqrt{1-y^2}} = 0;$$

$$\text{or } -y\sqrt{1-x^2} + \int dy\sqrt{1-x^2} - x\sqrt{1-y^2} + \int dx\sqrt{1-x^2} = -c;$$

which since $dy\sqrt{1-x^2} + dx\sqrt{1-y^2} = 0$, reduces itself to

$$y\sqrt{1-x^2} + x\sqrt{1-y^2} = c,$$

the required algebraic relation between y and x.

111. Again, let $\frac{dy}{\sqrt{a+by+cy^2}} + \frac{dx}{\sqrt{a+bx+cx^3}} = 0$;

make y and x functions of another variable t, so that

$$\frac{dy}{dt} = \sqrt{a+by+cy^2}; \quad \therefore \frac{dx}{dt} = -\sqrt{a+bx+cx^2};$$

let $x + y = p$, then squaring and differentiating,

$$2\frac{d^2y}{dt^2} = b + 2cy; \quad 2\frac{d^2x}{dt^2} = b + 2cx;$$

$$\therefore 2\frac{d^2y}{dt^2} + 2\frac{d^2x}{dt^2} = 2\frac{d^2p}{dt^2} = 2b + 2cp.$$

Multiply both sides by $\frac{dp}{dt}$, and integrate;

$$\therefore \frac{dp^2}{dt^2} = C + 2bp + cp^2; \quad \therefore \frac{dp}{dt} = \frac{dx}{dt} + \frac{dy}{dt} = \sqrt{C + 2bp + cp^2};$$

$$\text{or } \sqrt{a+bx+cx^2} - \sqrt{a+by+cy^2} = \sqrt{C+2b(x+y)+c(x+y)^2}.$$

Ex. 2. $$\frac{dy}{\sqrt{a+by+cy^2+ey^3+fy^4}} + \frac{dx}{\sqrt{a+bx+cx^2+ex^3+fx^4}} = 0.$$

Let $$\frac{dx}{dt} = \sqrt{a + bx + cx^2 + ex^3 + fx^4} = \sqrt{X};$$

$$\therefore \frac{dy}{dt} = -\sqrt{a + by + cy^2 + ey^3 + fy^4} = \sqrt{Y}.$$

Make $x + y = p$; $x - y = q$; whence squaring and differentiating,

$$\frac{d^2x}{dt^2} + \frac{d^2y}{dt^2} = \frac{d^2p}{dt^2} = \tfrac{1}{2}\left(\frac{dX}{dx} + \frac{dY}{dy}\right)$$

$$= b + c(x + y) + \tfrac{3}{2}e(x^2 + y^2) + 2f(x^3 + y^3)$$

$$= b + cp + \tfrac{3}{4}e(p^2 + q^2) + \tfrac{1}{2}fp(p^2 + 3q^2),$$

$$\frac{dx^2}{dt^2} - \frac{dy^2}{dt^2} = \frac{dp \cdot dq}{dt^2} = bq + cpq + \frac{e}{4}(3p^2q + q^3) + \frac{f}{2}qp(p^2 + q^2);$$

$$\therefore q\frac{d^2p}{dt^2} - \frac{dp \cdot dq}{dt^2} = \frac{e}{2}q^3 + fpq^3;$$

$\therefore$ multiplying both sides by $\frac{2}{q^3} \cdot \frac{dp}{dt}$;

$$\frac{2}{q^3} \cdot \frac{d^2p}{dt^2} \cdot \frac{dp}{dt} - \frac{1}{q^3} \cdot \frac{dp^2}{dt^2} \cdot 2\frac{dq}{dt} = e\frac{dp}{dt} + 2fp \cdot \frac{dp}{dt};$$

whence integrating

$$\frac{1}{q^2}\frac{dp^2}{dt^2} = C + ep + fp^2; \quad \therefore \frac{dp}{dt} = q\sqrt{C + ep + fp^2};$$

or $\because \frac{dp}{dt} = \frac{dx}{dt} + \frac{dy}{dt}$, $q = x - y$, and $p = x + y$,

$$\sqrt{a + bx + cx^2 + ex^3 + fx^4} - \sqrt{a + by + cy^2 + ey^3 + fy^4}$$

$$= (x - y)\sqrt{c + e(x + y) + f(x + y)^2},$$

an algebraic equation, which may be put also under a rational form: for writing $(x-y)\sqrt{P}$ instead of the right-hand side of the equation, inverting and multiplying by $X-Y$;

$$\therefore \frac{X-Y}{\sqrt{X}-\sqrt{Y}} = \frac{X-Y}{(x-y)\sqrt{P}};$$

$$\text{or } \sqrt{X}+\sqrt{Y} = \frac{X-Y}{(x-y)\sqrt{P}}; \quad \therefore \text{ by addition,}$$

$$2\sqrt{X} = (x-y)\sqrt{P} + \frac{X-Y}{(x-y)\sqrt{P}} = \frac{(x-y)^2 P + (X-Y)}{(x-y)\sqrt{P}};$$

and squaring both sides, the equation becomes rational.

Ex. 3. $$\frac{d\phi}{\sqrt{1-e^2\sin^2\phi}} + \frac{d\theta}{\sqrt{1-e^2\sin^2\theta}} = 0.$$

Let $$\frac{d\phi}{dt} = \sqrt{1-e^2\sin^2\phi}; \quad \therefore \frac{d\theta}{dt} = -\sqrt{1-e^2\sin^2\theta};$$

$$\therefore \frac{d^2\phi}{dt^2} = -e^2\sin\phi\,.\cos\phi; \quad \frac{d^2\theta}{dt^2} = -e^2\sin\theta\,.\cos\theta;$$

$$\therefore \frac{d^2\phi}{dt^2} + \frac{d^2\theta}{dt^2} = -\frac{e^2}{2}(\sin 2\phi + \sin 2\theta),$$

$$\frac{d^2\phi}{dt^2} - \frac{d^2\theta}{dt^2} = -\frac{e^2}{2}(\sin 2\phi - \sin 2\theta);$$

$\therefore$ by making $p = \phi + \theta$, $q = \phi - \theta$,

$$\frac{d^2p}{dt^2} = -\frac{e^2}{2}\cdot\{\sin(p+q) + \sin(p-q)\} = -e^2\sin p\,.\cos q,$$

$$\frac{d^2q}{dt^2} = -\frac{e^2}{2}\cdot\{\sin(p+q) - \sin(p-q)\} = -e^2.\sin q\,.\cos p.$$

$$\text{And } \frac{dp}{dt}\frac{dq}{dt} = \frac{d\phi^2}{dt^2} - \frac{d\theta^2}{dt^2} = -e^2(\sin^2\phi - \sin^2\theta)$$

$$= \frac{e^2}{2}\cdot(\cos 2\phi - \cos 2\theta) = \frac{e^2}{2}\cdot\{\cos(p+q) - \cos(p-q)\}$$

$$= -e^2\sin p\sin q;$$

$$\therefore \frac{d^2p}{dt^2} \div \frac{dp}{dt}\cdot\frac{dq}{dt} = \frac{\cos q}{\sin q}, \quad \frac{d^2q}{dt^2} \div \frac{dp}{dt}\frac{dq}{dt} = \frac{\cos p}{\sin p};$$

$$\therefore \frac{d.\left(\frac{dp}{dt}\right)}{\frac{dp}{dt}} = \frac{\cos q}{\sin q} \cdot \frac{dq}{dt}, \quad \frac{d.\left(\frac{dq}{dt}\right)}{\frac{dq}{dt}} = \frac{\cos p}{\sin p} \cdot \frac{dp}{dt};$$

$$\therefore \log\left(\frac{dp}{dt}\right) = \log \sin q + c' = \log \sin q + \log a = \log (a \sin q);$$

$$\therefore \frac{dp}{dt} = a \sin q; \quad \text{also } \frac{dq}{dt} = a' \sin p;$$

$$\therefore \sqrt{1 - e^2 \sin^2 \phi} - \sqrt{1 - e^2 \sin^2 \theta} = a \sin (\phi - \theta);$$

$$\text{and } \sqrt{1 - e^2 \sin^2 \phi} + \sqrt{1 - e^2 \sin^2 \theta} = a' \cdot \sin (\phi + \theta).$$

Cor. 1. The constants a and a' have a mutual dependence;

$$\text{for } \because \frac{dp}{dt} \cdot \frac{dq}{dt} = -e^2 \sin p \cos p = a a' \sin p \cdot \cos p;$$

$$\therefore a a' = -e^2.$$

Cor. 2. The preceding equation may be put under a simple form;

$$\text{for } \because \frac{dp}{dq} = \frac{a \sin q}{a' \sin p}; \quad \therefore a \cos q = a' \cos p + a'' \ldots\ldots (1).$$

But a, a', a'' are reducible to one constant; for if μ be the value of ϕ when $\theta = 0$,

$$a = \frac{-1 + \sqrt{1 - e^2 \sin^2 \mu}}{\sin \mu}, \quad a' = \frac{1 + \sqrt{1 - e^2 \sin^2 \mu}}{\sin \mu},$$

$$a'' = a \cos q - a' \cos p = (a - a') \cos \mu = -\frac{2 \cos \mu}{\sin \mu}.$$

Substituting in (1) for a, a', a'', we have

$$\cos (\phi - \theta) \left\{-1 - \sqrt{1 - e^2 \sin^2 \mu}\right\}$$

$$= \cos (\phi + \theta) \left\{1 + \sqrt{1 - e^2 \sin^2 \mu}\right\} - 2 \cos \mu;$$

$$\therefore \cos (\phi - \theta) + \cos (\phi + \theta)$$

$$+ \left\{\cos (\phi + \theta) - \cos (\phi - \theta)\right\} \sqrt{1 - e^2 \sin^2 \mu} = 2 \cos \mu;$$

$$\text{or } \cos \phi \cdot \cos \theta - \sin \phi \cdot \sin \theta \sqrt{1 - e^2 \sin^2 \mu} = \cos \mu.$$

Cor. 3. If $\int \frac{d\phi}{\sqrt{1-e^2\sin^2\phi}} = f(\phi)$,

and $\therefore \int \frac{d\theta}{\sqrt{1-e^2\sin^2\theta}} = f(\theta)$;

$\therefore f(\phi)+f(\theta) = \text{a constant} = \beta.$

But if $\phi = \mu$, $\theta = 0$, and $f(\theta) = 0$; $\therefore \beta = f(\mu)$;

$\therefore f(\phi)+f(\theta) = f(\mu).$

Integration of Differential Equations of the first Order and of the n^{th} Degree.

112. Let $\left(\frac{dy}{dx}\right)^n + P\left(\frac{dy}{dx}\right)^{n-1} + Q\left(\frac{dy}{dx}\right)^{n-2} + \&c. + U = 0$

be the equation; P, Q, &c. and U, being rational functions of x and y.

Let the equation be solved with regard to $\frac{dy}{dx}$; and let X_1, X_2, X_3, &c. be the values of $\frac{dy}{dx}$, or p thus found; then each of the equations $p = X_1$, $p = X_2$, $p = X_3$, &c. when integrated will satisfy the proposed equation, as also will the equation formed of the product of all these integrals.

Since the differential equation arises from eliminating a single constant*, raised to the n^{th} power, from the primitive equation; and since each simple integral introduces a constant, the solution will contain n constants, and therefore be more general than that from which it is derived. But if we consider that the constants are arbitrary, we may make each, equal to the constant belonging to the primitive equation, and then the result will be of the required form.

Ex. 1. Let $\frac{dy^2}{dx^2} = a^2$; $\therefore \frac{dy}{dx} = a$, and $\frac{dy}{dx} = -a$;

$\therefore y = ax + c$, and $y = -ax + c'$,

either of which satisfies the equation. Also their product

$$(y - ax - c)(y + ax - c') = 0$$

will satisfy it.

* For suppose $y - cx + c^2 = 0$; $\therefore p = c$; $\therefore y - px + p^2 = 0$, an equation of the first order and of the second degree.

For differentiating we obtain

$$\left(\frac{dy}{dx} - a\right)(y + ax - c') + \left(\frac{dy}{dx} + a\right)(y - ax - c) = 0,$$

and making successively $y = ax + c$, and $y = -ax + c'$, we get the results $\frac{dy}{dx} = a$; $\frac{dy}{dx} = -a$; as we ought.

Again from the original equation, since $\frac{dy}{dx} = \pm a$; $\therefore$ $y - c = \pm ax$, and squaring both sides, $(y - c)^2 = a^2x^2$.

This equation gives two lines, inclined at different directions to the axis of x, but both cutting the axis of y in the same point; and by giving to (c) different values, we may have groups of such lines in pairs. And the integral $(y - ax + c)(y + ax - c')$ gives the same result, except that each factor represents only lines inclined in the same direction; but by giving to c and c' all possible values, and taking care to collect together those straight lines in which c and c' are equal, we shall find the solutions comprised in the equation $(y - c)^2 = a^2x^2$, which is limited to the single constant c.

Ex. 2. Let $\frac{dy^2}{dx^2} = ax$, or $p = \pm\sqrt{ax}$;

$$\therefore \frac{dy}{dx} = \sqrt{ax}, \text{ and } \frac{dy}{dx} = -\sqrt{ax};$$

$$\therefore y = \frac{2}{3}\sqrt{a}x^{\frac{3}{2}} + c, \text{ and } y = -\frac{2}{3}\sqrt{a}x^{\frac{3}{2}} + c',$$

each of which is comprised in $(y - c)^2 = \frac{4}{9}ax^3$.

Ex. 3. Find the curve when $s = ax + by$.

$$\text{Here } \frac{ds}{dx} = \sqrt{1 + \frac{dy^2}{dx^2}} = a + b\frac{dy}{dx}.$$

And $\because \frac{dy}{dx}$ is obviously constant, let $\frac{dy}{dx} = m$;

$\therefore y = mx + c$, the equation to a straight line;

$$\therefore \frac{y - c}{x} = m, \text{ and } \sqrt{1 + \left(\frac{y - c}{x}\right)^2} = a + b\left(\frac{y - c}{x}\right).$$

Ex. 4. $p^2y + 2px = y$; $y^2 = 2ax + a^2$.

113. When the equation only involves x and p, and can be solved with regard to x, we proceed thus:

$$\text{Since } x = f(p) = P, \text{ and } \frac{dy}{dx} = p;$$

$$\therefore y = px - \int_p x = pP - \int_p P;$$

whence y is a function of p, and therefore of x.

Ex. 1. Let $x + ap = b\sqrt{1+p^2}$;

$$\therefore y = -ap^2 + bp\sqrt{1+p^2} - \int_p (-ap + b\sqrt{1+p^2})$$

$$= -\frac{ap^2}{2} + \frac{bp}{2}\sqrt{1+p^2} - \frac{b}{2}\int_p (p + \sqrt{1+p^2}) + c.$$

The elimination of p will give y in terms of x.

Ex. 2. Let $(1+p^2)x = 1$; $\therefore x = \dfrac{1}{1+p^2}$, and $p = \sqrt{\dfrac{1-x}{x}}$;

$$\therefore y = px - \int_p \frac{1}{1+p^2} = px - \tan^{-1} p + C$$

$$= \sqrt{x - x^2} - \tan^{-1}\sqrt{\frac{1-x}{x}} + C.$$

Ex. 3. Let $x\dfrac{dy}{dx} = \sqrt{1 + \dfrac{dy^2}{dx^2}}$;

$$\therefore px = \sqrt{1+p^2}; \quad \therefore x = \frac{\sqrt{1+p^2}}{p};$$

$$\therefore y = px - \int_p x = px - \int_p \frac{\sqrt{1+p^2}}{p}$$

$$= px - \int_p \frac{1}{p\sqrt{1+p^2}} - \int_p \frac{p}{\sqrt{1+p^2}}$$

$$= px - \log\left(\frac{cp}{1+\sqrt{1+p^2}}\right) - \sqrt{1+p^2};$$

$$\therefore y = \log\left(\frac{1+\sqrt{1+p^2}}{cp}\right) = \log\left(\frac{x+\sqrt{x^2-1}}{c}\right).$$

Ex. 4. $y = a\sqrt{1+p^2}$; $x + c = a\log(y + \sqrt{y^2 - a^2})$.

114. When the differential equation contains y, x and p, and is homogeneous with respect to y, the variables can be

separated by making $y = zx$; for then x will disappear, and we shall have $z = f(p)$.

But $\because y = xz$; $\therefore p - z = x\dfrac{dz}{dx}$; $\therefore \dfrac{1}{x}\dfrac{dx}{dz} = \dfrac{1}{p-z}$;

$$\therefore \frac{1}{x}\frac{dx}{dp} = \frac{\dfrac{dz}{dp}}{p - f(p)} = \frac{f'(p)}{p - f(p)}.$$

And x being found a function of p; $y = px - \int x\dfrac{dp}{dx}$, y may be determined in terms of p and therefore in terms of x.

Ex. Let $y - px = x\sqrt{1 + p^2}$.

Make $y = xz$; $\therefore z - p = \sqrt{1 + p^2}$;

$$\therefore z^2 - 2zp = 1, \text{ or } p = \frac{z}{2} - \frac{1}{2z} = z + x\frac{dz}{dx};$$

$$\therefore x\frac{dz}{dx} = -\frac{z}{2} - \frac{1}{2z} = -\frac{1}{2}\cdot\frac{z^2+1}{z};$$

$$\therefore \frac{dx}{xdz} = -\frac{2z}{1+z^2};$$

$$\therefore \log\left(\frac{x}{2c}\right) = \log\frac{1}{1+z^2} = \log\frac{x^2}{x^2+y^2};$$

$$\therefore x^2 + y^2 - 2cx = 0,$$

the equation to a circle, the origin being in the circumference. This is the solution of the problem; Find the curve in which the perpendicular from the origin upon the tangent is equal to the abscissa.

115. *Integration of the equation, called Clairaut's Formula.*

$$y = px + f(p) = px + P.$$

Differentiate, when we have

$$\frac{dy}{dx} = p + x\frac{dp}{dx} + \frac{dP}{dx};$$

$$\therefore \text{since } \frac{dy}{dx} = p, \text{ and } \frac{dP}{dx} = P'\frac{dp}{dx}, \text{ we have}$$

$$0 = (x + P')\frac{dp}{dx}; \quad \therefore \frac{dp}{dx} = 0, \text{ or } x + P' = 0.$$

$$\text{If we make } \frac{dp}{dx} = 0; \quad p = c; \quad \therefore\ y = cx + c'.$$

This equation appears to have two arbitrary constants; but if c be put for p in the original equation, and C for P, C being what P becomes when c is substituted for p, then $y = cx + C$; $\therefore\ C = c'$, and the equation has but one arbitrary constant. This is the general solution.

Again, from $x + P' = 0$, a value of p will be obtained which is a function of x or y, and does not introduce into the original equation the constant by the elimination of which the differential equation was formed; such a solution of the equation is called a singular or particular solution.

The particular value may be derived from the general solution, by making c to vary; and as $y = cx + C$ is the equation to a straight line, the particular solution gives the equation to the curve which is the locus of the intersections of the straight lines denoted by the general solution.

Ex. 1. $y - px = a\sqrt{1 + p^2}$;

$$\therefore\ \frac{dy}{dx} - p - x\frac{dp}{dx} = \frac{ap}{\sqrt{1 + p^2}}\frac{dp}{dx};$$

$$\therefore\ \left\{x + \frac{ap}{\sqrt{1 + p^2}}\right\} = 0, \text{ and } \frac{dp}{dx} = 0;$$

$$\therefore\ p = c, \text{ and } y = cx + a\sqrt{1 + c^2},$$

which is the general solution.

$$\text{But } x = \frac{-ap}{\sqrt{1 + p^2}}; \quad \therefore \frac{a^2}{x^2} = \frac{1 + p^2}{p^2}; \quad \therefore \frac{\sqrt{a^2 - x^2}}{x} = \frac{1}{p};$$

$$\therefore\ p = \frac{x}{\sqrt{a^2 - x^2}}; \quad \sqrt{1 + p^2} = \frac{-ap}{x} = \frac{-a}{\sqrt{a^2 - x^2}};$$

$$\therefore\ y = \frac{x^2}{\sqrt{a^2 - x^2}} - \frac{a^2}{\sqrt{a^2 - x^2}} = -\frac{a^2 - x^2}{\sqrt{a^2 - x^2}}$$

$$= -\sqrt{a^2 - x^2}; \quad \therefore\ y^2 + x^2 = a^2,$$

which is the solution of the problem: "Find the curve in which each of the perpendiculars drawn from a given point upon the tangent is equal to a given line."

Ex. 2. Let $y = px + \frac{a}{p}(1 + p^2)$; $\therefore\ y^2 = 4a(a + x)$.

Ex. 3. $y = px + a\sqrt[3]{1+p^3}$; $y = cx + a\sqrt[3]{1+c^3}$.

Ex. 4. Let $(y - px)^m = a^{m-1}\left(\frac{y}{p} - x\right)$;

$$\therefore \left(\frac{y}{m}\right)^m = \left(\frac{a}{m-1}\right)^{m-1} x;$$

this is the curve in which $AD^m = a^{m-1}AT$.

Ex. 5. Find the equation to the curve, when the rectangle contained by the perpendiculars on the tangent, one from the origin and the other from a point in the axis of x, at a distance $2c$ from the origin shall equal a given quantity b^2.

$$\text{Here } \frac{y - px}{\sqrt{1+p^2}} \times \frac{y + (2c - x)p}{\sqrt{1+p^2}} = b^2,$$

whence if $b^2 + c^2 = a^2$; $y^2 = \frac{b^2}{a^2}\{a^2 - (c-x)^2\}$.

Prob. Find the equation to the curve PQR which cuts any number of ellipses APB, AQB, &c. described upon a common major-axis AB, so that the areas APN, AQM, &c. shall be of constant magnitude, PN, QM being ordinates.

Let $AN = x$, $NP = y$, $AB = 2a$, m^2 the given area, b the axis minor of one of the ellipses, but which varies as we pass from one ellipse to the other;

$$\therefore y = \frac{b}{a}\sqrt{2ax - x^2}; \quad \therefore m^2 = \frac{1}{a}\int_x b\sqrt{2ax - x^2}.$$

Now differentiating with regard both to b and x,

$$\frac{b}{a}\sqrt{2ax - x^2} + \frac{1}{a}\cdot\frac{db}{dx}\int_x \sqrt{2ax - x^2} = 0;$$

$$\therefore \text{since } \int_x b\sqrt{2ax - x^2} = b\int_x \sqrt{2ax - x^2} = m^2 a:$$

for $\int_x$ refers only to a particular curve, for which $\therefore b$ is constant;

$$\therefore \frac{b}{a}\sqrt{2ax - x^2} + \frac{db}{dx}\cdot\frac{m^2}{b} = 0,$$

$$\text{or } \sqrt{2ax - x^2} + \frac{db}{dx}\cdot\frac{m^2 a}{b^2} = 0;$$

$$\therefore \int_x \sqrt{2ax - x^2} - \frac{m^2 a}{b} = C,$$

$$\text{or } \int_x \sqrt{2ax - x^2} - \frac{m^2\sqrt{2ax - x^2}}{y} = C.$$

Now when $x = 0$, $\int_x \sqrt{2ax - x^2} = 0$, it being a circular area; also since the area APM is of constant magnitude, y must be infinitely great when x is indefinitely small; therefore since $-m^2\sqrt{2ax - x^2} = y \times C$, C must $= 0$, and hence the equation to the required curve is

$$y = \frac{m^2\sqrt{2ax - x^2}}{\int_x \sqrt{2ax - x^2}}.$$

Integration of Differential Equations of the second and higher Orders.

116. The integration of differential equations of the higher orders is effected only in a few instances. We shall begin with the most simple.

To integrate $\frac{d^n y}{dx^n} = X$, X being a function of x.

(1). Let $\frac{d^2 y}{dx^2} = X$; $\therefore \frac{d}{dx}\left(\frac{dy}{dx}\right) = X$;

$$\therefore \frac{dy}{dx} = \int_x X; \quad \therefore y = \int_x \int_x X.$$

(2). Let $\frac{d^3 y}{dx^3} = X$;

$$\therefore \frac{d^2 y}{dx^2} = \int_x X; \quad \frac{dy}{dx} = \int_x \int_x X; \quad \therefore y = \int_x \int_x \int_x X,$$

and so on; the constants have been omitted.

Next to integrate $\frac{d^2 y}{dx^2} = Y$.

Let $\frac{dy}{dx} = p$; $\therefore \frac{d^2 y}{dx^2} = \frac{dp}{dx} = \frac{dp}{dy}\frac{dy}{dx} = p\frac{dp}{dy}$;

$$\therefore p\frac{dp}{dy} = Y, \text{ and } \frac{p^2}{2} = C + \int_y Y.$$

Ex. 1. Let $\frac{d^3 y}{dx^3} = x^3$; make $\frac{dy}{dx} = p$, $\frac{d^2 y}{dx^2} = q$;

$$\therefore \frac{dq}{dx} = x^3; \quad \therefore q = \frac{x^4}{4} + C = \frac{dp}{dx};$$

$$\therefore p = \frac{x^5}{4.5} + cx + c'; \text{ and } y = \frac{x^6}{4.5.6} + \frac{cx^2}{2} + c'x + c''.$$

Ex. 2. Integrate $\frac{d^2y}{dx^2} = \frac{1}{\sqrt{ay}}$.

$$\therefore \frac{dp}{dx} = p\frac{dp}{dy} = \frac{1}{\sqrt{ay}};$$

$$\therefore \frac{p^2}{2} = 2\sqrt{\frac{y}{a}} + c = 2\frac{\sqrt{y} + \sqrt{b}}{\sqrt{a}} \text{ by substitution};$$

$$\therefore \frac{dx}{dy} = \frac{\sqrt[4]{a}}{\sqrt{\sqrt{y} + \sqrt{b}}};$$

which may be integrated by making $\sqrt{y} + \sqrt{b} = z$.

117. To integrate equations involving p and q, put $\frac{dp}{dx}$ for q, and the equations will be transformed to those of the first order.

Ex. 1. Find the curve in which the radius of curvature is inversely as the abscissa.

$$\text{Here } \frac{1 + \left(\frac{dy^2}{dx^2}\right)}{-\frac{d^2y}{dx^2}} = \frac{a^2}{2x};$$

$$\text{or } \because \frac{dy}{dx} = p; \quad \frac{d^2y}{dx^2} = \frac{dp}{dx}; \quad \therefore \frac{\frac{dp}{dx}}{(1+p^2)^{\frac{3}{2}}} = -\frac{2x}{a^2};$$

$$\therefore \frac{p}{\sqrt{1+p^2}} = C - \frac{x^2}{a^2} = \frac{b^2 - x^2}{a^2};$$

$$\therefore \frac{1}{p^2} = -1 + \frac{a^4}{(b^2-x^2)^2} = \frac{a^4 - (b^2-x^2)^2}{(b^2-x^2)^2};$$

$$\therefore p = \frac{b^2 - x^2}{\sqrt{a^4 - (b^2-x^2)^2}}.$$

Ex. 2. Find the same when radius of curvature $= \frac{x^2}{a}$.

$$\therefore \frac{\frac{dp}{dx}}{(1+p^2)^{\frac{3}{2}}} = -\frac{a}{x^2}.$$

$$\frac{p}{\sqrt{1+p^2}} = \frac{a}{x} + c = \frac{a + cx}{x};$$

$$\therefore \frac{1}{p^2} = \frac{x^2 - (a+cx)^2}{(a+cx)^2};$$

$$\therefore p = \frac{a+cx}{\sqrt{x^2 - (a+cx)^2}}.$$

Ex. 3. $\dfrac{d^2y}{dx^2} = \sqrt{1+\dfrac{dy^2}{dx^2}}$; $x = \log\{c(y-c_1) + \sqrt{(y-c_1)^2-1}\}$.

Ex. 4. Integrate $\dfrac{d^2y}{dx^2} = g + m\dfrac{dy^2}{dx^2}$.

Let $\dfrac{dy}{dx} = p$; $\therefore \dfrac{dp}{dx} = g + mp^2$;

$$\therefore \frac{dx}{dp} = \frac{1}{g+mp^2}; \text{ and } \frac{dy}{dp} = \frac{p}{g+mp^2};$$

which are integrable, and p may be eliminated.

Ex. 5. Integrate $\dfrac{d^2y}{dx^2} = Y + m\dfrac{dy^2}{dx^2}$, or $\dfrac{dp}{dx} = Y + mp^2$.

But $\because \dfrac{dp}{dx} = p\dfrac{dp}{dy}$; $\therefore p\dfrac{dp}{dy} = Y + mp^2$;

whence if $p^2 = 2z$; $\dfrac{dz}{dy} - 2mz = Y$;

a linear equation of the first order and of the first degree.

This equation is used to find the velocity of a body moving along a circular arc in a resisting medium.

118. To integrate the equation

$$\frac{d^2y}{dx^2} + P\frac{dy}{dx} + Qy = 0.$$

Make $y = e^{\int u}$; $\therefore \dfrac{dy}{dx} = ue^{\int u}$; $\dfrac{d^2y}{dx^2} = \left(\dfrac{du}{dx} + u^2\right)e^{\int u}$;

$$\therefore e^{\int u}\left\{u^2 + Pu + Q + \frac{du}{dx}\right\} = 0;$$

whence $u^2 + Pu + Q + \dfrac{du}{dx} = 0$,

an equation of the first degree and order; but which is seldom integrable when P and Q are functions of x. It however is, when P and Q are constant; let $P = A$; $Q = B$;

$$\therefore \frac{du}{dx} + u^2 + Au + B = 0;$$

$$\text{or } \frac{du}{dx} + (u-a)(u-b) = 0;$$

which is satisfied by making $u = a$ and $u = b$;

$$\therefore y = e^{\int a} = e^{ax+c'} = c_1 e^{ax},$$

$$\text{and } y = e^{\int b} = e^{bx+c''} = c_2 e^{bx};$$

either of these values substituted for y will satisfy the conditions of the differential equation; but the complete solution, which must comprise two constants, is $y = c_1 e^{ax} + c_2 e^{bx}$; which by substitution we find also satisfies it.

Cor. 1. If the roots of $u^2 + Au + B = 0$ be impossible,

$$a = \alpha + \beta\sqrt{-1}, \text{ and } b = \alpha - \beta\sqrt{-1};$$

$$\therefore y = e^{\alpha x}\left\{c_1 e^{\beta x\sqrt{-1}} + c_2 e^{-\beta x\sqrt{-1}}\right\}$$

$$= e^{\alpha x}\left\{(c_1 + c_2)\cos\beta x + (c_1 - c_2)\sqrt{-1}\sin\beta x\right\}.$$

Make $c_1 + c_2 = A\sin\delta$, $(c_1 - c_2)\sqrt{-1} = A\cos\delta$;

$$\therefore y = Ae^{\alpha x}\left\{\sin\delta\cos\beta x + \cos\delta\sin\beta x\right\} = Ae^{\alpha x}\sin(\beta x + \delta).$$

Cor. 2. Let the roots be equal; or $a = b$.

Then $y = e^{ax}(c_1 + c_2) = c_1 e^{ax}$ which has but one constant. To find the second constant.

Suppose $b = a + h$; $\therefore y = c_1 e^{ax} + c_2 e^{ax+hx}$

$$= e^{ax}\left\{c_1 + c_2 e^{hx}\right\} = e^{ax}\left\{c_1 + c_2 + c_2 hx + \frac{c_2 h^2 x^2}{1.2} + \&c.\right\};$$

make $c_1 + c_2 = c'$, $c_2 h = c''$, and $h = 0$;

$$\therefore y = e^{ax}(c' + c''x).$$

119. The equation $\frac{d^2y}{dx^2} + P\frac{dy}{dx} + Qy = 0$,

is seldom integrable when P and Q are functions of x; it can however be solved when $P = \frac{A}{a+bx}$; and $Q = \frac{B}{(a+bx)^2}$.

For make $a + bx = e^{bz}$;

$$\therefore \frac{dz}{dx} = \frac{1}{a+bx}; \quad \therefore \frac{dy}{dx} = \frac{dy}{dz}\frac{dz}{dx} = \frac{dy}{dz}\frac{1}{a+bx};$$

$$\therefore \frac{d^2y}{dx^2} = \frac{d^2y}{dz^2}\frac{dz}{dx}\frac{1}{a+bx} - \frac{dy}{dz}\frac{b}{(a+bx)^2}$$

$$= \left(\frac{d^2y}{dz^2} - b\frac{dy}{dz}\right)\frac{1}{(a+bx)^2};$$

whence by substitution, and multiplying by $(a+bx)^2$,

$$\frac{d^2y}{dz^2} + (A-b)\frac{dy}{dz} + By = 0;$$

which may be integrated by the preceding methods.

120. To integrate the general equation

$$\frac{d^ny}{dx^n} + A\frac{d^{n-1}y}{dx^{n-1}} + B\frac{d^{n-2}y}{dx^{n-2}} + \&c. + Ly = 0;$$

where A, B, C, &c. L, are constant.

Let $y = e^{mx}$; $\therefore \frac{dy}{dx} = me^{mx}$; $\frac{d^2y}{dx^2} = m^2e^{mx}$, &c.

$$\therefore m^n + Am^{n-1} + Bm^{n-2} + Cm^{n-3} + \&c. + L = 0.$$

Let a, b, c, &c. be the roots of this equation; then

$$y = e^{ax},\ y = e^{bx},\ y = e^{cx};\ \&c.$$

will be particular integrals of the general equation, and the substitution of each in it will satisfy it. Hence the complete integral will be, by the introduction of n constants,

$$y = c_1e^{ax} + c_2e^{bx} + c_3e^{cx} + \&c.$$

Cor. 1. Should any of the roots be equal, as $a = b$; then for $c_1e^{ax} + c\,e^{bx}$, put $e^{ax}(c_1 + c_2x)$;

$$\therefore y = e^{ax}(c_1 + c_2x) + c_3e^{cx} + \&c.$$

And if three roots be equal, and a be the equal root, put

$$e^{ax}(c_1 + c_2x + c_3x^2) \text{ for } c_1e^{ax} + c_2e^{bx} + c_3e^{cx},$$

and so on for any number of equal roots.

Cor. 2. If pairs of roots be impossible, substitute for the impossible exponential functions, the cosines and sines of the circular arcs, to which they are equivalent.

Ex. 1. $\frac{d^2u}{d\theta^2} + n^2u = 0.$

Let $u = e^{m\theta}$; $\therefore \frac{du}{d\theta} = me^{m\theta}$; $\frac{d^2u}{d\theta^2} = m^2e^{m\theta}$;

$\therefore\ m^2e^{m\theta} + n^2e^{m\theta} = 0\ ;\ \therefore\ m^2 + n^2 = 0,$ and $m = \pm n\sqrt{-1}$;

$$\therefore\ u = c'e^{n\theta\sqrt{-1}} + c''e^{-n\theta\sqrt{-1}}$$
$$= (c' + c'')\cos n\theta + (c' - c'')\sqrt{-1}\sin n\theta$$
$$= A\cos(n\theta + B).$$

If $c' + c'' = A\cos B,$ and $(c' - c'')\sqrt{-1} = -A\sin B.$

Ex. 2. $\dfrac{d^2u}{d\theta^2} + n^2u + \alpha^2 = 0.$

Make $\alpha^2 = n^2\beta,$ and $u + \beta = w$;

$$\therefore\ \frac{d^2w}{d\theta^2} + n^2w = 0\ ;\quad \therefore\ u = -\beta + A\cos(n\theta + B).$$

Ex. 3. $\dfrac{d^2s}{dt^2} + 2k\dfrac{ds}{dt} + fs = 0.$

Make $s = e^{mt}\ ;\quad \therefore\ m^2 + 2km + f = 0\ ;$

$$\therefore\ m = -k \pm \sqrt{-1}\sqrt{f - k^2} = -k \pm \alpha\sqrt{-1}\ ;$$

$$\therefore\ s = e^{-kt}(c'e^{\alpha t\sqrt{-1}} + c''e^{-\alpha t\sqrt{-1}}) = Ae^{-kt}\cos(\alpha t + B).$$

Examples (1) and (2) are useful in *Physical Astronomy*; Ex. (3) gives the space a function of the time, when a body moves through the arc of a cycloid, the resistance varying as the velocity.

Ex. 4. $\dfrac{d^3y}{dx^3} - 6\dfrac{d^2y}{dx^2} + 11\dfrac{dy}{dx} - 6y = 0.$

Let $y = e^{mx}\ ;\quad \therefore\ m^3 - 6m^2 + 11m - 6 = 0\ ;$

$$\therefore\ y = c_1e^x + c_2e^{2x} + c_3e^{3x}.$$

Ex. 5. Let $\dfrac{d^3y}{dx^3} - 3\dfrac{d^2y}{dx^2} + 3\dfrac{dy}{dx} - y = 0.$

$$\therefore\ y = e^x(c_1 + c_2x + c_3x^2).$$

Ex. 6. $\dfrac{d^2y}{dx^2} + 8\dfrac{dy}{dx} + 16y = 0\ ;\quad \therefore\ y = e^{-4x}(c_1 + c_2x).$

Ex. 7. $\dfrac{d^2y}{dx^2} - 6\dfrac{dy}{dx} + 34y = 0\ ;\quad \therefore\ y = Ae^{3x}\cos(B + 5x).$

Ex. 8. $\dfrac{d^2y}{dx^2} - \dfrac{1}{x}\dfrac{dy}{dx} + \dfrac{y}{x^2} = 0.$ Let $x = e^z$;

$$\therefore\ y = e^z(c_1 + c_2z) = x(c_1 + c_2\log x).$$

Ex. 9. $\dfrac{d^2y}{dx^2} + \dfrac{1}{x}\dfrac{dy}{dx} - \dfrac{y}{x^2} = 0$;

$$\therefore\ y = c_1e^x + c_2e^{-x} = c_1x + \frac{c_2}{x}.$$

Ex. 10. Integrate $\dfrac{d^4y}{dx^4} - a^4y = 0.$

$$\therefore\ y = c_1e^{ax} + c_2e^{-ax} + A\cos(B + ax).$$

Ex. 11. Integrate $\dfrac{d^4y}{dx^4} + a^4y = 0.$

Here $m^4 + a^4 = 0$; $\dfrac{m}{a} = \dfrac{1 \pm \sqrt{-1}}{\sqrt{2}}$, and $= \dfrac{-1 \pm \sqrt{-1}}{\sqrt{2}}$;

$$\therefore\ y = Ae^{\frac{ax}{\sqrt{2}}}\cos\left(B + \frac{ax}{\sqrt{2}}\right) + A_1e^{-\frac{ax}{\sqrt{2}}}\cos\left(B_1 + \frac{ax}{\sqrt{2}}\right).$$

Ex. 12. Integrate $\dfrac{d^4y}{dx^4} = \dfrac{1}{a^2}\dfrac{d^2y}{dx^2}$.

$$\therefore\ y = c_1e^{\frac{x}{a}} + c_2e^{-\frac{x}{a}} + c_3 + c_4x.$$

Ex. 13. Integrate $\dfrac{d^ny}{dx^n} - y = 0.$

Make $y = e^{mx}$; $\therefore\ m^n - 1 = 0$,

let 1, a_1, a_2, a_3, a_4, &c. a_{n-1}, be the roots of this equation;

$$\therefore\ y = c_1e^x + c_2e^{a_1x} + c_3e^{a_2x} + \&c. + c_ne^{a_{n-1}x}.$$

121. To solve the equation,

$$\frac{d^2y}{dx^2} + P\frac{dy}{dx} + Qy = R.....(1).$$

We shall shew that the solution of this equation may be made to depend upon that of the equation,

$$\frac{d^2y}{dx^2} + P\frac{dy}{dx} + Qy = 0.....(2).$$

To effect this, we proceed to apply to this equation, a method called by Lagrange, "The Variation of the Parameters;" which consists in this, that if $y = c'y_1 + c''y_2$ be the solution of the equation (2), we may assume it to be that of equation (1), if c' and c'' be considered functions of x.

Let $\therefore\ y = c'y_1 + c''y_2$ be the solution of (1);

$$\therefore \frac{dy}{dx} = c'\frac{dy_1}{dx} + c''\frac{dy_2}{dx} + y_1\frac{dc'}{dx} + y_2\frac{dc''}{dx}.$$

But as we have made but one supposition to determine c' and c'', we may make another; let therefore

$$y_1\frac{dc'}{dx} + y_2\frac{dc''}{dx} = 0; \quad \therefore \frac{dy}{dx} = c'\frac{dy_1}{dx} + c''\frac{dy_2}{dx};$$

$$\therefore \frac{d^2y}{dx^2} = c'\frac{d^2y_1}{dx^2} + c''\frac{d^2y_2}{dx^2} + \frac{dc'}{dx}\frac{dy_1}{dx} + \frac{dc''}{dx}\frac{dy_2}{dx};$$

whence by substitution in the original equation (1),

$$c'\left(\frac{d^2y_1}{dx^2} + P\frac{dy_1}{dx} + Qy_1\right) + c''\left(\frac{d^2y_2}{dx} + P\frac{dy_2}{dx} + Qy_2\right)$$

$$+ \frac{dc'}{dx}\frac{dy_1}{dx} + \frac{dc''}{dx}\frac{dy_2}{dx} = R,$$

which by means of equation (2) is reduced to

$$\frac{dc'}{dx}\frac{dy_1}{dx} + \frac{dc''}{dx}\frac{dy_2}{dx} = R,$$

$$\text{or } \because \frac{dc''}{dx} = -\frac{y_1}{y_2}\frac{dc'}{dx},$$

$$\frac{dc'}{dx}\left(\frac{dy_1}{dx} - \frac{y_1}{y_2}\frac{dy_2}{dx}\right) = R;$$

whence $\dfrac{dc_1}{dx}$ is found to be a function of x, and $c' = X_1 + C_1$,

also similarly $c'' = X_2 + C_2$;

$$\therefore y = C_1y_1 + C_2y_2 + y_1X_1 + y_2X_2.$$

A similar proof applies to equations of a higher order.

Ex. 1. Integrate $\dfrac{d^2y}{dx^2} + a^2y = \cos\beta x.$

The solution of the equation $\dfrac{d^2y}{dx^2} + a^2y = 0$ is

$$y = c'\cos ax + c''\sin ax;$$

let this be the solution of the proposed equation;

$$\therefore \frac{dy}{dx} = -c'a\sin ax + c''a\cos ax + \frac{dc'}{dx}\cos ax + \frac{dc''}{dx}\sin ax$$

$$= -c'a\sin ax + c''a\cos ax.$$

$$\text{Since } \frac{dc'}{dx}\cos ax + \frac{dc''}{dx}\sin ax = 0;$$

$$\therefore \frac{d^2y}{dx^2} = -c'a^2\cos ax - c''a^2\sin ax - a\frac{dc'}{dx}\sin ax + a\frac{dc''}{dx}\cos ax$$

$$= -a^2y - a\frac{dc'}{dx}\sin ax + a\frac{dc''}{dx}\cos ax;$$

$$\therefore -a\frac{dc'}{dx}\sin ax + a\frac{dc''}{dx}\cos ax = \cos \beta x \ldots\ldots(1).$$

$$\text{But } \frac{dc''}{dx} = -\frac{\cos ax}{\sin ax}\frac{dc'}{dx};$$

$$\therefore -a\frac{dc'}{dx}\left(\sin ax + \frac{\cos^2 ax}{\sin ax}\right) = \cos \beta x;$$

$$\therefore \frac{dc'}{dx} = -\frac{1}{a}\cos \beta x \sin ax = -\frac{1}{2a}\{\sin(a+\beta)x + \sin(a-\beta)x\},$$

$$\text{and } \frac{dc''}{dx} = \frac{1}{a}\cos \beta x \cos ax = \frac{1}{2a}\{\cos(a+\beta)x + \cos(a-\beta)x\};$$

$$\therefore c' = c_1 + \frac{1}{2a}\left\{\frac{\cos(a+\beta)x}{a+\beta} + \frac{\cos(a-\beta)x}{a-\beta}\right\},$$

$$c'' = c_2 + \frac{1}{2a}\left\{\frac{\sin(a+\beta)x}{a+\beta} + \frac{\sin(a-\beta)x}{a-\beta}\right\};$$

$$\therefore y = c_1\cos ax + c_2\sin ax + \frac{1}{2a}\left(\frac{\cos \beta x}{a+\beta} + \frac{\cos \beta x}{a-\beta}\right)$$

$$= c_1\cos ax + c_2\sin ax + \frac{\cos \beta x}{a^2-\beta^2}.$$

Ex. 2. Integrate $\frac{d^2y}{dx^2} + a^2y = X$.

Let $y = c'\cos ax + c''\sin ax$, be the solution. Proceeding as in Example 1,

$$\therefore \frac{dc'}{dx}\sin ax - \frac{dc''}{dx}\cos ax = -\frac{X}{a}.$$

$$\text{And } \frac{dc''}{dx} = -\frac{dc'}{dx}\frac{\cos ax}{\sin ax};$$

$$\therefore \frac{dc'}{dx} = -\frac{1}{a}X\sin ax, \text{ and } \frac{dc''}{dx} = \frac{1}{a}X\cos ax;$$

$$\therefore\ c' = c_1 - \frac{1}{a}\int_x X \sin ax,$$

$$c'' = c_2 + \frac{1}{a}\int_x X \cos ax;$$

$$\therefore\ y = c_1 \cos ax + c_2 \sin ax - \frac{\cos ax}{a}\int_x X \sin ax + \frac{\sin ax}{a}\int_x X \cos ax.$$

Ex. 3. Integrate $\frac{d^2y}{dx^2} + A\frac{dy}{dx} + By = X;\ X = f(x)$.

Let a and b be the roots of the equation $m^2 + Am + B = 0$;

$\therefore$ let $y = c'e^{ax} + c''e^{bx}$ be the solution of the equation;

$$\therefore\ \frac{dy}{dx} = ac'e^{ax} + bc''e^{bx} + e^{ax}\frac{dc'}{dx} + e^{bx}\frac{dc''}{dx}.$$

$$\text{Make } e^{ax}\frac{dc'}{dx} + e^{bx}\frac{dc''}{dx} = 0;$$

$$\therefore\ \frac{dy}{dx} = ac'e^{ax} + bc''e^{bx}.$$

$$\frac{d^2y}{dx^2} = a^2c'e^{ax} + b^2c''e^{bx} + ae^{ax}\frac{dc'}{dx} + be^{bx}\frac{dc''}{dx};$$

$$\therefore\ c'e^{ax}(a^2 + Aa + B) + c''e^{bx}(b^2 + Ab + B) + ae^{ax}\frac{dc'}{dx} + be^{bx}\frac{dc''}{dx} = X.$$

And $a^2 + Aa + B = 0;\ b^2 + Ab + B = 0;$

$$\therefore\ ae^{ax}\frac{dc'}{dx} + be^{bx}\frac{dc''}{dx} = X.$$

But $be^{ax}\frac{dc'}{dx} + be^{bx}\frac{dc''}{dx} = 0;$

$$\therefore\ (a-b)e^{ax}\frac{dc'}{dx} = X, \text{ and } -(a-b)e^{bx}\frac{dc''}{dx} = X;$$

$$\therefore\ c' = c_1 + \frac{1}{a-b}\int_x Xe^{-ax},\ c'' = c_2 - \frac{1}{a-b}\int_x Xe^{-bx};$$

$$\therefore\ y = c_1e^{ax} + c_2e^{bx} + \frac{e^{ax}}{a-b}\int_x Xe^{-ax} - \frac{e^{bx}}{a-b}\int_x Xe^{-bx}.$$

Ex. 4. Integrate $\frac{d^2y}{dx^2} - 5\frac{dy}{dx} + 6y = x.$

$$\therefore y = c_1 e^{2x} + c_2 e^{3x} + \frac{1}{6}\left(x + \frac{5}{6}\right).$$

SIMULTANEOUS DIFFERENTIAL EQUATIONS.

122. In the applications of the Differential Calculus to physical problems, mutually dependent equations are frequently found in which $n + 1$ variables are involved, and n equations are given: as most commonly the unknown quantities are x, y, and t; x and y being functions of t; we shall first solve the system of equations which involve these quantities. The method of solution is due to D'Alembert.

Let $A\frac{dx}{dt} + B\frac{dy}{dt} + Cx + Dy = \theta,$

and $A_1\frac{dx}{dt} + B_1\frac{dy}{dt} + C_1x + D_1y = \theta_1,$

be the two equations: A, B, &c. being constant, θ and θ_1 functions of t. By the successive elimination of $\frac{dx}{dt}$ and $\frac{dy}{dt}$ these may be reduced to the form,

$$\frac{dx}{dt} + ax + by = T \ (1), \quad \frac{dy}{dt} + a_1x + b_1y = T_1 \ (2).$$

Now multiply (2) by m and add the product to (1);

$$\therefore \frac{d}{dt}(x + my) + (a + ma_1)\left(x + \frac{b + mb_1}{a + ma_1}y\right) = T + mT_1.$$

Let $\frac{b + mb_1}{a + ma_1} = m$: and let m_1, m_2 be the two values of m resulting from the equation; also let $a + m_1a_1 = r_1$; $a + m_2a_1 = r_2$; $\therefore$ we shall have the two linear equations of the first order,

$$\frac{d}{dt}(x + m_1y) + r_1(x + m_1y) = T + m_1T_1;$$

$$\frac{d}{dt}(x + m_2y) + r_2(x + m_2y) = T + m_2T_1;$$

$$\therefore x + m_1y = e^{-r_1t}\left\{\int_t e^{r_1t}(T + m_1T_1) + C\right\},$$

$$x + m_2y = e^{-r_2t}\left\{\int_t e^{r_2t}(T + m_2T_1) + C_1\right\}.$$

Ex. Let $\frac{dx}{dt} + 4y + 5x = e^t$; $\frac{dy}{dt} + x + 2y = e^{2t}$;

$$\therefore \frac{d}{dt}(x + my) + (4 + 2m)y + (5 + m)x = e^t + me^{2t}.$$

Let $\frac{4 + 2m}{5 + m} = m$; $\therefore m = 1$ or -4, $5 + m = 6$ or 1;

$$\therefore x + y = e^{-6t}\left\{\int_t e^{6t}(e^t + e^{2t}) + C\right\} = \tfrac{1}{7}e^t + \tfrac{1}{8}e^{2t} + Ce^{-6t},$$

or $x - 4y = e^{-t}\left\{\int_t e^t(e^t + e^{2t}) + C_1\right\} = \frac{1}{2}e^t + \frac{1}{3}e^{2t} + C_1e^{-t}$;

which give x and y in terms of t.

123. Next to integrate the simultaneous equations,

$$\frac{dx}{dt} + (Ax + By + Cz) = T \quad (1),$$

$$\frac{dy}{dt} + (A_1x + B_1y + C_1z) = T_1 \quad (2),$$

$$\frac{dz}{dt} + (A_2x + B_2y + C_2z) = T_2 \quad (3):$$

where A, B, C, &c. are constant, and T, T_1, T_2 functions of t: multiply (2) by m and (3) by m' and add;

$$\therefore \frac{d}{dt}(x + my + m'z) + (A + A_1m + A_2m')\left\{x + B'y + C'z\right\} = U,$$

$$\text{where } B' = \frac{B + mB_1 + m'B_2}{A + A_1m + A_2m'}, \quad C' = \frac{C + mC_1 + m'C_2}{A + mA_1 + m'A_2},$$

$$U = T + T_1m + T_2m';$$

$$\therefore \text{if } B' = m;\ C' = m';\ A + A_1m + A_2m' = M;\ x + my + m'z = v;$$

$$\frac{dv}{dt} + Mv = U, \text{ a linear equation,}$$

which integrated will give the relation between v and t: also since from $B' = m$, and $C' = m$, two cubic equations will arise if m_1, m_2, m_3; m_1', m_2', m_3' be their roots, and if U_1, U_2, U_3 be the values of the right-hand side of the equation when integrated;

$$\therefore x + m_1y + m_1'z = U_1,$$

$$x + m_2y + m_2'z = U_2,$$

$$x + m_3y + m_3'z = U_3.$$

124. To integrate the simultaneous equation of the second order.

$$\frac{d^2x}{dt^2} + ax + by + c = 0 \quad (1),$$

$$\frac{d^2y}{dt^2} + a_1x + b_1y + c_1 = 0 \quad (2);$$

multiply (2) by m and add,

$$\therefore \frac{d^2}{dt^2}(x + my + c) + (a + ma_1)\left\{x + \frac{b + mb_1}{a + ma_1}y + \frac{c + mc_1}{a + ma_1}\right\} = 0.$$

Make $m = \dfrac{b + b_1m}{a + ma_1}$; $u = x + my + c$; $a + a_1m = -n^2$;

$$\therefore \frac{d^2u}{dt^2} - n^2u = 0; \quad \therefore u = Ce^{nt} + C_1e^{-nt};$$

therefore if m_1 and m_2 be the two values of m,

$$(a + a_1m_1)(x + m_1y) + c + m_1c_1 = (a + m_1a_1)\{c_1e^{nt} + c_2e^{-nt}\},$$

$$(a + a_1m_2)(x + m_2y) + c + m_2c_1 = (a + m_2a_1)\{c_1'e^{nt} + c_2'e^{-nt}\}.$$

Ex. $\dfrac{d^2x}{dt^2} = 3x + 4y - 3$: $\dfrac{d^2y}{dt^2} = 8y - x - 5$;

$$\therefore x = \tfrac{1}{7} + 4c_1e^{2t} + 4c_2e^{-2t} - 3c_1'e^{t\sqrt{7}} - c_2'e^{t\sqrt{7}},$$

$$y = \tfrac{9}{14} + c_1e^{2t} + c_2e^{-2t} - c_1'e^{t\sqrt{7}} - c_2'e^{-t\sqrt{7}}.$$

Differential Equations containing more than two Variables.

125. Equations of this description, of the first order, which, we in general shall suppose, involve the three variables x, y, z, may be divided into (1) Total Differential Equations, (2) Partial Differential Equations.

Total Differential Equations.

Let $du = Pdx + Qdy + Rdz$ be the equation, which may be supposed to arise from the differentiation of

$$u = f(xyz); \text{ whence } P = \frac{du}{dx}; \quad Q = \frac{du}{dy}; \quad R = \frac{du}{dz};$$

and since when this is the case,

$$\frac{dP}{dy} = \frac{dQ}{dx}; \quad \frac{dP}{dz} = \frac{dR}{dx}; \text{ and } \frac{dQ}{dz} = \frac{dR}{dy};$$

we can always ascertain when an equation is a total differential.

Ex. Let $du = \frac{y}{a-z}dx + \frac{x}{a-z}dy + \frac{xy}{(a-z)^2}dz.$

Here $\frac{dP}{dy} = \frac{1}{a-z} = \frac{dQ}{dx}$, $\frac{dP}{dz} = \frac{y}{(a-z)^2} = \frac{dR}{dx}$;

$$\therefore \frac{du}{dx} = \frac{y}{a-z}; \quad \therefore u = \frac{xy}{a-z} + f(yz);$$

$$\therefore \frac{du}{dy}.dy + \frac{du}{dz}dz$$

$$= \frac{x}{a-z}dy + \frac{xy}{(a-z)^2}dz + \frac{d}{dy}f(yz)\,dy + \frac{d}{dz}f(yz)\,dz;$$

$$\therefore \frac{d}{dy}f(yz)\,dy + \frac{d}{dz}f(yz)\,dz = 0;$$

$$\therefore f(yz) = C;$$

$$\therefore u = \frac{xy}{a-z} + C.$$

126. Next to integrate the equation $Pdx + Qdy + Rdz = 0$, which may be put under the form $dz = -\frac{P}{R}dx - \frac{Q}{R}dy$; or which, by making $p = -\frac{P}{R}$, $q = -\frac{Q}{R}$,

may be written $dz = pdx + qdy$.

Now if this equation can be expressed by an equation

$$z = f(x, y, c), \text{ or } f(x, y, z) = c,$$

we ought to have $\frac{d(p)}{dy} = \frac{d(q)}{dx}$,

$$\text{or } \frac{dp}{dy} + \frac{dp}{dz}.\frac{dz}{dy} = \frac{dq}{dx} + \frac{dq}{dz}.\frac{dz}{dx},$$

or $\because \frac{dz}{dy} = q$, and $\frac{dz}{dx} = p$, we have

$$\frac{dp}{dy} - \frac{dq}{dx} + q\frac{dp}{dz} - p\frac{dq}{dz} = 0 \ldots (1),$$

an equation of condition, by which we can ascertain whether

the proposed equation admits of the solution $f(x, y, z) = c$. If we restore the values of p and q the equation becomes

$$P\left(\frac{dQ}{dz} - \frac{dR}{dy}\right) + Q.\left(\frac{dR}{dx} - \frac{dP}{dz}\right) + R.\left(\frac{dP}{dy} - \frac{dQ}{dx}\right) = 0 \ldots (2);$$

when this equation or the preceding one holds, one of the variables must be considered as constant, and the remaining part of the equation integrated according to the rules given for the integration of functions of two variables.

Ex. 1. $(y + z)\,dx + (x + y)\,dz + (z + x)\,dy = 0.$

Here $P = y + z$, $\frac{dP}{dy} = 1 = \frac{dP}{dz}$, $Q = x + z$, $\frac{dQ}{dx} = \frac{dQ}{dz} = 1$;

$R = x + y$; $\frac{dR}{dx} = 1 = \frac{dR}{dy}$; $\therefore$ equation (2) is satisfied;

$\therefore$ making $dz = 0$; $\frac{dx}{x + z} + \frac{dy}{y + z} = 0$;

$\therefore \log(x + z) + \log(y + z) = \phi(z) = \log(Z)$;

$\therefore (x + z).(y + z) = Z$;

$\therefore (y + z)\,dx + (x + z)\,dy + (x + y + 2z)\,dz = dZ$;

$\therefore \frac{dZ}{dz} = 2z$; $\therefore Z = z^2 + C$; $\therefore xy + xz + yz = C.$

Ex. 2. $(ay - bz)\,dx + (cz - ax)\,dy + (bx - cy)\,dz = 0.$

Make z constant; $\therefore dz = 0$, then

$$\frac{a\,dx}{cz - ax} + \frac{a\,dy}{ay - bz} = 0; \quad \therefore \log\left(\frac{ay - bz}{cz - ax}\right) = \log Z;$$

$$\therefore \frac{a\,dy}{cz - ax} + \frac{a(ay - bz)\,dx}{(cz - ax)^2} + \frac{a(bx - cy)}{(cz - ax)^2}\,dz = dZ;$$

$$\therefore (cz - ax)\,dy + (ay - bz)\,dx + (bx - cy)\,dz = 0$$

$$= \{(cz - ax)\}^2 dZ;$$

$$\therefore dZ = 0; \quad \therefore Z = C'; \quad \therefore (ay - bz) = C'(cz - ax).$$

127. If the equation $P\,dx + Q\,dy + R\,dz = 0$ is not a complete differential, but may be rendered so by the means of a factor F, the equation (2) must still be satisfied: for, multiplying by F,

$FPdx + FQdy + FRdz$ is an exact differential;

$$\therefore \frac{d.FP}{dy} = \frac{d.FQ}{dx}; \quad \frac{d.FR}{dx} = \frac{d.FP}{dz}; \quad \frac{d.FQ}{dz} = \frac{d.FR}{dy};$$

$$\therefore F\left(\frac{dP}{dy} - \frac{dQ}{dx}\right) + P.\frac{dF}{dy} - Q.\frac{dF}{dx} = 0,$$

$$F\left(\frac{dR}{dx} - \frac{dP}{dz}\right) + R.\frac{dF}{dx} - P.\frac{dF}{dz} = 0,$$

$$F\left(\frac{dQ}{dz} - \frac{dR}{dy}\right) + Q.\frac{dF}{dz} - R.\frac{dF}{dy} = 0.$$

Multiply the first of these equations by R, the second by Q, and the third by P, and add: we have

$$R\left(\frac{dP}{dy} - \frac{dQ}{dx}\right) + Q.\left(\frac{dR}{dx} - \frac{dP}{dz}\right) + P.\left(\frac{dQ}{dz} - \frac{dR}{dy}\right) = 0,$$

the same equation as in the preceding article.

128. When the differentials dx, dy, dz exceed the first degree, it must be solved with respect to dz; and can then only be integrated when the factors of the equation so solved are of the form $dz - pdx - qdy = 0$.

Partial Differential Equations.

129. It is here required to find $z = f(xy)$ from one of the partial differential coefficients, or from some relation existing between them.

To integrate $\frac{dz}{dx} = P$; P being a function of x, y, z: we first integrate it on the supposition that y is constant, and instead of adding an arbitrary constant after the integration we add $\phi(y)$: similarly we add $\phi(x)$, if the equation be $\frac{dz}{dy} = P$.

Ex. 1. $\frac{dz}{dx} = a$; $\therefore z = ax + \phi(y)$.

Ex. 2. $\frac{dz}{dx} = \frac{az}{x}$; $\therefore \log z = \log x^a \phi(y)$; $\therefore z = x^a \phi(y)$.

Ex. 3. $\frac{dz}{dx} = \frac{y^2 + z^2}{x^2 + y^2}$; $\therefore \frac{dz}{y^2 + z^2} - \frac{dx}{x^2 + y^2} = 0$;

$$\therefore \tan^{-1}\frac{z}{y} - \tan^{-1}\frac{x}{y} = \tan^{-1}\phi(y); \quad \frac{y(z-x)}{y^2+xz} = \phi(y).$$

Ex. 4. $\dfrac{dz}{dx} = \dfrac{x^2-y^2}{x^2+y^2}\cdot\dfrac{z}{x}; \quad z = \dfrac{x^2+y^2}{x}\phi(y).$

130. To integrate the equation $Pp + Qq = R$, in which P, Q, R, contain at once, x, y, z;

$$\because dz = pdx + qdy; \quad \therefore p = \frac{dz - qdy}{dx};$$

$$\therefore \text{substituting, } Pdz - Rdx = q.(Pdy - Qdx).$$

Here there are two cases: 1st, $Pdz - Rdx$ may contain only z and x, $Pdy - Qdx$ only y and x; 2nd, either or both of these factors may contain all the variables.

Case 1. Let F be the factor which will make $Pdz - Rdx$ a complete differential dM, and F_1 the factor which will make $Pdy - Qdx = dN$;

$$\therefore dM = \frac{q.F}{F_1}.dN, \text{ which cannot be integrated unless } \frac{qF}{F_1}$$

is a function of N, $= \phi'(N)$; whence $dM = \phi'(N)\,dN$; $\therefore M = \phi(N)$.

Ex. 1. $px + qy = nz$; also $p = \dfrac{dz - qdy}{dx}$;

$$\therefore xdz - nzdx = q(xdy - ydx);$$

and to integrate $xdz - nzdx$, and $xdy - ydx$, we must multiply the former by $\dfrac{1}{x^{n+1}}$, the latter by $\dfrac{1}{x^2}$;

$$\therefore M = \frac{z}{x^n}; \quad N = \frac{y}{x}; \quad \therefore z = x^n\phi\left(\frac{y}{x}\right).$$

Ex. 2. $px + qy = 0$; $\therefore z = \phi\left(\dfrac{y}{x}\right)$.

Ex. 3. $qx - py = 0$; $\therefore z = \phi(x^2 + y^2)$.

Ex. 4. $ap + bq = c$; $\therefore z = \dfrac{cx}{a} + \dfrac{1}{a}\phi.(ay - bx)$.

Ex. 5. $px - q = x^2$; $\therefore z = \dfrac{x^2}{2} + \phi.(y + \log x)$.

Case 2. Next let the variables x, y, z be found in both of the functions $Pdy - Qdx$, and $Pdz - Rdx$; we can no longer

integrate them separately, since z cannot be considered constant in the former, nor x in the latter.

Lagrange observed that if these equations were integrated conjointly: and if we call the integral of the former N: and of the latter M; so that $N = a$, and $M = b$, a and b being arbitrary constants; then the complete integral will be $M = \phi(N)$. But that this method may succeed, one of the equations must involve two of the variables only; and its integral will enable us to eliminate one of the three variables x, y, z, from the remaining equation.

The truth of this proposition may be thus shewn.

Since the equations $N = a$, and $M = b$, are derived from $Pdx - Qdy = 0$, and $Pdz - Rdx = 0$, the differentials of $N = a$ and $M = b$ will be satisfied by the values of dz, dy, deduced from these latter equations: hence differentiating and putting

$$M_x = \frac{dM}{dx}, \quad M_y = \frac{dM}{dy}, \quad M_z = \frac{dM}{dz};$$

$$M_x dx + M_y dy + M_z dz = 0; \quad \text{and} \quad N_x dx + N_y dy + N_z dz = 0.$$

But $\frac{dy}{dx} = \frac{Q}{P}$; $\frac{dz}{dx} = \frac{R}{P}$; $\therefore$ substituting

$$M_x = -M_y \cdot \frac{Q}{P} - M_z \cdot \frac{R}{P}; \quad N_x = -N_y \cdot \frac{Q}{P} - N_z \cdot \frac{R}{P}.$$

But from the equation $M = \phi(N)$;

$$\therefore M_x dx + M_y dy + M_z dz = \phi'(N)\{N_x dx + N_y dy + N_z dz\};$$

hence $M_y \cdot (Pdy - Qdx) + M_z \cdot (Pdz - Rdx)$

$$= \phi'(N)\{N_y (Pdy + Qdx) + N_z (Pdz - Rdx)\};$$

$$\therefore Pdz - Rdx = -\frac{M_y - N_y \phi'(N)}{M_z - M_z \phi'(N)}(Pdy - Qdx) = -\omega(Pdy - Qdx);$$

$$\therefore dz = \frac{R + \omega \cdot Q}{P} dx - \omega \cdot dy;$$

whence $p = \frac{R + \omega Q}{P}$, $q = -\omega$: which substituted in the original equation $Pp + Qq = R$ satisfy it; and therefore the assumption that $M = \phi(N)$ (which is derived from the integration of $Pdy - Qdx = 0$; and $Pdz - Rdx = 0$) is the solution of the problem, is completely justified.

Ex. 1. $px^2 - qxy + y^2 = 0$;

$\therefore\ x^2 dz + y^2 dx = q\,(x^2 dy + yx dx)$;

$\therefore\ x^2 dy + yx dx = 0\,(1)$; and $x^2 dz + y^2 dx = 0\,(2)$:

from (1), $x dy + y dx = 0$; $\therefore\ xy = a = N$; $\therefore\ y = \dfrac{a}{x}$:

from (2), $x^2 dz + \dfrac{a^2}{x^2} dx = 0$; $\therefore\ z - \dfrac{a^2}{3x^3} = b = M$,

or $z - \dfrac{y^2}{3x} = \phi(N)$; $\therefore\ z = \dfrac{y^2}{3x} + \phi(xy)$.

Ex. 2. $px + qy = n\sqrt{x^2 + y^2}$;

$\therefore\ x dz - n\sqrt{x^2 + y^2}\, dx = q\,.\,(x dy - y dx)$:

from $x dy - y dx = 0$; $\dfrac{y}{x} = a = N$; $\therefore\ y = ax$;

$\therefore\ x dz - n\sqrt{x^2 + a^2 x^2}\, dx = 0$; $\therefore\ z - nx\sqrt{1 + a^2} = b = M$;

$\therefore\ z - n\sqrt{x^2 + y^2} = \phi(N)$; $z = n\sqrt{x^2 + y^2} + \phi\left(\dfrac{y}{x}\right)$.

Ex. 3. $px^2 + qy - y^2 = 0$; $z = \dfrac{y^2}{2} + \phi\left(\dfrac{1}{x} + \log y\right)$.

Ex. 4. $qx + py = nz$; $z = (x + y)^n\,.\,\phi(y^2 - x^2)$.

This equation as well as the more general one $Pp + Qq = Rz$, is best solved by making $z = e^u$.

Ex. 5. $px^2 + qy^2 = nxy$; $z = \dfrac{nxy}{x - y} \log\left(\dfrac{x}{y}\right) + \phi\left(\dfrac{y - x}{xy}\right)$.

Ex. 6. $p + q\,.\,\dfrac{y}{x} = x^m z^n$; $\dfrac{1}{(n-1)z^{n-1}} + \dfrac{x^{m+1}}{m+1} = \phi\left(\dfrac{y}{x}\right)$.

Ex. 7. $px + qz + y = 0$; $x e^{\sin^{-1}\cdot \frac{z}{\sqrt{y^2+z^2}}} = \phi(y^2 + z^2)$.

Ex. 8. $(c - z) = (a - x)p + (b - y)q$; $\therefore\ \dfrac{y - b}{z - c} = \phi\left(\dfrac{x - a}{z - c}\right)$.

Ex. 9. $px + qy = \dfrac{xy}{z}$; $z^2 = xy + f\left(\dfrac{y}{x}\right)$.

Ex. 10. $z - px - qy = m\,(x + y + z)$;

$\therefore\ x^{m-1}(x + y + mz) = c^m \phi\left(\dfrac{y}{x}\right)$.

Ex. 11. $(y - bz)\,p - (x - az)\,q = bx - ay$;

$$\therefore\ (y - bz)\,dy + (x - az)\,dx = 0 \ldots (1),$$

$$(y - bz)\,dz - (bx - ay)\,dx = 0 \ldots (2),$$

$$(x - az)\,dz + (bx - ay)\,dy = 0 \ldots (3);$$

$\therefore$ (2) $\times\, a$ − (3) $\times\, b$ and $\div$ by $bx - ay$;

$$\therefore\ dz + a\,dx + b\,dy = 0.$$

(2) $\times\, x - 3 \times y$ and $\div$ by $bx - ay$;

$$\therefore\ x\,dx + y\,dy + z\,dz = 0;$$

$$\therefore\ x^2 + y^2 + z^2 = \phi\,(z + ax + by),$$

this is the general equation to surfaces of revolution.

131. The same method applies to partial differential equations containing a greater number of variables.

Ex. 1. Let $nu + px + qy = az$,

where $n = \dfrac{dz}{du}$ and $z = f(xyu)$:

$$\because\ dz = p\,dx + q\,dy + n\,du;\quad \therefore\ p = \frac{dz - q\,dy - n\,du}{dx};$$

$$\therefore\ n(u\,dx - x\,du) + q(y\,dx - x\,dy) + x\,dz - az\,dx = 0;$$

$$\therefore\ u\,dx - x\,du = 0;\quad \therefore\ \frac{x}{u} = a,$$

$$y\,dx - x\,dy = 0;\quad \therefore\ \frac{x}{y} = \beta,$$

$$x\,dz - az\,dx = 0;\quad \therefore\ \frac{z}{x^a} = \gamma;$$

$$\therefore \text{ since } \gamma = \phi\,(a\beta);\quad z = x^a \cdot \phi\left(\frac{x}{u}, \frac{x}{y}\right).$$

Ex. 2. $ap + bq + cn = 0$; put $n = \dfrac{dz - p\,dx - q\,dy}{du}$;

$$\therefore\ z = \phi\,\{cx - au),\ (cy - bu)\}.$$

132. When the partial differential coefficients p and q exceed the first degree, q must be considered as a function of p, x, y, z, and the values of $\dfrac{dq}{dx}$, $\dfrac{dq}{dz}$ substituted in the

equation $\frac{dp}{dy} - \frac{dq}{dx} + q\frac{dp}{dz} - p\frac{dq}{dz} = 0$, which is derived from $dz = p\,dx + q\,dy$, considered to be a complete differential. This substitution will give an equation which integrated will give a value of p, and $\therefore$ of q, in terms of x, y, z, and a constant a: substitute for p and q in $dz = p\,dx + q\,dy$, and integrate, whence $f(x, y, z, a) = \beta = \phi(a)$; and to eliminate (a) differentiate $b = \phi(a)$ with regard to (a).

Ex. 1. Let $\frac{dz}{dx} \cdot \frac{dz}{dy} = 1$, or $pq = 1$;

$$\therefore\ q = \frac{1}{p};\quad \therefore\ \frac{dq}{dz} = -\frac{1}{p^2}\cdot\frac{dp}{dz};\ \frac{dq}{dx} = -\frac{1}{p^2}\cdot\frac{dp}{dx};$$

$$\therefore \text{ substituting in } \frac{dp}{dy} - \frac{dq}{dx} + q\frac{dp}{dz} - p\frac{dq}{dz} = 0,$$

$$\frac{dp}{dy} + \frac{1}{p^2}\frac{dp}{dx} + \frac{2}{p}\cdot\frac{dp}{dz} = 0;$$

$$\text{or if } \frac{dp}{dy} = q';\ \frac{dp}{dx} = p';\ \frac{dp}{dz} = n';$$

$$q' + \frac{1}{p^2}p' + \frac{2}{p}n' = 0;$$

whence, treating it as an equation of three variables and a function of $(pxyz)$,

$$dp = 0;\quad \therefore\ p = a;\quad q = \frac{1}{a};$$

$$\therefore\ dz - a\,dx - \frac{dy}{a} = 0;\quad \therefore\ z - ax - \frac{y}{a} = (\phi\, a);$$

whence, by differentiation, $\frac{y}{a^2} - x = \phi'(a)$.

Ex. 2. Let $\frac{dz^2}{dx^2} + \frac{dz^2}{dy^2} = 1$; or $p^2 + q^2 = 1$;

$$\therefore\ z - ax - y\sqrt{1 - a^2} = \phi(a),$$

$$x - \frac{ay}{\sqrt{1 - a^2}} = -\phi'(a).$$

Partial Differential Equations of the second Order.

133. Here make $\frac{d^2z}{dx^2} = r$; $\frac{d^2z}{dx\,dy} = s$; $\frac{d^2z}{dy^2} = t$.

To integrate $\frac{d^2z}{dx^2} = M$, M being a function of x and y:

$$\because\ p = \frac{dz}{dx};\quad \therefore\ \frac{dp}{dx} = \frac{d^2z}{dx^2} = M;\quad \therefore\ p = \int_x M + \phi(y);$$

$$\therefore\ z = \int_x . \int_x M + x\phi(y) + f(y).$$

If $\frac{d^2z}{dxdy} = M$: then $\frac{dz}{dx} = \int_y M + \phi'(x)$;

$$\therefore\ z = \int_x \int_y M + \phi(x) + f(y).$$

Ex. 1. Let $\frac{d^2z}{dxdy} = xy$; $\therefore\ p = \frac{xy^2}{2} + \phi'(x)$;

$$\therefore\ z = \frac{x^2y^2}{4} + \phi(x) + f(y).$$

Ex. 2. $\frac{d^2z}{dxdy} = ax + by$; $\therefore z = \frac{ax^2y}{2} + \frac{by^2x}{2} + \phi(x) + f(y)$.

134. Let $\frac{d^2z}{dx^2} = P . \frac{dz}{dx}$, P being a function of x and y;

$$\therefore\ \frac{dp}{dx} = Pp;\quad \therefore\ \log p = \int_x P + \phi(y);$$

$$\therefore\ p = e^{\int_x P + \phi(y)};\quad \therefore\ z = \int_x e^{\int_x P + \phi(y)} + f(y).$$

Ex. $x\frac{d^2z}{dx^2} = (n-1)\frac{dz}{dx}$; $nz = x^n\phi(y) + nf(y)$.

135. To integrate $\frac{d^2z}{dxdy} = P\frac{dz}{dx}$ or $\frac{dp}{dy} = Pp$;

$$\therefore\ z = \int_x e^{\int_x P + \phi(x)} + f(y).$$

136. To integrate $\frac{d^2z}{dx^2} = P\frac{dz}{dx} + Q$, P and Q being functions of x and y.

If $\frac{dz}{dx} = p$; $\therefore\ \frac{dp}{dx} - Pp = Q$, a linear equation;

whence $p = e^u\{\int_x e^{-u}Q + \phi(y)\}$ where $u = \int_x P$;

$$\therefore\ z = \int_x e^u\{\int_x e^{-u}Q + \phi y\} + f(y).$$

Ex. 1. Let $xy\frac{d^2z}{dx^2} = (n-1) . y . \frac{dz}{dx} + a$;

$$\therefore\ z = \frac{x^n \cdot \{\phi(y)\}^{n-1}}{n \cdot (n-1)} - \frac{ax}{(n-1)y} + f(y).$$

Ex. 2. $xy\dfrac{d^2z}{dxdy} = bx\dfrac{dz}{dx} + ay;$

$$\therefore\ z = -\frac{ay\log x}{b-1} + \phi(x) + f(y).$$

137. To integrate $Rr + Ss + Tt = V$, where R, S, T and V are functions of x, y, z, p and q:

$$\because\ p = \frac{dz}{dx};\quad \therefore\ dp = \frac{d^2z}{dx^2}dx + \frac{d^2z}{dydx}dy = rdx + sdy,$$

$$\text{and } q = \frac{dz}{dy};\quad \therefore\ dq = \frac{d^2z}{dy^2}dy + \frac{d^2z}{dxdy}dx = tdy + sdx;$$

$$\therefore\ r = \frac{dp - sdy}{dx};\quad t = \frac{dq - sdx}{dy};\quad \therefore \text{ substituting}$$

$$Rdpdy + Tdqdx - Vdxdy = s(Rdy^2 - Sdxdy + Tdx^2);$$

it is unnecessary to integrate the two members of this equation separately; for if we can integrate one of them so as to have the integral $N = a$, and by combining this with the other arrive at the integral $M = b$, M and N being functions of x, y, z, p and q, we may prove, as in a preceding article, that $M = \phi(N)$: and this result will give an equation with which we must proceed, as with an equation of partial differences of the first order.

Ex. 1. To integrate $\dfrac{d^2z}{dx^2} = c^2 \cdot \dfrac{d^2z}{dy^2}$; or $r = c^2t$:

since $dp = rdx + sdy;\quad dq = tdy + sdx;$

$$\therefore\ dpdy - c^2dqdx = s(dy^2 - c^2dx^2);$$

$$\therefore\ \frac{dy^2}{dx^2} = c^2;\quad \therefore\ \frac{dy}{dx} = \pm c;\quad \therefore\ y - cx = a:\quad y + cx = a_1:$$

$$\text{and } dp \cdot \frac{dy}{dx} - c^2dq = 0;\quad \therefore\ cdp - c^2dq = 0;$$

$$\therefore\ p - cq = b = \phi'(a) = \phi'(y - cx).$$

$$\text{But } p = \frac{dz - qdy}{dx};\quad \therefore dz - \phi'(y - cx)\,dx = q(dy + cdx);$$

$$\therefore\ dy + cdx = 0;\quad \therefore\ y + cx = a,\ y = a - cx;$$

$$\therefore\ dz - \phi'(y - cx)\,dx = dz - \phi'(a - 2cx)\,dx = 0;$$

$$\therefore\ z - \phi(a - 2cx) = \beta = f(a) = f(y + cx);$$

$$\therefore\ z = \phi(y - cx) + f(y + cx).$$

This is the equation of vibrating chords.

Ex. 2. Integrate $x^2 r + y^2 t + 2xys = 0$;

$$\therefore\ x^2\,dp\,dy + y^2\,dq\,dx = s\,.\,(x^2\,dy^2 + y^2\,dx^2 - 2xy\,dx\,dy);$$

$$\therefore\ (x\,dy - y\,dx)^2 = 0;\quad \therefore\ x\,dy - y\,dx = 0;\quad \therefore\ \frac{y}{x} = a = \frac{dy}{dx},$$

$$\text{and } \frac{x^2}{y^2}\,.\,dp\,.\,\frac{dy}{dx} + dq = 0;\quad dp + a\,dq = 0;$$

$$\therefore\ p + aq = b = \phi(a) = \phi\left(\frac{y}{x}\right);\quad \text{and } \because\ p = \frac{dz - q\,dy}{dx};$$

$$dz - dx\,\phi\left(\frac{y}{x}\right) = q(dy - a\,dx) = 0;$$

$$\therefore\ dz - dx\,\phi(a) = 0;\quad \therefore\ z - x\,\phi(a) = f(a);$$

$$\therefore\ z = x\phi\left(\frac{y}{x}\right) + f\left(\frac{y}{x}\right).$$

Ex. 3. Let $q^2 r + p^2 t - 2pqs = 0$; $y = \phi(z) - xf(z)$.

Ex. 4. Let $r + As + Bt + C = 0$;

$$z = -\frac{Cx^2}{2} + \phi(y - mx) + f(y - m_1 x).$$

Ex. 5. $r - a^2 t = xy$;

$$z = \frac{1}{6}x^3 y + f(y + ax) + \phi(y - ax).$$

Ex. 6. $x^2 r - y^2 t = 0$;

$$z = \sqrt{xy}\,.\,f\left(\frac{y}{x}\right) + \phi xy.$$

CHAPTER VIII.

THE CALCULUS OF VARIATIONS.

1. In the problems of Maxima and Minima hitherto solved, the form of the function which possessed the required property has always been given. But there is a class of problems, in which it is not only required to find when there is a maximum or minimum, but also the nature of the function which possesses the property.

Thus if the shortest distance between two given points be required, we must, in order to find the minimum distance, ascertain the nature of the curve that possesses the property; and thus, to continue the illustration, if C and D be the two points, we must select from the curves CPD, CQD, that which is the shortest: in our reasoning we must therefore pass from one curve to another curve, from a point P in one curve to a point Q in another curve: the change from P to Q is called a *variation* of the ordinate PN: the symbol of variation being δ.



Thus if $NP = y$, and if PQ be indefinitely small, the symbols δ of variation, and d of differentiation, differ in this respect; by means of dy we pass from a point P to another P_1 indefinitely near to it, but in the same curve: by δy we transfer P to a point Q in another curve. We have here supposed the variation to be confined to the ordinate y: but $x = AN$ may also vary at the same time that y does.

2. Since $NP = y$; $\therefore N_1P_1 = y + dy$: also $NQ = y + \delta y$;

$$\therefore N_1Q_1 = NQ + d(NQ) = y + \delta y + d(y + \delta y),$$

and $N_1Q_1 = N_1P_1 + \delta(N_1P_1) = y + dy + \delta(y + dy)$;

$$\therefore \delta y + d(y + \delta y) = dy + \delta(y + dy); \quad \therefore d\delta y = \delta dy;$$

or the symbols of variation and differentiation are interchanged.

Cor. Hence also if dy be put for y,

$$\therefore d\delta dy = \delta d^2y; \text{ or } \because d\delta dy = dd\delta y = d^2\delta y;$$

$$\therefore d^2\delta y = \delta d^2y, \text{ and thus } d^n\delta y = \delta d^n y.$$

If the differential of y be taken $= y_1 - y$, and the variation of y be also supposed to be very small, the theorem will admit of the following proof:

$$\delta dy = \delta(y_1 - y) = \delta y_1 - \delta y = d\delta y.$$

3. There is a similar theorem with regard to integration:

For let $\int u = u_1$; $\therefore\ u = du_1$; $\therefore\ \delta u = d\delta u_1$;

$$\therefore\ \int \delta u = \int \delta d u_1 = \int d \delta u_1 = \delta u_1 = \delta \int u.$$

Also $\int^2 \delta u = \int \delta \int u = \delta \iint u = \delta \int^2 u$,

and thus $\int^n \delta u = \delta \int^n u$.

4. From the preceding we may perceive, that variation is only differentiation under a new symbol; and that to find the variation of a function of y, we must put $y + \delta y$ for y, and that term of the expanded function which involves δy will be the variation of the function required: or, what amounts to the same thing, the variation is the differential coefficient of $u = f(y)$ multiplied by δy; thus if

$u = y^n$, $\delta u = n y^{n-1} \delta y$; and if

$u = f(x, y, p, q,$ &c.$)$ p, q, &c. being $\dfrac{dy}{dx}$, $\dfrac{d^2y}{dx^2}$, &c., and

$$\therefore\ du = \frac{du}{dx}dx + \frac{du}{dy}dy + \frac{du}{dp}dp + \frac{du}{dq}dq + \&c.$$

$$= M dx + N dy + P dp + Q dq + \&c.$$

by putting $M = \dfrac{du}{dx}$, $N = \dfrac{du}{dy}$, $P = \dfrac{du}{dp}$, &c.;

$$\therefore\ \delta u = M \delta x + N \delta y + P \delta p + Q \delta q + \&c.$$

5. To find $\delta \int u$, or $\int \delta u$; u being a function of y and x and their differentials; and y and x being dependent upon some other variable as s or t.

$$\therefore\ du = M dx + N d^2x + P d^3x + Q d^4x + \&c.$$

$$+ m dy + n d^2y + p d^3y + q^4 dy + \&c.;$$

therefore $\because\ d^2x = d dx$, $d^3x = d d^2x$;

$$\therefore\ \delta u = M \delta x + N \delta dx + P \delta d^2x + Q \delta d^3x + \&c.$$

$$+ m \delta y + n \delta dy + p \delta d^2y + q \delta d^3y + \&c.$$

$$\therefore\ \int \delta u = \int (M \delta x + N \delta d.x + P \delta d^2x + Q \delta d^3x + \&c.)$$

$$+ \int (m \delta y + n \delta dy + p \delta d^2y + q \delta d^3y + \&c.);$$

$\therefore$ integrating by parts, and placing δ after d,

$$\int N\delta dx = \int Nd\delta x = N\delta x - \int dN\delta x,$$

$$\int P\delta d^2x = \int Pd^2\delta x = Pd\delta x - \int dPd\delta x,$$

$$= Pd\delta x - dP\delta x + \int d^2P\delta x,$$

$$\int Q\delta d^3x = \int Qd^3\delta x = Qd^2\delta x - \int dQd^2\delta x$$

$$= Qd^2\delta x - dQd\delta x + \int d^2Qd\delta x$$

$$= Qd^2\delta x - dQd\delta x + d^2Q\delta x - \int d^3Q\delta x.$$

&c. &c. &c.

Similarly $\int n\delta dy = n\delta y - \int dn\delta y,$

$$\int p\delta d^2y = pd\delta y - dp\delta x + \int d^2p\delta x,$$

$$\int q\delta d^3y = qd^2\delta y - dqd\delta y + d^2q\delta y - \int d^3q\delta y,$$

and substituting these values in $\int\delta u$, we have

$$\int\delta u = (N - dP + d^2Q - \&c.)\,\delta x + (n - dp + d^2q - \&c.)\,\delta y$$
$$+ (P - dQ + \&c.)\,d\delta x + (p - dq + \&c.)\,d\delta y$$
$$+ (Q - dR + \&c.)\,d^2\delta x + (q - dr + \&c.)\,d^2\delta y + \&c.$$
$$+ \int(M - dN + d^2P - d^3Q + \&c.)\,\delta x$$
$$+ \int(m - dn + d^2p - d^3r + \&c.)\,\delta y.$$

6. The result just obtained is composed of two similar parts, one due to the variation of δx, the other due to that of δy; and we thus see that had there been a third variable z, there must be added to the preceding expression, a series of terms similar to that which involves δx.

7. When u is of the form Vdx, to find the variation of $\int Vdx$.

Let $dV = Mdx + Ndy + Pdp + Qdq + Rdr + \&c.$

where $p = \dfrac{dy}{dx}$; $q = \dfrac{d^2y}{dx^2}$; $r = \dfrac{d^3y}{dx^3}$; &c.

and $M = \dfrac{dV}{dx}$; $N = \dfrac{dV}{dy}$; $P = \dfrac{dV}{dp}$; &c.

$$\therefore \delta V = M\delta x + N\delta y + P\delta p + Q\delta q + \&c.$$

Now $\delta\int Vdx = \int\delta . Vdx = \int(V\delta dx + dx . \delta V)$

$$= \int(Vd\delta x + dx\delta V) = \int Vd\delta x + \int dx\delta V$$
$$= V\delta x + \int(dx\delta V - \delta x dV).$$

But $\int(dx\delta V - \delta x dV) = \int dx\,(M\delta x + N\delta y + P\delta p + \&c.)$

$$- \int \delta x\,(M dx + N dy + P dp + \&c.)$$

$$= \int_x N(\delta y - p\delta x) + \int_x P(\delta p - q\delta x) + \int_x Q(\delta q - r\delta x) + \&c.$$

But $\because\ p = \frac{dy}{dx};\ q = \frac{dp}{dx};\ r = \frac{dq}{dx};$

$$\therefore\ \delta p = \frac{dx\delta dy - dy\delta dx}{dx^2} = \frac{\delta dy - p\delta x}{dx};$$

$$\therefore\ \delta p - q\delta x = \frac{\delta dy - p d\delta x - dp\delta x}{dx} = \frac{d}{dx}(\delta y - p\delta x),$$

$$\delta q = \frac{dx\delta dp - dp\delta dx}{dx^2} = \frac{d\delta p - q d\delta x}{dx};$$

$$\therefore\ \delta q - r\delta x = \frac{d\delta p - q d\delta x - dq\delta x}{dx} = \frac{d}{dx}(\delta p - q\delta x).$$

Now let $\delta y - p dx = w$; $\therefore\ \delta p - q\delta x = \frac{dw}{dx}$,

$$\delta q - r dx = \frac{d^2w}{dx^2};\quad \delta r - s dx = \frac{d^3w}{dx^3};$$

$$\therefore\ \delta \int_x V = V\delta x + \int_x N w + \int_x P \cdot \frac{dw}{dx} + \int_x Q \cdot \frac{d^2w}{dx^2} + \&c.$$

But $\int_x P\frac{dw}{dx} = Pw - \int_x \frac{dP}{dx} w,$

$$\int_x Q\frac{d^2w}{dx^2} = Q\frac{dw}{dx} - \int_x \frac{dQ}{dx}\cdot\frac{dw}{dx}$$

$$= Q\frac{dw}{dx} - w\frac{dQ}{dx} + \int w \cdot \frac{d^2Q}{dx^2};$$

and $\int_x R \cdot \frac{d^3w}{dx^3} = R\frac{d^2w}{dx^2} - \frac{dR}{dx}\cdot\frac{dw}{dx} + \frac{d^2R}{dx^2}w - \int w \cdot \frac{d^3R}{dx^3};$

$$\therefore\ \delta \int_x V = V\delta x + \left(P - \frac{dQ}{dx} + \frac{d^2R}{dx^2} - \&c.\right) w$$

$$+ \left(Q - \frac{dR}{dx} + \frac{d^2S}{dx^2} - \&c.\right)\frac{dw}{dx}$$

$$+ \left(R - \frac{dS}{dx} + \&c.\right)\frac{d^2w}{dx^2} + \&c.$$

$$+ \int_x \left(N - \frac{dP}{dx} + \frac{d^2Q}{dx^2} - \frac{d^3R}{dx^3} + \&c.\right) w.$$

8. Thus the variation of $\int_x V$ consists of two distinct parts, one of which is under the sign of integration, and the other is not; the latter is affected only by the variations of the extreme values of y and x, the former is dependent upon all the values between those extreme ones. Let x_1, y_1, x_2, y_2 be the values of x and y at these limits, then the total variation of $\int_x V$

$$= V_2\delta x_2 - V_1\delta x_1 + (P_2 - \frac{dQ_2}{dx_2} + \frac{d^2R_2}{dx_2^2} - \&c.)w_2$$
$$- (P_1 - \frac{dQ_1}{dx_1} + \frac{d^2R_1}{dx_1^2} - \&c.)w_1 + \&c.$$
$$+ \int_x (N - \frac{dP}{dx} + \frac{d^2Q}{dx^2} - \&c.)w.$$

The part under the sign of integration must be taken between the same limits.

9. If $u = f(xyz)$ and x still be the independent variable, then, since the differential of V, may be put

$$dV = Mdx + Ndy + Pdp + Qdq + \&c.$$
$$+ N'dz + P'dp' + Qdq' + \&c.;$$
$$\therefore \delta V = M\delta x + Ndy + Pdp + Q\delta q + \&c.$$
$$+ N'dz + P'dp' + Q'\delta q' + \&c.$$

We shall have for the new variable z, a series of terms, involving N', P', Q', &c. similar to that in which N, P, Q, &c. are introduced; $\therefore$ if $\delta z - p'\delta x = w'$, the total variation will be expressed by

$$\delta\int_x V = V\delta x + (P - \frac{dQ}{dx} + \frac{d^2R}{dx^2} - \&c.)w$$
$$+ (P' - \frac{dQ'}{dx} + \frac{d^2R'}{dx^2} - \&c.)w'$$
$$+ (Q - \frac{dR}{dx} + \&c....)\frac{dw}{dx}$$
$$+ (Q' - \frac{dR'}{dx} + \&c....)\frac{dw'}{dx} + \&c.$$
$$+ \int_x (N - \frac{dP}{dx} + \frac{d^2Q}{dx^2} - \&c.)w$$
$$+ \int_x (N_1 - \frac{dP_1}{dx} + \frac{d^2Q_1}{dx^2} - \&c.)w_1.$$

Maxima and Minima of Integral Formulæ.

10. We proceed to apply the results of the preceding article to the solution of some geometrical problems, involving the lengths and areas of curves, the surfaces and volumes of solids; when these quantities are, within certain limits of the variables, the greatest or least possible.

Now we know that if u, any function of x and y, be a maximum or minimum, $du = 0$; and by the same kind of reasoning which has been used to establish this proposition, it may be shewn that under the same circumstances the variation of u also vanishes; but if $u = V dx$, we have seen that between the limits of x_1, y_1, x_2, y_2,

$$\delta . \int_x V = V_2 \delta x_2 - V_1 \delta x_1 + (P_2 - \frac{dQ_2}{dx_2} + \&c.)\, w_2$$

$$- (P_1 - \frac{dQ_1}{dx_1} + \&c.)\, w_1 + \&c.$$

$$+ \int_x (N - \frac{dP}{dx} + \frac{d^2Q}{dx^2} - \frac{d^3R}{dx^3} + \&c.)\, w.$$

And since when $\int_x V$ is a maximum or minimum, $\delta \int_x V = 0$; therefore the two parts of which the variation of $\int_x V$ is composed must separately be put $= 0$; one part will determine the relations between the co-ordinates of the extreme values of the required function, the other the function which possesses the required maximum or minimum property.

11. Thus from $N - \frac{dP}{dx} + \frac{d^2Q}{dx^2} - \&c. = 0$, and the equation $dV = M dx + N dy + P dp + \&c.$ may be found the curve or function, which is the object of our enquiry, and from $V_2 \delta x_2 - V_1 \delta x_1 + P_2 w_2 - P_1 w_1 \&c. = 0$, the position of its extreme points may be determined; if however the extreme points be fixed, $\delta x_1 = 0$, and $\delta x_2 = 0$, and the latter equation disappears.

12. Thus, if the shortest distance between two given points be required, the former equation will be quite sufficient for the problem; the constants of the integral being determined by the co-ordinates of the given points; but if we wish to find the shortest distance between two given curves the latter equation is also necessary, since it determines the points

in the two curves to which the shortest distance is to be drawn.

Its use may be thus illustrated; let PP_1, QQ_1 be the given curves, and PQ, P_1Q_1 two curves drawn between them, and let P_1Q_1 be derived from PQ by writing $x+\delta x$, $y+\delta y$ for x and y: also let $\frac{dy}{dx}=m$; and $\frac{dy}{dx}=n$ be the equations to PP_1, and QQ_1; then if x_1y_1, x_2y_2 be the co-ordinates of P and Q; $\therefore \frac{\delta y_1}{\delta x_1}=m$; and $\frac{dy_2}{dx_2}=n$; (since the point P is always in PP_1 and Q in QQ_1;) and between these equations and that of the limits, $V_2\delta x_2 - V_1\delta x_1 + \&c. = 0$; two of the quantities, as δy_1 and δy_2 may be eliminated, and the two independent variations, δx_1 and δx_2 will be left, the coefficients of which, being separately put $=0$, will give equations, by which and from the given equations to the curves, the points P and Q, may be determined.



We have here tacitly assumed that $\delta dy_1 = 0$ and $\delta dx_1 = 0$: if this be not the case, some new conditions must be fulfilled by the limits, which will enable us to introduce the higher differentials of the equations of the given curves: by means of which some of the variations δdx, δdy, &c. may be eliminated, and the coefficient of the remaining variations separately put $=0$, and the co-ordinates of the extreme points and the given conditions fulfilled.

13. We shall now deduce from the equation

$$N - \frac{dP}{dx} + \frac{d^2Q}{dx^2} - \&c. = 0,$$

some formulæ of great use in the solution of Problems of maxima and minima.

Let $dV = Mdx + Ndy + Pdp + Qdq + Rdr + \&c.$,

and $N - \frac{dP}{dx} + \frac{d^2Q}{dx^2} - \frac{d^3R}{dx^3} + \&c. = 0.$

(1) Let all but N and $P=0$;

$$\therefore \frac{dV}{dx} = Np + \frac{Pdp}{dx}; \text{ and } N - \frac{dP}{dx} = 0; \quad \therefore N = \frac{dP}{dx};$$

$$\therefore \frac{dV}{dx} = p\frac{dP}{dx} + P\frac{dp}{dx}; \quad \therefore V = Pp + c.$$

(2) Let all but M, N, and P be $= 0$;

$$\therefore \frac{dV}{dx} = M + \frac{d}{dx}(Pp + c); \quad \therefore V = \int_x M + Pp + c;$$

(3) Let $M = 0$, $N = 0$; and all the terms after $Q = 0$;

$$\therefore \frac{dV}{dx} = P\frac{dp}{dx} + Q\frac{dq}{dx}; \quad \frac{dP}{dx} - \frac{d^2Q}{dx^2} = 0; \quad \therefore P = \frac{dQ}{dx} + c;$$

$$\therefore \frac{dV}{dx} = q\frac{dQ}{dx} + Q\frac{dq}{dx} + c\frac{dp}{dx}; \quad \therefore V = Qq + cp + c_1.$$

Cor. If M does not $= 0$; $V = Qq + cp + c_1 + \int_x M$.

Prob. 1. Find the shortest distance between two given points in the same plane.

$$\text{Here } \int_x V = \int_x \sqrt{1 + \frac{dy^2}{dx^2}} = \int_x \sqrt{1 + p^2};$$

$$\therefore V = \sqrt{1 + p^2}; \quad \therefore dV = \frac{p}{\sqrt{1 + p^2}}\,dp;$$

$$\therefore M = 0; \quad N = 0; \quad P = \frac{p}{\sqrt{1 + p^2}}, \quad Q = 0.$$

$$\text{But } N - \frac{dP}{dx} + \&c. = 0; \quad \therefore \frac{dP}{dx} = 0; \quad \therefore P = c = \frac{p}{\sqrt{1 + p^2}};$$

$$\therefore p = \frac{c}{\sqrt{1 - c^2}} = a; \quad \therefore y = ax + b,$$

the equation to a straight line; the constants a, b may be determined by the co-ordinates of the given points.

Prob. 2. Required the curve of quickest descent between two given points.

Let y be vertical and be measured downwards.

$$\text{Then time} = \int_x \frac{\frac{ds}{dx}}{\sqrt{2gy}} = \frac{1}{\sqrt{2g}}\int_x \frac{\sqrt{1 + p^2}}{\sqrt{y}} = \frac{1}{\sqrt{2g}}\int_x V;$$

$$\therefore V = \frac{\sqrt{1 + p^2}}{\sqrt{y}}, \quad dV = -\frac{\sqrt{1 + p^2}}{2y^{\frac{3}{2}}}\,dy + \frac{p}{\sqrt{y}\sqrt{1 + p^2}}\,dp;$$

$$\therefore\ M = 0;\ N = -\frac{\sqrt{1+p^2}}{2y^{\frac{1}{2}}};\ P = \frac{p}{\sqrt{y}\sqrt{1+p^2}},\ Q = 0;$$

$$\therefore\ V = Pp + c;\ \frac{\sqrt{1+p^2}}{\sqrt{y}} = \frac{p^2}{\sqrt{y}\sqrt{1+p^2}} + c;$$

$$\therefore\ \frac{1}{\sqrt{y}\sqrt{1+p^2}} = c = \frac{1}{\sqrt{2a}};\ \therefore\ \sqrt{1+p^2} = \sqrt{\frac{2a}{y}};$$

$$\therefore\ p = \sqrt{\frac{2a-y}{y}}$$ the equation to the cycloid.

PROB. 3. To find the shortest distance between two given curves.

From Prob. (1) $V = \sqrt{1+p^2}$; $p = c$; $y = ax + b$ the equation to the line which is the least distance required.

Let $\frac{dy}{dx} = m$, and $\frac{dy}{dx} = n$, be the equations to the two curves, and $y_1\ x_1$, $y_2\ x_2$ the co-ordinates of the points in which the shortest line intersects them; then since δy_1, δx_1 are the variations of y_1 and x_1 as we pass from one point to another adjacent one in the curve $\frac{dy}{dx} = m$;

$$\frac{\delta y_1}{\delta x_2} = \frac{dy_1}{dx_1} = m;\ \text{and}\ \frac{\delta y_2}{\delta x_2} = \frac{dy_2}{dx_2} = n.$$

But $V_2\delta x_2 - V_1\delta x_1 + P_2 w_2 - P_1 w_1 = 0$,

whence since the variations of the extreme points = 0,

$$V_1\delta x_1 + P_1 w_1 = 0;\quad V_2\delta x_2 + P_2 w_2 = 0;$$

$\therefore\ V_1\delta x_1 + P_1(\delta y_1 - p_1\delta x_1) = 0$ (1); $V_2\delta x_2 + P_2(\delta y_2 - p_2\delta x_2) = 0$ (2);

from (1) $V_1 + P_1 m - P_1 p_1 = 0$; $\therefore\ m = p_1 - \frac{V_1}{P_1} = -\frac{1}{p_1} = -\frac{1}{c}$;

from (2) $V_2 + P_2 n - P_2 p_2 = 0$; $\therefore\ n = p_2 - \frac{V_2}{P_2} = -\frac{1}{p_2} = -\frac{1}{c}$;

$$\therefore\ 1 + cm = 0;\ \text{and}\ 1 + cn = 0;$$

which shew that the line must cut both curves at right angles.

Also the equation to the line being $y - y_1 = \frac{y_2 - y_1}{x_2 - x_1}(x - x_1)$;

$$\therefore\ c = \frac{y_2 - y_1}{x_2 - x_1};$$ whence substituting for c, in $1 + cm = 0$;

and $1 + cn = 0$; we shall have with the equations to the given curves four equations to determine the four quantities, y_1, x_1, y_2, x_2, and thus the line is completely determined.

PROB. 4. Find the curve, which within its own arc, its evolute and radius of curvature, contains the least area.

Here $\frac{dA}{dx} = \frac{R}{2}\frac{ds}{dx} = \frac{(1+p^2)^{\frac{3}{2}}}{-2q}\sqrt{1+p^2} = \frac{(1+p^2)^2}{-2q}$;

$\therefore V = \frac{(1+p^2)^2}{q}$; whence the curve is the cycloid.

PROB. 5. Find the curve when $\int \frac{p^3 y}{\sqrt{1+p^2}}$ is a minimum. This is the solid of least resistance,

$$y = \frac{c(1+p^2)^2}{p^3}; \quad x = c_1 + c\left(\frac{3}{4p^4} + \frac{1+p^2}{p^2} + \log p\right).$$

PROB. 6. Find the curve of quickest descent from one curve to another curve, the velocity being that from a horizontal line.

Here $V = \frac{\sqrt{1+p^2}}{\sqrt{y}}$; $\therefore p = \frac{\sqrt{2a-y}}{y}$, the equation to the cycloid; the equations of the limits give

$$V_1\delta x_1 + P_1\delta y_1 - P_1 p_1 \delta x_1 = 0; \quad V_2\delta x_2 + P_2\delta y_2 - P_2 p_2 \delta x_2 = 0,$$

from which since $\frac{\delta y_1}{\delta x_1} = m$, and $\frac{\delta y_2}{\delta x_2} = n$, we have

$$V_1 - P_1 p_1 + P_1 m = 0; \text{ and } V_2 - P_2 p_2 + P_2 n = 0.$$

But $V_1 - P_1 p_1 = c = \frac{1}{\sqrt{2a}}$, $P = \frac{p}{\sqrt{y}\sqrt{1+p^2}} = \frac{p}{\sqrt{2a}}$;

$$\therefore \frac{1}{\sqrt{2a}} + \frac{p_1 m}{\sqrt{2a}} = 0; \text{ and } \frac{1}{\sqrt{2a}} + \frac{p_2 n}{\sqrt{2a}} = 0;$$

$$\therefore 1 + p_1 m = 0; \text{ and } 1 + p_2 n = 0,$$

which equations shew that the cycloid cuts both the curves at right angles, and from $p = \sqrt{\frac{2a-y}{y}}$, we see that the base of the cycloid coincides with the horizontal line from which we have supposed the body to have commenced its motion.

PROB. 7. To find the curve of quickest descent from one given curve to another given curve, the motion commencing from the upper curve.

Let y_1 be the value of the ordinate at the point at which the motion commences, y the value at the end of time t.

$$\text{Then time} = \frac{1}{\sqrt{2g}} \int_x \frac{\sqrt{1+p^2}}{\sqrt{y-y_1}}; \quad \therefore V = \frac{\sqrt{1+p^2}}{\sqrt{y-y_1}}.$$

In this problem the function V involves y_1 one of the variable co-ordinates of the limits; in such a case we must add the term $\delta y_1 \int_x \frac{dV}{dy}$ to the equation of the limits, and then the whole variation of the $\int_x V$ will become

$$\delta \int_x V = V_2 \delta x_2 - V_1 \delta x_1 + \delta y_1 \int_x \frac{dV}{dx_1} + P_2 w_2 - P_1 w_1 + \&c.$$

$$+ \int_x \left(N - \frac{dP}{dx} + \frac{d^2 Q}{dx^2} - \&c. \right) w.$$

Now referring to the problem,

$$dV = -\frac{\sqrt{1+p^2}}{2(y-y_1)^{\frac{3}{2}}} dy + \frac{p\,dp}{\sqrt{y-y_1}\sqrt{1+p^2}} + \frac{\sqrt{1+p^2}}{2(y-y_1)^{\frac{3}{2}}} dy_1;$$

$$\therefore N = -\frac{\sqrt{1+p^2}}{2(y-y_1)^{\frac{3}{2}}}; \quad P = \frac{p}{\sqrt{y-y_1}\sqrt{1+p^2}}; \quad \frac{dV}{dy_1} = \frac{\sqrt{1+p^2}}{2(y-y_1)^{\frac{3}{2}}},$$

whence from the formula $V = Pp + c$,

$$\frac{\sqrt{1+p^2}}{\sqrt{y-y_1}} = \frac{p^2}{\sqrt{y-y_1}\sqrt{1+p^2}}; \quad \therefore \sqrt{y-y_1}\sqrt{1+p^2} = \frac{1}{C} = \sqrt{2a},$$

the equation to a cycloid, the cusp being at the point from which the motion commences.

Next, to find $\int_x \frac{dV}{dy_1}$; $\frac{dV}{dy_1} = -N = -\frac{dP}{dx}$, $\therefore N - \frac{dP}{dx} = 0$;

$$\therefore \int_x \frac{dV}{dy_1} = -P; \quad \therefore \int_{x_1}^{x_2} \frac{dV}{dx} = P_1 - P_2;$$

$$\therefore V_2 \delta x_2 - V_1 \delta x_1 + P_2 w_2 - P_1 w_1 + (P_1 - P_2)\delta y_1 = 0,$$

$$\text{or } (V_2 - P_2 p_2)\delta x_2 - (V_1 - P_1 w_1)\delta x_1 + P_2 \delta y_2 - P_2 \delta y_1 = 0;$$

$$\therefore \frac{1}{\sqrt{2a}} (\delta x_2 - \delta x_1) + p_2 \delta y_2 - p_2 dy_1 = 0.$$

But if $\frac{dy}{dx} = m$ and $\frac{dy}{dx} = n$, be the equations to the given curves, $\frac{\delta y_1}{\delta x_1} = m$ and $\frac{\delta y_2}{\delta x_2} = n$; $\therefore$ substituting

$$(1 + p_2 n)\,\delta x_2 - (1 + p_2 m)\,\delta x_1 = 0,$$

whence $\because \delta x_2$ and δx_1 are independent variables,

$$1 + p_2 n = 0; \quad 1 + p_2 m = 0; \quad \therefore\ m = n,$$

or the tangents at the points where the cycloid cuts the two curves are parallel; also since $p_2 = -\frac{1}{n}$, it cuts the second curve at right angles.

14. We shall now consider problems in which z is also a function of x, and shall make use of the general method investigated in Art. 5.

Prob. 1. Required the shortest distance between two points in space.

Let s be the distance; $\therefore \int ds = \int \sqrt{dx^2 + dy^2 + dz^2}$;

$$\therefore\ \delta \int ds = \int \delta ds = \int \frac{dx}{ds}\,\delta dx + \int \frac{dy}{ds}\,\delta dy + \int \frac{dz}{ds}\,\delta dz$$

$$= \frac{dx}{ds}\,\delta x + \frac{dy}{ds}\,\delta y + \frac{dz}{ds}\,\delta z - \int \left\{ d \cdot \frac{dx}{ds}\,\delta x + d \cdot \frac{dy}{ds} \cdot \delta y + d \cdot \frac{dz}{ds}\,\delta z \right\},$$

whence $M = 0$, $m = 0$, $M' = 0$;

$$N = \frac{dx}{ds}; \quad n = \frac{dy}{ds}; \quad N' = \frac{dz}{ds}; \quad dN = 0 = dn = dN';$$

$$\therefore \frac{dx}{ds} = a; \quad \frac{dy}{ds} = b; \quad \frac{dz}{ds} = c.$$

Also $\frac{dx^2}{ds^2} + \frac{dy^2}{ds^2} + \frac{dz^2}{ds^2} = 1 = a^2 + b^2 + c^2$;

an equation which connects the constants a, b, c;

also $\because \frac{dx}{dz} = \frac{a}{c}; \quad \frac{dy}{dz} = \frac{b}{c}; \quad \therefore\ x = \frac{a}{c}z + c'; \quad y = \frac{b}{c}z + c''$,

the equations to the projections of a straight line.

Prob. 2. Find the equation to the shortest line that can be drawn upon a given curve surface.

Let $dz = pdx + qdy$ be the equation of the surface; then the variations of the co-ordinates x, y, z which are under the sign $\int$ must satisfy the differential equation; $\therefore \delta z = p\delta x + q\delta y$;

$$\therefore \delta \int ds = \left(\frac{dx}{ds} + p\frac{dz}{ds}\right)\delta x + \left(\frac{dy}{ds} + q \cdot \frac{dz}{ds}\right)\delta y$$

$$- \int\left\{\left(d \cdot \frac{dx}{ds} + p \cdot d \cdot \frac{dz}{ds}\right)\delta x + \left(d \cdot \frac{dy}{ds} + q \cdot d \cdot \frac{dz}{ds}\right)\delta y\right\},$$

whence we have from the part under the sign of integration,

$$d \cdot \frac{dx}{ds} + p \cdot d \cdot \frac{dz}{ds} = 0; \quad \text{and } d \cdot \frac{dy}{ds} + q \cdot d \cdot \frac{dz}{ds} = 0;$$

whence, having found p and q from the equation to the given surface, the equations to the curve may be found.

Ex. 1. Let the surface be a surface of revolution;

$$\therefore z = \phi(x^2 + y^2);$$

$$\therefore p = 2x\phi' \cdot (x^2 + y^2); \quad q = 2y \cdot \phi'(x^2 + y^2);$$

and considering s to be the independent variable,

$$\therefore \frac{d^2x}{ds^2} + p \cdot \frac{d^2z}{ds^2} = \frac{d^2x}{ds^2} + 2x \cdot \phi'(x^2 + y^2)\frac{d^2z}{ds^2} = 0, \quad (1)$$

$$\text{and } \frac{d^2y}{ds^2} + q \cdot \frac{d^2z}{ds^2} = \frac{d^2y}{ds^2} + 2y \cdot \phi'(x^2 + y^2)\frac{d^2z}{ds^2} = 0, \quad (2)$$

$$(1) \times y - (2) \times x; \quad \therefore y\frac{d^2x}{ds^2} - x\frac{d^2y}{ds^2} = 0.$$

$$\text{But if } x^2 + y^2 = r^2, \text{ and } \theta = \cos^{-1}\frac{x}{r};$$

$$y d^2x - x d^2y = d \cdot (r^2 d\theta);$$

$$\therefore \text{integrating, } r^2\frac{d\theta}{ds} = c; \quad \therefore \frac{r d\theta}{ds} = \frac{c}{r}.$$

But $\dfrac{r d\theta}{ds}$ is the sine of the angle at which the shortest line cuts the generating curve or meridian, hence if ϕ be this angle,

$$\sin\phi = \frac{c}{r} = \frac{c}{\sqrt{x^2 + y^2}}.$$

Cor. Since $r^2\dfrac{d\theta}{ds} = c$;

$$\therefore\ r^4\frac{d\theta^2}{dr^2} = c^2\frac{ds^2}{dr^2} = c^2\left(1 + r^2\frac{d\theta^2}{dr^2} + \frac{dz^2}{dr^2}\right);$$

$$\therefore\ r^2(r^2 - c^2)\frac{d\theta^2}{dr^2} = c^2\left\{1 + \phi'(r)^2\right\};$$

$$\therefore\ \frac{d\theta}{dr} = \frac{c}{r}\sqrt{\frac{1 + \phi'(r)^2}{r^2 - c^2}},$$

the equation to the projection on the plane of xy.

Ex. 2. Let the surface be a sphere.

Then if $a\psi$ be the distance of the point of intersection of the shortest line with the meridian, from the point where the axis of z meets the sphere, $r = a\sin\psi$;

$$\therefore\ \sin\phi = \frac{c}{a\sin\psi};\quad \sin\phi\,.\sin\psi = \frac{c}{a},$$

or is constant, which is a property of an arc of a great circle. Hence the shortest line is the arc of a great circle.

Ex. 3. If the surface be a cone,

$$z = e\sqrt{x^2 + y^2} = er;\quad \frac{dz}{dr} = e;$$

$$\therefore\ \frac{d\theta}{dr} = \frac{c}{r}\frac{\sqrt{1 + e^2}}{\sqrt{r^2 - c^2}};$$

$$\therefore\ \theta + c_1 = \sqrt{1 + e^2}\sec^{-1}\left(\frac{r}{c}\right);\quad \therefore\ r = c\sec\left(\frac{\theta + c_1}{\sqrt{1 + e^2}}\right),$$

the equation to the elliptic spiral.

15. Resuming the equations, Art. 14, Prob. 2, since

$$\frac{d^2x}{ds^2} = -p\cdot\frac{d^2z}{ds^2},$$

$$\frac{d^2y}{ds^2} = -q\cdot\frac{d^2z}{ds^2},$$

square, add, and add $\left(\frac{d^2z}{ds^2}\right)^2$;

$$\therefore\ \left(\frac{d^2x}{ds^2}\right)^2 + \left(\frac{d^2y}{ds^2}\right)^2 + \left(\frac{d^2z^2}{ds^2}\right)^2 = (1 + p^2 + q^2)\left(\frac{d^2z}{ds^2}\right)^2;$$

or if R be the radius of absolute curvature, and γ the angle which the normal to the curve makes with the axis of z,

$$\therefore \frac{1}{R^2} = \frac{1}{\cos^2\gamma} \cdot \left(\frac{d^2x}{ds^2}\right)^2; \quad \therefore \pm \frac{d^2x}{ds^2} = \frac{\cos\gamma}{R}.$$

But if $\frac{dx}{ds}$ be the direction cosine of the angle which the tangent makes with x,

$$\therefore \frac{dx}{ds} = \sin\gamma; \quad \frac{d^2x}{ds^2} = \cos\gamma \cdot \frac{d\gamma}{ds} = \pm\frac{\cos\gamma}{R};$$

$$\therefore \frac{1}{R} = \pm\frac{d\gamma}{ds}.$$

Now we know, if θ be the angle which the tangent to a plane curve makes with the axis of x, that $\pm\frac{d\theta}{ds} = \frac{1}{R}$, or that the rate at which the inclination of the tangent increases or decreases, varies as the curvature, hence the preceding result shews that this is also true for the curve of double curvature under consideration, possessing the property of being the minimum distance between any two points on a surface.

Isoperimetrical Problems.

16. The preceding problems have been those of absolute maxima and minima; we now treat of questions of relative maxima and minima. Of this kind is the problem, 'Given the length of a curve, find its equation, when the area included by it is a maximum.' This Problem was first proposed and solved by James Bernoulli, and from its nature was called Isoperimetrical, a name which was extended to all problems of this kind.

The problem of relative maxima and minima may be defined to be this. 'Find $y = f(x)$; so that $\int_x u$ may be a maximum while $\int_x u_1 = c$.' To solve this problem, we multiply $\int_x u_1$ by a constant a, and add the product to $\int_x u$, and make $\delta \cdot \{\int_x u + a\int_x u_1\} = 0$; or $\delta\int_x (u + au_1) = 0$; for since $\int_x u$ is a maximum and $\int_x u_1 = c$, their separate variations will $= 0$; and as the former is limited by the values of the variables in the latter, all the conditions of the problem will be fulfilled and included in the variation of $\int_x (u + au_1)$; so also if there be another equation of condition $\int_x u_2 = c_1$, we must add $b\int_x u_2$ to the former integrals, and take the variation of $\int_x (u + au_1 + bu_2)$.

Hence instead of V, we must write $V + au_1$ or $V + au_1 + bu_2$ in $\delta \int_x V$, and then proceed as in absolute maxima and minima: in these examples the total variation will be expressed by $\delta \int_x V_1$.

PROB. 1. Of all curves of equal perimeters, find that which has the greatest area.

Here $\int_x u_1 = \int_x \sqrt{1+p^2} = c$; $\int_x V = \int_x y$;

$\therefore \int_x V_1 = \int_x (y + a\sqrt{1+p^2})$; $\therefore V_1 = y + a\sqrt{1+p^2}$;

$\therefore dV_1 = dy + \dfrac{ap}{\sqrt{1+p^2}}$; $\therefore N=1$, $P = \dfrac{ap}{\sqrt{1+p^2}}$; $Q = 0$;

$\therefore V_1 = Pp + C$; $\therefore y + a\sqrt{1+p^2} = \dfrac{ap^2}{\sqrt{1+p^2}} + c$;

$\therefore y - c = -\dfrac{a}{\sqrt{1+p^2}}$; $\therefore p = \dfrac{\sqrt{a^2-(y-c)^2}}{y-c}$;

$\therefore \dfrac{dx}{dy} = \dfrac{y-c}{\sqrt{a^2-(y-c)^2}}$; $x - c_1 = -\sqrt{a^2-(y-c)^2}$;

$\therefore (x-c_1)^2 + (y-c)^2 = a^2$. The equation to the circle.

PROB. 2. Find the curve in which a chain of given length l may hang, that its centre of gravity may be the lowest possible.

X = depth of centre of gravity $= \dfrac{\int_x x\sqrt{1+p^2}}{l}$;

$\therefore V_1 = \dfrac{x\sqrt{1+p^2}}{l} + a\sqrt{1+p^2}$;

$\therefore dV_1 = \dfrac{\sqrt{1+p^2}}{l}.dx + \left(\dfrac{xp}{l\sqrt{1+p^2}} + \dfrac{ap}{\sqrt{1+p^2}}\right)dp$;

$\therefore M = \dfrac{\sqrt{1+p^2}}{l}$; $N = 0$, $P = \dfrac{px + lap}{l\sqrt{1+p^2}}$,

$N - \dfrac{dP}{dx} = 0$; $\therefore \dfrac{dP}{dx} = 0$; $\therefore P = c = \dfrac{p.(x+\beta)}{l\sqrt{1+p^2}}$;

$\therefore \dfrac{\sqrt{1+p^2}}{p} = \dfrac{x+\beta}{lc} = \dfrac{x+\beta}{a}$; $\therefore p = \dfrac{a}{\sqrt{(x+\beta)^2 - a^2}}$;

$\therefore \dfrac{y}{a} = \log.\left\{\dfrac{x+\beta+\sqrt{(x+\beta)^2-a^2}}{a_1}\right\}$, the catenary.

PROB. 3. Find the curve, which of all those that have the same length and include the same area, shall by rotation round the axis of x generate the greatest solid.

Here $\int_x V_1 = \int_x (\pi y^2 + a\sqrt{1+p^2} + by)$.

PROB. 4. Find the Brachystochrone when the length of the curve is given;

$$\int_x V_1 = \int_x \left(\frac{\sqrt{1+p^2}}{\sqrt{y}} + a\sqrt{1+p^2} \right).$$

PROB. 5. Of all curves which include a given area, the circle has the least perimeter.

PROB. 6. The curve which by revolution round its axis generates the greatest solid under a given surface, is the circle.

PROB. 7. The length of curve being given, shew that it will generate the least surface, when it is the catenary.

PROB. 8. If from a point two straight lines be drawn, and their extremities be joined by a curve, so that the area included is constant, the curve will be a circular arc, when its length is a minimum.

THE END.

New Works and New Editions*

printed for

John W. Parker,

West Strand.

LONDON: MAY, M.DCCC.XLVI.

A Charge,

Delivered at his Annual Visitation in 1845.

By John George, Lord Archbishop of Armagh.

Octavo, 2*s*.

Religio Quotidiana:

Daily Prayer the Law of God's Church, and heretofore the Practice of Churchmen.

By the Right Rev. Richard Mant, D.D.,

Lord Bishop of Down and Connor, and Dromore.

Foolscap Octavo. Cloth, gilt edges, 4*s*. 6*d*.

The Mission of the Comforter, and other Sermons.

With Notes.

By Julius Charles Hare, M.A.,

Archdeacon of Lewes.

Two Volumes, Octavo.

---

* The Publisher's Catalogue of Educational Books, for the use of Schools and Families, including the Works published under the Sanction of the Committee of Council on Education, and those issued by the Committee of General Literature and Education, appointed by the Society for Promoting Christian Knowledge, may be had, gratis, on application.

Preparing for the Press, New Editions, thoroughly revised and corrected, of The

History and Philosophy of the Inductive Sciences.

By the Rev. William Whewell, D.D., F.R.S., etc.,
Master of Trinity College, and Professor of Moral Philosophy,
in the University of Cambridge.

By the same Author.

Lectures on Systematic Morality,

Delivered in the University of Cambridge, in the Lent Term, 1846.
Octavo. 7*s.* 6*d.*

Elements of Morality, including Polity.

Two Volumes, Octavo, 24*s.*

Indications of the Creator.

Theological Extracts from the *History* and the *Philosophy of the Inductive Sciences.* With a Preface.
Second Edition, with an additional Preface. 5*s.* 6*d.*

Of a Liberal Education in general,

and with particular reference to the leading Studies of the University of Cambridge.
Octavo. 9*s.*

On the Principles of English University Education.

Second Edition. 5*s.*

Architectural Notes on German Churches;

With Notes of a Tour in Normandy and Picardy.
Third Edition, with Additions. 12*s.*

Lectures on the Prophecies,

Proving the Divine Origin of Christianity;

Delivered in the Chapel of Lincoln's Inn.

By Alexander M'Caul, D.D., T.C.D.,

Professor of Divinity in King's College, London, and Prebendary of St. Paul's.

Octavo, 7*s.*

Lectures on the Epistle to the Hebrews;

Delivered in the Chapel of Lincoln's Inn,

By the Rev. F. D. Maurice, M.A.,

Professor of Divinity in King's College, and Chaplain of Lincoln's Inn.

With a Preface, containing a Review of Mr. Newman's Theory of Development.

Octavo, 7*s.* 6*d.*

Notes on the Miracles.

Octavo, 12*s.*

Exposition of the Sermon on the Mount,

Drawn from the Writings of St. Augustine, with Observations.

Foolscap Octavo, 3*s.* 6*d.*

By the Rev. Richard Chenevix Trench, M.A.,

Professor of Divinity in King's College, London, and Examining Chaplain to the Lord Bishop of Oxford.

Sermons preached at Jerusalem, in the years 1842 and 1843.

By the Rev. George Williams, M.A.,

Fellow of King's College, Cambridge; sometime Chaplain to the late Bishop Alexander.

Octavo, 10*s.* 6*d.*

Theory of Developments in Christian Doctrine, applied and tested.

By Mortimer O'Sullivan, D.D.,

Rector of Kellyman.

Dedicated, by permission, to His Grace the Archbishop of Armagh.

Octavo. 7*s*. 6*d*.

Bishop Jeremy Taylor;

His Predecessors, Contemporaries, and Successors.

By the Rev. Robert Aris Willmott, M.A.

Incumbent of St. Catherine's, Bear Wood; Author of *Lives of Sacred Poets*.

Foolscap Octavo.

Bishop Heber and His Works;

With some Account of Christian Missions in India.

By the Rev. James Chambers, M.A.,

late of All Souls' College, Oxford.

Foolscap Octavo. 2*s*. 6*d*.

Life of Archbishop Sancroft.

By the late George D'Oyly, D.D., F.R.S.,

Rector of Lambeth.

The New Edition, revised. Octavo. 9*s*

Modern Hagiology:

An Examination of the Nature and Tendency of some Legendary and Devotional Works lately published under the Sanction of the Rev. J. H. Newman, the Rev. Dr. Pusey, and the Rev. F. Oakeley.

By the Rev. J. C. Crosthwaite, M.A.,

Rector of St. Mary-at-Hill and St. Andrew Hubbard, London.

Two Volumes, foolscap Octavo. 9*s*.

The Gospel-Narrative,

according to the Authorized Text of the Evangelists, without Repetition or Omission. With a Continuous Exposition, Marginal Proofs in full, and Notes, briefly collected from the best Critics and Commentators.

Dedicated, by Special Permission, to Her Majesty the Queen.

Second Edition, Super Royal Octavo. 16*s.*, boards.

II.

The Churchman's Guide;

A Copious Index of Sermons and other Works, by eminent Church of England Divines; arranged according to their subjects.

Octavo. 7*s.*

By the Rev. John Forster, M.A.,
Her Majesty's Chaplain of the Savoy.

Two Series of Discourses,

I. *On Christian Humiliation.* II. *On the City of God.*

By the Right Rev. C. H. Terrot, D.D.,
Bishop of Edinburgh; late Fellow of Trinity College, Cambridge.

Octavo. 7*s.* 6*d.*

Practical Sermons

By Dignitaries and other Clergymen of the United Church of England and Ireland.

Printed verbatim from the Authors' MSS.

Edited by the Rev. J. C. Crosthwaite, M.A.,
Rector of St. Mary-at-Hill, and St. Andrew Hubbard, London.

Complete in Three Volumes, Octavo, 7*s.*, in cloth; or in Eighteen Monthly Parts.

Sacra Domestica:

A Course of Family Prayers.

In Octavo; Large Type, with provision for Additions in Manuscript. 3*s.*

College Lectures on Christian Antiquities, and the Ritual of the English Church.

II.

College Lectures on Ecclesiastical History.

Each with complete Sets of Cambridge, Dublin, and Durham University Examination Papers.

By the Rev. W. Bates, M.A.,

Fellow, Lecturer, and Hebrew Lecturer of Christ's College, Cambridge.

Post Octavo, 9*s*. each.

A Manual of Christian Antiquities;

An Account of the Constitution, Ministers, Worship, Discipline, and Customs of the Early Church: with a Complete Analysis of the Works of the Antenicene Fathers.

By the Rev. J. E. Riddle, M.A.

The Second Edition, Octavo. 18*s*.

Ordo Sæclorum;

A Treatise on the Chronology of the Holy Scriptures.

By the Rev. Henry Browne, M.A.,

Canon of Chichester, and Chaplain to the Lord Bishop.

In a large Volume, Octavo. 20*s*.

JAMES'S TREATISE

On the Corruption of Scripture, Councils, and Fathers,

by the Prelates, Pastors, and Pillars of the Church of Rome, for the maintenance of Popery.

Revised and Corrected from the Editions of 1612 and 1688,

By the Rev. J. E. Cox, M.A.

Octavo, 12*s*.

The Anglo-Saxon Church;

Its History, Revenues, and General Character.

By the Rev. Henry Soames, M.A.

The Third Edition, with Additions and Corrections. 10*s.* 6*d.*

History of the Christian Church.

From the Ascension of Jesus Christ to the Conversion of Constantine.

By the late Professor Burton.

The Sixth Edition. 6*s.* 6*d.*

History of the Church of England.

By the Right Rev. T. Vowler Short, D.D.,

Lord Bishop of Sodor and Man.

The Fourth Edition, Revised. 16*s.*

History of the Church of Ireland.

By the Right Rev. Richard Mant, D.D.,

Lord Bishop of Down and Connor, and Dromore.

Two large Volumes, to be had together or separately, 17*s.* each.

The Choral Service of the Church;

Being an Enquiry into the Liturgical System of the Cathedral and Collegiate Foundations of the Anglican Communion.

By the Rev. John Jebb, M.A.,

Rector of Peterstow.

Octavo. 16*s.*

The Literature of the Church of England,

exhibited in Specimens of the Writings of Eminent Divines, with Memoirs of their Lives, and Historical Sketches of the Times in which they lived.

By the Rev. Richard Cattermole, B.D.

Two Volumes, Octavo. 25*s.*

The Statutes relating to the Ecclesiastical & Eleemosynary Institutions

of England, Wales, Ireland, India, and the Colonies;
with the Decisions thereon.

By Archibald J. Stephens, M.A., F.R.S.
Barrister at Law.

Two Volumes, Royal Octavo. 3*l.* 3*s.* boards, or 3*l.* 13*s.* 6*d.* in law calf.

Horæ Ecclesiasticæ;

The Position of the Church with regard to Romish Error.

II.

Horæ Liturgicæ;

I. *Liturgical Discrepancy.* II. *Liturgical Harmony.*

By the Right Rev. Richard Mant, D.D.,
Lord Bishop of Down and Connor, and Dromore.
Printed uniformly, Foolscap Octavo, 3*s.* 6*d.* each.

Specimens of Ecclesiastical Architecture in Great Britain,

Edited by Henry Bowman, Architect.

Illustrated by Views, Plans, Elevations, Sections, and Det il.
Royal Quarto, 2*l.* 5*s* ; or, in Twelve Parts, at 3*s.* 6*d* each

---

The Churchman's Theological Dictionary.

By the Rev. Robert Eden, M.A., F.S.A.;
late Fellow of Corpus Christi College, Oxford.

The design of this Work is to give plain and simple explanations of the Theological and Ecclesiastical Terms which are used in describing and discussing religious Ordinances, Doctrines, and Institutions, without entering into the controversies which have arisen respecting their object and import.

Second Edition, Medium 12mo. 5*s.*

Biographia Britannica Literaria:

A Literary History of the United Kingdom.

By T. Wright, M.A., Corresponding Member of the Institute of France.

Published under the Superintendence of the ROYAL SOCIETY OF LITERATURE.

VOL. I. THE ANGLO-SAXON PERIOD.
VOL. II. THE ANGLO-NORMAN PERIOD.

To be continued. Octavo. 12*s.* each.

The Holy City;

Historical and Topographical Notices of Jerusalem;

By the Rev. George Williams, M.A.,

Fellow of King's College, Cambridge; and sometime Chaplain to the late Bishop Alexander, at Jerusalem.

With numerous Illustrations. Octavo, 18*s.*

Travels and Researches among the Christian Tribes

In Asia Minor, Mesopotamia, Chaldea, and Armenia;

By W. F. Ainsworth, F.G.S., F.R.G.S.,

Two Volumes, with Maps and numerous Illustrations. 24*s.*

Lusitanian Sketches of the Pen and Pencil;

By William H. G. Kingston, Esq.

Two Volumes, Post Octavo. 18*s.*

Exercises for the Cultivation of the Voice;

By John Hullah.

I. For Soprano and Tenor. II. For Contralto and Bass Voices.

Two separate Volumes. Imperial Octavo, 4*s.* 6*d.* each.

A System of Logic;

By John Stuart Mill.

New Edition, Two Volumes. Octavo. 30*s.*

Creation by the Agency of God, as opposed to Creation by Natural Law;

being a Refutation of the Work entitled *Vestiges of the Natural History of Creation.*

By Thomas Monck Mason, B.A.

Post Octavo. 5*s.*

A Cycle of Celestial Objects;

By Captain W. H. Smyth, R.N.; K.S.F., D.C.L., F.R.S.
President of the Astronomical Society.

Two Volumes, Octavo; with numerous Illustrations, 2*l.* 2*s.*

Elements of Meteorology;

By the late J. F. Daniell, D.C.L., For. Sec. R.S.;
Professor of Chemistry in King's College, London.

Two Volumes, Octavo, with Coloured Charts and other Plates. 32*s.*

Practical Geodesy;

comprising Chain Surveying, the Use of Surveying Instruments, together with Levelling, and Trigonometrical, Mining and Maritime Surveying.

By Butler Williams, C.E., F.G.S.,
late Professor of Geodesy in the College of Civil Engineers.

Octavo, with numerous Illustrations. 12*s.* 6*d.*

Second Edition, with new Chapters on Railway, Parish, and Estate Surveying.

On the Nature of Thunderstorms,

and on the Means of Protecting Churches and other Buildings, and Shipping, against the Destructive Effects of Lightning.

By W. Snow Harris, F.R.S.

Octavo. 10*s.* 6*d.*

Lectures on Astronomy,

delivered at King's College, London.

By the Rev. H. Moseley, M.A., F.R.S.

late Professor of Astronomy and Natural Philosophy in the College.

The Second Edition, revised, with numerous Wood Cuts. 5*s.* 6*d.*

Practical Geology and Mineralogy & the Chemistry of Metals.

By Joshua Trimmer, F.G.S.

Octavo, with Two Hundred Illustrations. 12*s.*

Lectures on the Principles and Practice of Physic.

By Thomas Watson, M.D.,

Fellow of the Royal College of Physicians; late Physician to the Middlesex Hospital; and formerly Fellow of St. John's College, Cambridge.

New Edition. Two Volumes, Octavo, 34*s.*

Physiological Anatomy and Physiology of Man.

By R. B. Todd, M.D., F.R.S., & W. Bowman, F.R.S.,

of King's College, London.

With numerous Original Illustrations.

The First Volume, Octavo, 15*s.*, cloth; also Parts I. and II. 7*s.*, each.

To be completed in Four Parts, forming Two Volumes.

An Introduction to the Study of Chemical Philosophy;

By the late Professor Daniell, F.R.S.

With numerous Illustrations. Second Edition, much Enlarged, 21*s*.

A Manual of Chemistry.

The Fifth Edition, Revised and Considerably Enlarged, with numerous Wood-Cuts, 35*s*.

II.

A Dictionary of the Materia Medica and Pharmacy.

Octavo. 15*s*.

By William Thomas Brande, F.R.S.,

Of Her Majesty's Mint; Professor of Chemistry at the Royal Institution

Practical Chemistry for Farmers and Landowners.

By Joshua Trimmer, F.G.S.,

Post Octavo. 5*s*.

The Practical Bee-keeper;

Or, Concise and Plain Instructions for the Management of Bees and Hives.

By John Milton.

Foolscap Octavo. 4*s*. 6*d*.

The Health and Sickness of Town Populations,

considered with reference to proposed Sanatory Legislation, and to the establishment of a comprehensive system of Medical Police and District Dispensaries. With Appendices and Statistical Tables.

Octavo. 3*s*. 6*d*.

The Fables of Babrius,

From the newly-discovered Manuscript; together with the Fragments of the lost Fables.

Edited by George Cornewall Lewis, Esq., M.A.,

late Student of Christ Church.

Post Octavo. *5s. 6d.*

Uniformly, in Post Octavo, with Illustrations, 12*s.* each work.

Becker's Charicles;

Illustrations of the Private Life of the Ancient Greeks.

II.

Becker's Gallus;

Roman Scenes of the Time of Augustus.

With Notes and Excursus.

Translated by the Rev. F. Metcalfe, M.A.,

Fellow of Lincoln College, Oxford.

Dahlmann's Life of Herodotus, drawn out from his Book.

Translated, with Notes, by G. V. Cox, M.A.,

Esquire Bedell in the University of Oxford.

Post Octavo. *5s.*

Travels in the Track of the Ten Thousand Greeks;

A Geographical and Descriptive Account of the Expedition of Cyrus and of the Retreat of the Ten Thousand, as related by Xenophon.

By W. F. Ainsworth, F.G.S.,

Surgeon to the Euphrates Expedition.

Post Octavo. *7s. 6d.*

A Selection from the Greek Verses of Shrewsbury School;

With an Account of the Iambic Metre and Style of Greek Tragedy, and Progressive Exercises in Greek Tragic Scenarii.

By B. H. Kennedy, D.D.,

Head Master of Shrewsbury School.

Second Edition, enlarged. *In the Press.*

Iliad of Homer: Books I. to III.

With the English Notes and Homeric Glossary of Professor ANTHON.

Edited, with his Sanction, by J. R. Major, D.D.,

Head Master of King's College School, London.

Nearly Ready.

The Æneid of Virgil, with the English Notes of Professor Anthon.

Edited, with his Sanction, by J. R. Major, D.D.,

Head Master of King's College School, London.

Foolscap Octavo. 7*s.* 6*d.*, strongly bound.

The Student's Manual of Ancient History;

Containing the Political History, Geographical Position, and Social State of the Principal Nations of Antiquity.

Fourth Edition, 10*s.* 6*d.*

II.

The Student's Manual of Modern History;

The Rise and Progress of the Principal European Nations, their Political History, and the Changes in their Social Condition; with a History of the Colonies founded by Europeans.

Third Edition, 10*s.* 6*d.*

By W. C. Taylor, LL.D.

Outlines of the History of France,

for Schools and Families.

By the Rev. O. Cockayne,

King's College, London.

Uniformly with the *Outlines of English, Roman,* and *Grecian History.* 1*s.* 3*d.*

Amusements in Chess.

I. The History, Antiquities and Curiosities of the Game. II. Easy Lessons in Chess, illustrative of the various Openings, analysed and explained. III. Chess Problems, or, Ends of Games won or drawn by brilliant and scientific Moves.

By Charles Tomlinson.

With numerous Illustrations. 4*s.* 6*d.*

The Merchant and the Friar;

or, Truths and Fictions of the Middle Ages.

By Sir Francis Palgrave.

New Edition. 3*s.*

A Sketch of the Military History of Great Britain.

By the Rev. G. R. Gleig, M.A.,

Chaplain-General to the Forces, and Chaplain of Chelsea Hospital.

Foolscap Octavo. 3*s.* 6*d.*

The Life and Services of General Lord Harris, G.C.B.

By the Right Hon. S. R. Lushington,

Private Secretary to Lord Harris, and late Governor of Madras.

A new Edition, revised, Foolscap Octavo. 6*s.* 6*d.*

Works Edited by John Hullah,

Professor of Vocal Music in King's College, London.

Vocal Scores;

Sacred and Secular Music, for four or more Voices.

In Monthly Numbers, containing 16 Folio pages, 1*s.* each.

Part Music.

Class A, in Score and in Parts, for Soprano, Alto, Tenor, and Bass Voices. Two Volumes Sacred and Two Secular. Cloth, Score, 9*s.*; Voice Parts, 3*s.* each. Also in Numbers (I. to XII.)—Score, 2*s.* 6*d.*; Voice Parts, 8*d.* each.

Class B.—In Score for the Voices of Women and Children. One Volume Sacred and one Secular. 3*s.* each. Also in Numbers (I. to VI.) 8*d* each.

Class C.—In Score for the Voices of Men. One Volume Sacred and One Secular. 3*s.* each. Also in Numbers (I. to VI.) 8*d.* each.

The Whole Book of Psalms,

With the Canticles and Hymns of the Church, for the Morning and Evening Service,

Set to appropriate Chants for four Voices,

Every Syllable being placed under its proper Note.

Imperial Octavo, 15*s.*, bound in cloth.

The Psalter;

Or, Psalms of David, in Metre, set to appropriate Tunes, arranged for four Voices.

The various Editions comprise the arrangements in Score, with and without Organ or Piano-forte Accompaniments, and the different Voice Parts, each in separate volumes. They may be had in fine Editions, in plain or elegant bindings, or cheap Editions, ornamentally or strongly bound, as may be required. For particulars and prices, see the Publisher's Catalogues.

Psalm Tune-Books,

Without Words, containing eighty-two Tunes from the Psalter.

The Tunes harmonized for Three Equal Voices, 2*s.* 6*d.*

The Tunes harmonized for four Voices, 3*s.*

London: John W. Parker, West Strand.